

The First Naturally Occurring Thiepinols and Thienol from an Endolichenic Fungus *Coniochaeta* sp.


Yinchao Wang,^{†,§} Shubin Niu,[†] Shuchun Liu,[†] Liangdong Guo,[†] and Yongsheng Che^{*,†,‡}

Key Laboratory of Systematic Mycology & Lichenology, Institute of Microbiology, Chinese Academy of Sciences, Beijing 100190, People's Republic of China, Beijing Institute of Pharmacology & Toxicology, Beijing 100850, People's Republic of China, and Graduate School of Chinese Academy of Sciences, Beijing 100039, People's Republic of China

cheys@im.ac.cn

Received September 10, 2010

ABSTRACT



Coniothiepinols A (1) and B (2) and coniothienol A (3), the first naturally occurring thiepinols (1 and 2) and thienol (3), have been isolated from the crude extract of an endolichenic fungus *Coniochaeta* sp. 1 possesses a unique 8-oxa-2-thia-bicyclo[3.2.1]octane skeleton, and its structure was assigned by NMR spectroscopy and X-ray crystallography. 1 showed significant activity against the Gram-positive bacteria, *Enterococcus faecium* and *Enterococcus faecalis*.

Analogous to plant endophytes living in the intercellular spaces of the hosts, endolichenic fungi are microbes that inhabit the thalli of lichens.¹ To date, only a limited number of secondary metabolites have been reported from the endolichenic fungi. Examples include five heptaketides isolated from the *Corynespora* sp.,^{2,3} ambuic acid and torreyanic acid derivatives from the *Pestalotiopsis* sp.,⁴ and allenyl and alkynyl phenyl ethers from *Neurospora terricola*.⁵ Our prior chemical study of the endolichenic fungus *Coniochaeta* sp. also afforded six new xanthone derivatives, such

as conioxepinol A (4), a cytotoxic oxepinochromenone, and coniofuro A (5), a furochromenone.⁶ The oxepinochromenones and furochromenones (ring-expanded and ring-contracted xanthenes, respectively) are relatively rare, with only a few precedents reported prior to our work.^{7–11}

Since the crude extract of *Coniochaeta* sp. also showed antimicrobial activities, and its HPLC chromatogram revealed minor components that could not be identified, the fungus was

* To whom correspondence should be addressed. Tel/Fax: +86 10 82618785.

[†] Institute of Microbiology.

[‡] Beijing Institute of Pharmacology & Toxicology.

[§] Graduate School of Chinese Academy of Sciences.

(1) Arnold, A. E. *Fungal Biol. Rev.* **2007**, *21*, 51–66.

(2) Paranagama, P. A.; Wijeratne, E. M. K.; Burns, A. M.; Marron, M. T.; Gunatilaka, M. K.; Arnold, A. E.; Gunatilaka, A. A. L. *J. Nat. Prod.* **2007**, *70*, 1700–1705.

(3) Wijeratne, E. M. K.; Bashyal, B. P.; Gunatilaka, M. K.; Arnold, A. E.; Gunatilaka, A. A. L. *J. Nat. Prod.* **2010**, *73*, 1156–1159.

(4) Ding, G.; Li, Y.; Fu, S.; Liu, S.; Wei, J.; Che, Y. *J. Nat. Prod.* **2009**, *72*, 182–186.

(5) Zhang, F.; Liu, S.; Lu, X.; Guo, L.; Zhang, H.; Che, Y. *J. Nat. Prod.* **2009**, *72*, 1782–1785.

(6) Wang, Y.; Zheng, Z.; Liu, S.; Zhang, H.; Guo, L.; Che, Y. *J. Nat. Prod.* **2010**, *73*, 920–924.

(7) Singh, S. B.; Ball, R. G.; Zink, D. L.; Monaghan, R. L.; Polishook, J. D.; Sanchez, M.; Pelaez, F.; Silverman, K. C.; Lingham, R. B. *J. Org. Chem.* **1997**, *62*, 7485–7488.


(8) Bugni, T. S.; Bernan, V. S.; Greenstein, M.; Janso, J. E.; Maiese, W. M.; Mayne, C. L.; Ireland, C. M. *J. Org. Chem.* **2003**, *68*, 2014–2017.

(9) Liermann, J. C.; Kolshorn, H.; Opatz, T.; Thines, E.; Anke, H. *J. Nat. Prod.* **2009**, *72*, 1905–1907.

(10) Krohn, K.; Kouam, S. F.; Kuigoua, G. M.; Hussain, H.; Cludius-Brandt, S.; Flörke, U.; Kurtán, T.; Pescitelli, G.; Di Bari, L.; Draeger, S.; Schulz, B. *Chem.—Eur. J.* **2009**, *15*, 12121–12132.

(11) Motai, T.; Kitanaka, S. *J. Nat. Prod.* **2005**, *68*, 1732–1735.

referred to on a larger scale on rice in which the oxepinochromenones and furochromenones were initially isolated. Bioassay-guided separation of an EtOAc extract afforded two thiopinols, coniothiopinols A (**1**) and B (**2**), and a thienol, coniothienol A (**3**). Details of their structure assignment and antimicrobial activities are reported herein.



Coniothiopinol A (**1**) was assigned a molecular formula of $C_{16}H_{14}O_7S$ (10 degrees of unsaturation) by HRESIMS (m/z 373.0353 $[M + Na]^+$). Its NMR spectra showed resonances for two exchangeable protons, two methyl groups (one methoxy), one methylene, two oxymethines, eight aromatic/olefinic carbons with two protonated, one oxygenated sp^3 quaternary carbon, one carboxylic carbon (δ_C 166.2), and one α,β -unsaturated ketone carbon (δ_C 177.2). The 1H and ^{13}C NMR data of **1** (Table 1)

Table 1. NMR Spectroscopic Data for **1** in Acetone- d_6

position	δ_H^a (J in Hz)	δ_C^b	HMBC (H \rightarrow C#)
1		161.5	
2	6.61, s	113.0	1, 4, 9a, 11
3		148.1	
4	6.76, s	107.8	2, 4a, 9, 9a, 11
4a		157.4	
5		96.9	
6	5.06, m	84.6	
7a	2.41, dd (8.0, 3.5)	46.3	6, 8, 8a
7b	2.75, dd (13.5, 8.0)		5, 8a
8	5.79, d (8.0)	73.0	5, 6, 8a, 9, 10a
8a		117.1	
9		177.2	
9a		108.6	
10a		165.4	
11	2.38, s	22.1	2, 3, 4
12		166.2	
13	3.88, s	53.4	12
OH-1	12.29, s		1, 2, 3
OH-6	5.42, d (7.0)		

^a Recorded at 500 MHz. ^b Recorded at 100 MHz.

revealed the same 5-hydroxy-7-methyl-4*H*-chromen-4-one unit as found in **4** and **5**,⁶ but the remaining portion was significantly different. The 1H - 1H COSY NMR data of **1** showed the isolated spin-system of C-6–C-8 (including OH-6). HMBC correlations

from H₂-7 and H-8 to C-8a, and from H-7b to C-5 led to the connections of C-8 to C-8a and C-5 to C-6, respectively. While that from H-8 to C-5 established an ether linkage between C-5 and C-8. Considering the chemical shifts of C-5 (δ_C 96.9) and C-10a (δ_C 165.4), the only sulfur atom in **1** was attached to both carbons to complete a 4,5-dihydro-2*H*-thiopyno[2,3-*b*]chromen-6(3*H*)-one skeleton. An HMBC cross peak from H₃-13 to C-12 connected the C-13 *O*-methyl group to C-12, whereas C-12 was attached to C-5 on the basis of unsaturation requirement, permitting assignment of the planar structure of **1** as shown.

Finally, **1** was further confirmed by single-crystal X-ray diffraction analysis (Figure 1), and the X-ray data allowed




Figure 1. Thermal ellipsoid representation of **1**. (Note: The numbering of structure **1** presented here is consistent with the backbone numbering for **1**. A different numbering system is used for the structural data deposited with the CCDC.)

determination of its relative configuration. The presence of a sulfur atom in **1** and the value of the Flack parameter 0.01(10)¹² determined by X-ray analysis also permitted assignment of the absolute configurations of all the chiral centers as 5*R*, 6*R*, and 8*S*.


Compound **2** was given a molecular formula of $C_{16}H_{16}O_6S$ by HRESIMS (m/z 359.0563 $[M + Na]^+$). Analysis of its NMR spectroscopic data showed structural similarity to **1**, except that the thiepane ring was different. Specifically, the C-8 oxymethine in **1** (δ_H/δ_C 5.79/73.0) was reduced and connected to the methyl formate unit as evidenced by its NMR shifts (δ_H/δ_C 3.93/44.6) and HMBC cross peaks from H-8 and H₃-13 to C-12. While the C-7 methylene in **1** was replaced by an oxymethine (δ_H/δ_C 4.25/66.9), and the C-5 oxygenated sp^3 quaternary carbon was replaced by a methylene (δ_H/δ_C 2.90/26.8), which were supported by relevant 1H - 1H COSY NMR data. Therefore, the gross structure of **2** was determined as depicted.

The relative configuration of **2** was deduced by analogy to **4**.⁶ Considering their biogenetic similarity, the C-7 and C-8 stereogenic centers in both compounds presumably have the same configuration, suggesting a *cis* relationship between OH-7 and the methyl formate group, which was partially supported by a NOESY correlation of OH-7 with H₃-13.

The absolute configuration of the C-7 secondary alcohol in **2** was first assigned via the circular dichroism data of an in situ

(12) Flack, H. D. *Acta Crystallogr., Sect. A* **1983**, 39, 876–881.

configuration. Considering the possible interference of the carbonyl functionality, the modified Mosher method was also applied.^{15,16} Treatment of **2** with (S)- and (R)-MTPA Cl afforded *R*-(**2a**) and *S*-MTPA (**2b**) monoesters, respectively. The difference in chemical shift values ($\Delta\delta = \delta_S - \delta_R$) for **2b** and **2a** was calculated to assign the 7*S* configuration (Figure 3). Therefore, the 7*S* and 8*R* absolute configuration



Compounds **1–3** were tested for activity against the Gram-positive bacteria, *Enterococcus faecium* (CGMCC 1.2025) and *Enterococcus faecalis* (CGMCC 1.2535), and the plant pathogenic fungus *Fusarium oxysporum* (CGMCC 3.2830) (Table 2). Com-

compd	IC ₅₀ (μg/mL)		
	<i>E. faecium</i>	<i>E. faecalis</i>	<i>F. oxysporum</i>
1	3.93 ± 0.18	11.51 ± 0.45	13.12 ± 0.46
2	>20	>20	>20
3	2.00 ± 0.06	4.89 ± 0.19	>20
ampicillin	0.51 ± 0.014	2.61 ± 0.23	
carbendazim			0.44 ± 0.008

OL102168Z

(16) Ohtani, I.; Kusumi, T.; Kashman, Y.; Kakisawa, H. *J. Am. Chem. Soc.* **1991**, *113*, 4092–4096.