

Evolutionary Comparisons of the Chloroplast Genome in Lauraceae and Insights into Loss Events in the Magnoliids

Yu Song^{1,2,†}, Wen-Bin Yu^{1,2,†}, Yunhong Tan^{1,2,†}, Bing Liu³, Xin Yao¹, Jianjun Jin⁴, Michael Padmanaba¹, Jun-Bo Yang^{4,*}, and Richard T. Corlett^{1,2,*}

¹Center for Integrative Conservation, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Mengla, China

²Southeast Asia Biodiversity Research Institute, Chinese Academy of Sciences, Yezin, Nay Pyi Taw, Myanmar

³State Key Laboratory of Systematic and Evolutionary Botany, Institute of Botany, Chinese Academy of Sciences, Beijing, China

⁴Germplasm Bank of Wild Species, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, China

*Corresponding authors: E-mails: corlett@xtbg.org.cn; jbyang@mail.kib.ac.cn.

†These authors contributed equally to this work.

Accepted: September 1, 2017

Data deposition: This project has been deposited at GenBank of NCBI under the accession number MF939337 to MF939351.

Abstract

Available plastomes of the Lauraceae show similar structure and varied size, but there has been no systematic comparison across the family. In order to understand the variation in plastome size and structure in the Lauraceae and related families of magnoliids, we here compare 47 plastomes, 15 newly sequenced, from 27 representative genera. We reveal that the two shortest plastomes are in the parasitic Lauraceae genus *Cassytha*, with lengths of 114,623 (*C. filiformis*) and 114,963 bp (*C. capillaris*), and that they have lost NADH dehydrogenase (*ndh*) genes in the large single-copy region and one entire copy of the inverted repeat (IR) region. The plastomes of the core Lauraceae group, with lengths from 150,749 bp (*Nectandra angustifolia*) to 152,739 bp (*Actinodaphne trichocarpa*), have lost *trnI*-CAU, *rpl23*, *rpl2*, a fragment of *ycf2*, and their intergenic regions in IRb region, whereas the plastomes of the basal Lauraceae group, with lengths from 157,577 bp (*Eusideroxylon zwageri*) to 158,530 bp (*Beilschmiedia tungfangensis*), have lost *rpl2* in IRa region. The plastomes of *Calycanthus* (Calycanthaceae, Laurales) have lost *rpl2* in IRb region, but the plastome of *Caryodaphnopsis henryi* (Lauraceae) remain intact, as do those of the nonLaurales magnoliid genera *Piper*, *Liriodendron*, and *Magnolia*. On the basis of our phylogenetic analysis and structural comparisons, different loss events occurred in different lineages of the Laurales, and fragment loss events in the IR regions have largely driven the contraction of the plastome in the Lauraceae. These results provide new insights into the evolution of the Lauraceae as well as the magnoliids as a whole.

Key words: Lauraceae, chloroplast, genome, phylogenetic relationship, loss event.

Introduction

In land plants, most chloroplast genomes are single, circular, double-stranded DNA sequences 100–220 kb in size, with a quadripartite structure including one large single-copy (LSC) region, one small single-copy (SSC) region, and a pair of inverted repeat (IR) regions (Bock 2007). Together these regions include >30 structural RNA genes and around 80 protein-coding genes, with the latter including genes related to photosynthesis, transcription or translation, and other functions (Gao et al. 2010). Generally, the ribosomal RNA genes are in the IR region, almost all of the photosynthesis related

genes in the LSC region, and a number of the NADPH dehydrogenase genes in the SSC region. The plastomes of land plants originated once, from a free-living algal ancestor (Turmel et al. 2006), but the gene contents and order vary considerably among species, and significant structural rearrangements and gene losses have been reported in several unrelated lineages, including ferns (Roper et al. 2007; Karol et al. 2010), gnetophytes (McCoy et al. 2008; Wu et al. 2009), and multiple angiosperm families (Goremykin et al. 2003a; Cai et al. 2006), as well as nonphotosynthetic plants (Wicke et al. 2016).

© The Author 2017. Published by Oxford University Press on behalf of the Society for Molecular Biology and Evolution.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

Comparative analyses of the plastomes of algae and embryophytes show that four genes, *tufA*, *ftsH*, *odpB*, and *rpl5*, have been lost or transferred to the nucleus and three genes, *matK*, *ycf1*, and *ycf2*, have been gained in charophyte algae and embryophytes (Turmel et al. 2006). For example, the *tufA* gene, encoding chloroplast protein synthesis elongation factor Tu, is encoded in the plastomes of most algae, but is a pseudogene in *Isoetes*, fragmented in *Anthoceros*, cycads, and *Ginkgo*, and completely lost in the angiosperms (Karol et al. 2010). Within the angiosperms, three genes, *ycf1*, *ycf2*, and *accD*, have been lost in the Poaceae (Guisinger et al. 2010), whereas *rpl22*, *infA*, and *accD* were lost in the legumes, Lemnoideae, and Acoraceae, respectively (Wang and Messing 2011; Goremykin et al. 2005; Doyle et al. 1995). In plants with a heterotrophic lifestyle, pseudogenization and entire loss events of *ndh*-genes were detected (Wickett et al. 2008; Barrett et al. 2014; Wicke et al. 2016). However, the *ndh*-gene loss events have also occurred in autotrophic orchids, gnetophytes, and Pinaceae (Braukmann et al. 2009; Kim et al. 2015; Wakasugi et al. 1994).

In addition to gene losses, large inversions, and other structural rearrangements have been also reported. In ferns and seed plants, a 30-kb fragment flanked by the complete *matK* and *rpoC2* has been identified as an inversion, with gene organization different from that in liverworts, mosses, hornworts, lycophytes, and *Chaetosphaeridium* (Wickett et al. 2011). In rice, maize, *Calamus*, and orchids, two identical *trnH-rps19* gene clusters were detected as a duplication event before the diversification of extant monocot lineages (Chang et al. 2006; Wang et al. 2008; Luo et al. 2016). In *Tetracentron* and *Trochodendron*, a 4-kb extra region containing the five genes *rpl22*, *rps3*, *rpl16*, *rpl14*, and *rps8* was found as evidence for unstable boundaries of the IR region across early-diverging eudicots (Sun et al. 2013, 2016). Interestingly, most of the rearrangements were detected in the boundary regions of IR, suggesting that the IR regions represent hotspots for structural rearrangements within the plastome (Wicke et al. 2011; Zhu et al. 2016).

The IR regions in the plastome of angiosperms have been used as evolutionary markers for elucidating relationships among some taxa, because they are frequently subject to contraction, expansion, or even complete loss (Lavin et al. 1990; Kim and Jansen 1994; Plunkett and Downie 2000; Luo et al. 2016; Sun et al. 2016; Zhu et al. 2016). In the early-diverging eudicots, the IR regions range from 24.3 to 36.4 kb in length and contain from 18 to 33 genes (Sun et al. 2016). In early-diverging monocots, the IR regions range from 25.2 to 33.3 kb in length and contain from 16 to 20 genes (Luo et al. 2016). As extreme examples, loss of one or two IR regions has been detected in Cephalotaxaceae (Yi et al. 2013), Pinaceae (Wu et al. 2011b), Taxodiaceae (Hirao et al. 2008), Leguminosae (Palmer et al. 1987; Lavin et al. 1990), Geraniaceae (Guisinger et al. 2011), and Cactaceae (Sanderson et al. 2015).

After the eudicots and monocots, the magnoliids is the third-largest group of Mesangiospermae, and includes four orders, 19 families, and over 9,000 woody species from all over the world (www.theplantlist.org). However, <30 species have assembled chloroplast genome sequences, and there has not been a systemic structural comparison of these plastomes. To improve understanding of the dynamics and evolution of plastome structure in magnoliids, we therefore focused on the plastomes of the important family Lauraceae and the related families Calycanthaceae (Laurales), Chloranthaceae (Chloranthales), Magnoliaceae (Magnoliales), Piperaceae (Piperales), and Winteraceae (Canellales). We included 15 newly sequenced and 33 previously reported plastomes in our study, representing 25 genera from all four orders of magnoliids. The main objectives of this study were 1) to reconstruct the phylogenetic relationships using the sequenced magnoliid plastomes, 2) to reveal plastome structural variations in Lauraceae, 3) to trace the evolutionary pattern of plastome contraction.

Materials and Methods

Plant Material and Plastome Sequencing

Fresh leaves and silica-gel dried materials were sampled from 15 species representing 10 genera of Lauraceae. The voucher specimens for the 15 sampled plants collected from China and Indonesia were deposited at the Herbarium of Xishuangbanna Tropical Botanical Garden (HITBC), Chinese Academy of Sciences (CAS; [table 1](#)). Genomic DNA was extracted from 2 g leaves using the CTAB method (Doyle and Dickson 1987), in which 4% CTAB was used, and we added ~1% polyvinyl polypyrrolidone (PVP) and 0.2% DL-dithiothreitol (DTT). From each purified sample of total DNA, 0.5 µg was fragmented to construct short-insert (500 bp) libraries following the manufacturer's manual (Illumina) and then used for sequencing. The DNA samples were indexed by tags and pooled together in one lane of a Genome Analyzer (Illumina HiSeq 2000) for sequencing at BGI-Shenzhen, and >4.0 Gb of reads for each sample were obtained.

Genome Annotation and Comparison

The paired-end reads were filtered using GetOrganelle pipeline (<https://github.com/Kinggerm/GetOrganelle>) to get plastid-like reads, then the filtered reads were assembled using SPAdes version 3.10 (Bankevich et al. 2012). To retain pure chloroplast contigs, the final “fastg” files were filtered using the “slim” script of GetOrganelle. The filtered De Bruijn graphs were viewed and edited using Bandage (Wick et al. 2015), then a circular chloroplast genome was generated. The genome was automatically annotated using CpGAVAS (Liu et al. 2012), then adjusted using Geneious version 9.1.7 (Kearse et al. 2012). The annotated chloroplast genomes

Table 1

Sampled Species of Lauraceae and Their Voucher Specimens Sequenced in This Study

No	Species	Herbarium	Taxon	Voucher	Geographic Origin	Accession Number in GenBank
1	<i>Eusideroxylon zwageri</i>	HITBC-BRG	<i>Eusideroxylon zwageri</i> Teijsm. & Binn.	SY34806	Sulawesi, Indonesia	MF939351
2	<i>Cryptocarya chinensis</i>	HITBC-BRG	<i>Cryptocarya chinensis</i> (Hance) Hemsl.	SY34239	Jianfenglin, Hainan	MF939349
3	<i>Cryptocarya hainanensis</i>	HITBC-BRG	<i>Cryptocarya hainanensis</i> Merr.	SY01426	Menghai, Yunnan	MF939350
4	<i>Beilschmiedia tungfangensis</i>	HITBC-BRG	<i>Beilschmiedia tungfangensis</i> S.K. Lee & L.F. Lau	SY34805	Wenshan, Yunnan	MF939348
5	<i>Beilschmiedia pauciflora</i>	HITBC-BRG	<i>Beilschmiedia pauciflora</i> H.W. Li	SY01364	Mengla, Yunnan	MF939347
6	<i>Cassytha filiformis</i>	HITBC-BRG	<i>Cassytha filiformis</i> Linnaeus	SY34802	Menghai, Yunnan	MF939337
7	<i>Cassytha capillaris</i>	HITBC-BRG	<i>Cassytha capillaris</i> Meisn.	SY34803	Sulawesi, Indonesia	MF939338
8	<i>Neocinnamomum caudatum</i>	HITBC-BRG	<i>Neocinnamomum caudatum</i> (Nees) Merr.	SY01561	Puer, Yunnan	MF939344
9	<i>Neocinnamomum lecomtei</i>	HITBC-BRG	<i>Neocinnamomum lecomtei</i> H. Liu	SY33249	Wenshan, Yunnan	MF939345
10	<i>Caryodaphnopsis henryi</i>	HITBC-BRG	<i>Caryodaphnopsis henryi</i> Airy Shaw	SY01542	Honghe, Yunnan	MF939346
11	<i>Caryodaphnopsis malipoensis</i>	HITBC-BRG	<i>Caryodaphnopsis malipoensis</i> Bing Liu & Y. Yang	SY32618	Wenshan, Yunnan	MF939343
12	<i>Actinodaphne trichocarpa</i>	HITBC-BRG	<i>Actinodaphne trichocarpa</i> C.K. Allen	SY32938	Emei, Sichuan	MF939342
13	<i>Neolitsea sericea</i>	HITBC-BRG	<i>Neolitsea sericea</i> (Blume) koidzumi	SY33307	Linan, Zhejiang	MF939341
14	<i>Nectandra angustifolia</i>	HITBC-BRG	<i>Nectandra angustifolia</i> (Schrad.) Nees & Mart.	SY34804	Sulawesi, Indonesia	MF939340
15	<i>Sassafras tzumu</i>	HITBC-BRG	<i>Sassafras tzumu</i> (Hemsl.) Hemsl.	SY34790	Anqing, Anhui	MF939339

have been submitted to GenBank (accession number: MF939337 to MF939351). The genome maps of all the 15 plastomes were drawn by OrganellarGenomeDRAW tool (OGDRAW; Lohse et al. 2013) and the gene organization maps were drawn by Gene Structure Display Server (GSDS) version 2.0 (Hu et al. 2015). Mauve version 2.4.0 software was used for alignment and determining the plastome rearrangements among the Magnoliids (Darling et al. 2004).

Phylogenetic Analysis

To estimate phylogenetic relationships within the magnoliids, 47 taxa with available complete plastomes were compared, including one taxon each from Canellales and Chloranthales, four from Piperales, six from Magnoliales, and 35 from Laurales. The 35 taxa included the 15 new plastomes and 20 complete plastomes which have been published elsewhere or adopted from NCBI (Song et al. 2015, 2016; Wu et al. 2017). *Amborella trichopoda* (AJ506156) was treated as the outgroup. For the species tree, maximum likelihood (ML) analyses were performed on data sets of 48 plastome sequences with single IR, SSC, and LSC regions. The whole genome matrix was aligned using MAFFT version 3.73 (Kato and Standley 2013), then manually edited using Geneious version 9.1.7 (Kearse et al. 2012). ML analysis was conducted using RAXML version 7.2.6 with the GTR + G model to search the best-scoring ML tree (Tamura et al. 2011). One thousand bootstrap replicates were performed to obtain the confidence support. Bayesian inference (BI) was performed using MrBayes version 3.2.6 (Ronquist and Huelsenbeck 2003).

The best-fit DNA substitution model of the Bayesian information criterion (BIC) was evaluated by using jModeltest version 2.1.10 (Darriba et al. 2012; Guindon et al. 2003). Markov Chain Monte Carlo (MCMC) analyses were run in MrBayes for 10,000,000 generations. The BI analysis started with a random tree and sampled every 1,000 generations. The first 25% of the trees was discarded as burn-in, and the remaining trees were used to generate a majority-rule consensus tree (supplementary fig. S1, Supplementary Material online). The trees were viewed and edited with the Fig tree version 1.4.0 software (<http://tree.bio.ed.ac.uk/software/figtree/>).

Results

Overall Structure and Gene Pool

Thirteen of the 15 newly sequenced Lauraceae plastomes displayed the typical quadripartite structure of angiosperms, including LSC, SSC, and a pair of IR regions, whereas the two plastomes from *Cassytha*, a genus of parasitic vines, have lost one copy of the IR (fig. 1). The complete plastome of *Cassytha filiformis* is 114,623 bp in length, 340 bp shorter than that of *Cassytha capillaris* (114,963 bp), and 42,954 bp shorter than that of *Eusideroxylon zwageri* (157,577 bp; table 2). Among the other 13 plastomes, genome size ranged from 150,749 bp (*Nectandra angustifolia*) to 158,530 bp (*Beilschmiedia tungfangensis*). In the LSC region, the length varied from 86,035 (*Caryodaphnopsis henryi*) to 93,803 bp (*Neolitsea sericea*), in the SSC region from 15,751 bp (*Caryodaphnopsis malipoensis*) to 19,222 bp (*Cryptocarya chinensis*), and in the IR region from 19,292 (*N. angustifolia*) to




FIG. 1.—Gene maps of the plastomes of *Cassytha*, *Eusideroxylon*, *Cryptocarya*, *Beilschmiedia*, *Caryodaphnopsis*, *Neocinnamomum*, *Nectandra*, *Sassafras*, *Neolitsea*, and *Actinodaphne* in the Lauraceae. The pink asterisks indicate the structural differences of IR loss.

25,601 bp (*C. henryi*). The plastomes of *Eusideroxylon*, *Cryptocarya*, *Beilschmiedia*, and *C. henryi* shared identical complements of coding genes; a total of 130 genes, including 8 rRNA genes, 37 tRNA, and 85 protein-coding genes, of which 17 are duplicated in IR regions. A total of 128 genes were detected on the plastomes of *Neocinnamomum*, *Nectandra*, *Sassafras*, *Actinodaphne*, *Neolitsea*, and *Caryodaphnopsis malipoensis*, 113 of which are single copy, while 15 are duplicated in IR regions. The different gene numbers reflect the duplication of *rpl23* and *trnI*-CAU in the first group. The plastomes of *Cassytha* have not only lost the duplicated genes in the IR region, but also six NADH dehydrogenase (*ndh*) genes, *ndhA*, *ndhC*, *ndhG*, *ndhI*, *ndhJ*, and *ndhK*, and their five *ndh* genes are pseudogenes.

Phylogenomic Analysis

The matrix of complete plastomes was used to reconstruct a phylogenetic tree of magnoliids (fig. 2). Magnoliids are divided into five main clades (ML-BS = 100%) corresponding to five orders: Canellales, Chloranthales, Laurales, Magnoliales, and Piperales. Sisterhood of Laurales and Magnoliales, with Piperales and Canellales being the next sister groups, was highly supported. Two major clades, including

Calycanthaceae and Lauraceae, were recognized within the Laurales. There was 100% support for the monophyly of Lauraceae family. Five well-supported groups were recovered within the Lauraceae (ML-BS = 100%). The basal group (ML-BS = 100%), including the genera *Eusideroxylon*, *Cryptocarya*, *Beilschmiedia*, and *Endiandra*, the *Cassytha* group (ML-BS = 100%), the *Neocinnamomum* group (ML-BS = 100%), the *Caryodaphnopsis* group (ML-BS = 100%), and the core group (ML-BS = 100%), including *Alseodaphne*, *Persea*, *Phoebe*, *Machilus*, *Lindera*, *Laurus*, *Actinodaphne*, *Neolitsea*, *Litsea*, *Nectandra*, *Sassafras*, and *Cinnamomum*.

Plastome Comparisons

Synteny and rearrangements were detected in ten plastomes of Lauraceae. A significant degree of synteny was found within the basal group, including *E. zwageri* and *B. tungfangensis*, and the core group, including *N. angustifolia*, *Laurus nobilis*, *Lindera communis*, *Machilus balansae*, *Alseodaphne semecarpifolia*, *Neocinnamomum caudatum*, and *C. capillaris*. However, the two groups differ in the orientation of a 13.7-kb fragment flanked by *rps7* and *rpl2* (fig. 3). In the basal group, the *rps7*-*ndhB*-*trnL*-*ycf2*-*trnI*-*rpl23*-*rpl2* segment has been

Table 2

Summary of 15 Complete Plastomes of Lauraceae

	<i>Eusideroxylon zwageri</i>	<i>Cryptocarya chinensis</i>	<i>Cryptocarya hainanensis</i>	<i>Beilschmiedia tungfangensis</i>	<i>Beilschmiedia pauciflora</i>	<i>Cassytha filiformis</i>	<i>Cassytha capillaris</i>
Total cpDNA size (bp)	157,577	157,675	157,145	158,530	157,901	114,623	114,963
Length of LSC region (bp)	89,231	89,199	89,002	89,351	88,673	–	–
Length of IR region (bp)	24,717	24,627	24,621	25,473	25,496	–	–
Length of SSC region (bp)	18,912	19,222	18,901	18,233	18,236	–	–
Total GC content	39.10%	39.10%	39.10%	39.00%	39.00%	36.90%	36.90%
Total number of genes (unique)	130 (113)	130 (113)	130 (113)	130 (113)	130 (113)	107 (107)	107 (107)
protein encoding	85	85	85	85	85	73	73
tRNA	37	37	37	37	37	30	30
rRNA	8	8	8	8	8	4	4
Length of <i>ycf1</i> (bp)	5,493	5,460	5,436	5,436	5,460	5,211	5,211
Length of truncated <i>ycf1</i> (bp)	971	977	974	1,863	1,863	–	–
Length of <i>ycf2</i> (bp)	6,882	6,885	6,885	6,843	6,849	5,583	5,583
Length of complete or truncated <i>ycf2</i> (bp)	6,882	6,885	6,885	6,843	6,849	–	–

<i>Neocinnamomum caudatum</i>	<i>Neocinnamomum lecomtei</i>	<i>Caryodaphnopsis henryi</i>	<i>Caryodaphnopsis malipoensis</i>	<i>Actinodaphne trichocarpa</i>	<i>Neolitsea sericea</i>	<i>Nectandra angustifolia</i>	<i>Sassafras tzumu</i>
150,842	150,838	154,938	149,239	152,739	152,442	150,749	151,798
91,881	91,912	86,035	91,901	93,783	93,803	93,783	92,752
20,257	20,257	25,601	20,036	20,078	20,067	19,292	20,096
18,447	18,412	17,701	17,266	18,800	18,505	18,382	18,854
38.80%	38.80%	39.00%	39.00%	39.20%	39.20%	39.20%	39.20%
128 (113)	128 (113)	131 (113)	128 (113)	128 (113)	128 (113)	128 (113)	128 (113)
84	84	86	84	84	84	84	84
36	36	37	36	36	36	36	36
8	8	8	8	8	8	8	8
5,517	5,517	5,526	5,526	5,574	5,568	5,535	5,586
928	928	1,473	1,473	1,378	1,372	1,372	1,419
6,831	6,831	6,894	6,894	6,876	6,846	6,909	6,294
3,110	3,110	6,894	3,186	3,168	3,162	2,478	3,168

combined with *trnH*-GUG, whereas the segment of the core group species has been combined with *rps19* (fig. 4), indicating that a rearrangement event occurred in Lauraceae plastome evolution. In the plastomes of *C. henryi* and in the basal group species, two unbroken protein-coding copies of *ycf2* were detected, suggesting that fragmentation of *ycf2* has occurred in other species of Lauraceae. Moreover, upstream of *rps19* adjoining the IR region, we detected one copy of a protein-coding gene *rpl23* and a tRNA gene *trnM*-CAU in the plastome of *C. henryi* and the basal group species, but not in the plastomes of other species, indicating that significant IR boundary changes occurred in Lauraceae plastome evolution.

IR Expansion and Contraction

In the sequenced plastomes of Lauraceae, two complete or fragmented copies of *ycf1* and *ycf2* were located at the

boundaries between the IR regions and the LSC or SSC regions. The full lengths of *ycf2* and *ycf1* ranged from 5,583 bp in *Cassytha filiformis* to 6,894 bp in *Caryodaphnopsis malipoensis* and from 5,211 bp in *Cassytha filiformis* to 5,586 bp in *Sassafras tzumu*, respectively (table 2). Double complete copies of the *ycf2* genes were detected in the seven sequenced Lauraceae plastomes of the basal group species, but only one complete copy and one fragment in the 24 plastomes of *C. malipoensis*, *Neocinnamomum*, and the core group species, except those of *C. henryi* and both *Cassytha* species. The length of the fragment of *ycf2* ranged from 2,478 bp in *N. angustifolia* to 3,168 bp in *Actinodaphne trichocarpa*. In contrast, all 32 sequenced Lauraceae plastomes, except the two species of *Cassytha*, had one complete copy and a fragment of *ycf1*. The length of the fragment of *ycf1* ranged from 971 bp in *E. zwageri* to 1,863 bp in *Beilschmiedia pauciflora*. Neither *Cassytha* plastome had fragments of *ycf1* and *ycf2*, but only one complete copy of each due to the IR loss.




FIG. 2.—Molecular phylogenetic tree of 47 taxa of Magnoliids based on complete plastome sequences using unpartitioned ML. Numbers at each node are bootstrap support value.

Discussion

Relationships in Lauraceae

This study included 47 complete chloroplast genomes for plants from all five orders (Canellales, Chloranthales, Laurales, Magnoliales, and Piperales) of the magnoliids. All of these complete plastome sequences of Lauraceae and related families yielded a fully resolved tree, consistent with the Angiosperm Phylogeny Group's most recent phylogeny, APG IV (Byng et al.

2016). Relationships among the five orders of the magnoliids are clarified as sisterhood of Laurales and Magnoliales, with Piperales and Canellales being the next sister groups, and Chloranthales the most basal group. Calycanthaceae and Lauraceae were recognized within the Laurales. All of these clades were recognized by Renner (Renner 1999).

The deep relationships of 34 Lauraceae taxa are separated into the following groups in our study. *Eusideroxylon*, *Cryptocarya*, *Beilschmiedia*, and *Endiandra* form the first




FIG. 3.—Synteny and rearrangements detected in Lauraceae plastomes using the Mauve multiple-genome alignment program. The yellow asterisks 1 indicate the varied gene locus of *rpl2*, the blue asterisks 2 indicate a rearrangement of the fragment flanked by *rpl2* and *rps7*, and the green asterisks 3 indicate an insert of intergenic region *rpl2*–*rpl23*.




FIG. 4.—Comparisons of gene loci in the fragments flanked by *rps19* and *trnL* in IRa and *trnL* and *trnH* in IRb among 15 plastomes of Magnoliids.

group in the phylogeny. *Cassytha*, *Neocinnamomum*, and *Caryodaphnopsis* form the second, third, and fourth groups, respectively. The fifth group includes *Alseodaphne*, *Persea*, *Phoebe*, and *Machilus*. The sixth group includes *Nectandra*, *Sassafras*, and *Cinnamomum*. And the last group includes

Lindera, *Laurus*, *Litsea*, *Actinodaphne*, and *Neolitsea*. The phylogenetic placements of the first, fourth, fifth, and sixth groups are consistent with previously published phylogenetic relationships (Chanderbali et al. 2001; Rohwer and Rudolph 2005). The position of *Cassytha*, considered as a ‘jumping

genus' by Rohwer and Rudolph (2005), was settled here in the way predicted from morphology (Chanderbali et al. 2001). The seventh group, equivalent to the tribe Laureae (Chanderbali et al. 2001), was confirmed as sister to the sixth group, tribe Cinnamomeae (including *Sassafras*), which has always been assumed based on morphological characters, although previous molecular analyses failed to prove it convincingly (Chanderbali et al. 2001; Rohwer and Rudolph 2005).

Unusual Structure of the *Cassytha* Plastomes

The sizes of the fifteen newly sequenced Lauraceae plastomes differed greatly, from 114,623 bp in the hemiparasitic vine, *C. capillaris*, to 158,530 bp in *B. tungfangensis*, as a result of the loss of one IR copy and six *ndh* genes in *Cassytha*. *Cassytha* is the only stem hemiparasitic genus with reduced leaves and roots in the magnoliids, and the only nonwoody member of the Lauraceae. We show that it is also unique in the Lauraceae the loss of one IR copy in its plastome, although similar losses have occurred independently in the Leguminosae (Cai et al. 2008), Pinaceae (Raubeson and Jansen 1992), Cephalotaxaceae (Yi et al. 2013), and cupressophytes (Wu et al. 2011a). In addition, six *ndh* genes, *ndhA*, *ndhC*, *ndhG*, *ndhI*, *ndhJ*, and *ndhK*, have been lost, and the other five, *ndhB*, *ndhD*, *ndhE*, *ndhF*, and *ndhH*, are clearly pseudogenes in both *Cassytha* taxa sequenced in this study. All eleven *ndh* genes encode independent subunits of a plastid NADPH-dehydrogenase complex (*Ndh* 1-complex) which carries out one of the recycled electron pathways around Photosystem I (Casano et al. 2000). Cyclic electron flow is vital for maintenance of efficient photosynthesis and enablement of photoprotection under environmental stresses in higher plants (Wang et al. 2006). The *ndh* genes are frequently pseudogenized or lost in plant groups with a degree of heterotrophy, such as *Aneura*, *Cuscuta*, *Epifagus*, *Hydnora*, and nonphotosynthetic orchid species, and in some autotrophic gymnosperms and ferns (dePamphilis and Palmer 1990; Wicke et al. 2011; Wickett et al. 2008; McNeal et al. 2007; Kim et al. 2015; Naumann et al. 2016), but this is first report for *Cassytha*, the only hemiparasitic genus in the Laurales. This adds to the evidence that the *Ndh* 1-complex is not essential for plant survival, while the *ndh*-independent antimycin-A-sensitive pathway, which functions in cyclic electron flow as another choice, could be more important under most conditions (Shikanai 2014).

Loss Events in the Laurales

Comparative genomic analysis indicated that missing segments of DNA in Lauraceae plastids mainly drive the genome contraction events. A fragment flanked by *rps7* and *rpl2* was detected as a rearrangement event between the basal group species and the other species except *C. henryi*. However, it looks more like two or more independent loss events when

we choose the plastomes of *C. henryi* or nonLaurales species as reference. Double IR fragments with the gene order of *trnL-ycf2-trnI-rpl23-rpl2* are highly conserved in the plastomes of *C. henryi* (fig. 4) and nonLaurales genera such as *Drimys*, *Piper*, *Liriodendron*, and *Magnolia* (Cai et al. 2006; Zhu et al. 2016; Yang et al. 2014), indicating the plastome of *C. henryi* is evolutionarily conserved. In *Calycanthus* (Laurales) plastome (Goremykin et al. 2003a), one copy of *rpl2* with the length of 1,480 bp disappeared from the *trnL-rpl2* fragment in IRb, but all of the sequenced Lauraceae plastomes of the basal group, including *Endiandra*, *Beilschmiedia*, *Cryptocarya*, and *Eusideroxylon*, lost another copy of *rpl2* from the *trnL-rpl2* fragment in the IRa region (fig. 4). More interesting are the sequenced Lauraceae plastomes of the core group, including *Alseodaphne*, *Persea* (Song et al. 2016), *Phoebe*, *Machilus* (Song et al. 2015), *Lindera*, *Laurus*, *Litsea*, *Nectandra*, *Sassafras*, *Cinnamomum* (Wu et al. 2017), *Actinodaphne*, and *Neolitsea*, which have further lost a segment of at least 4,500 bp which contains a fragment of *ycf2* and one copy of *rpl23* and *trnL-CAU* in IRb of *Calycanthus*. This segment was also lost in the plastomes of *Neocinnamomum* species and *C. malipoensis*. Taken together, these independent loss events show that in the Lauraceae the plastomes of *Neocinnamomum*, *Cassytha*, the core group, and the basal group could share a common ancestral genome structure like that of *C. henryi*, but have subsequently evolved independently with different loss patterns.

Evolutionary Pattern in Angiosperms

To put these results in a wider phylogenetic context, we traced the fragments flanked by *trnL-CAA* and *rps19* in the IRa region and by *trnL-CAA* and *trnH-GUG* in the IRb region in the six major groups of the angiosperms and found that the gene backbone and order are conserved (fig. 5). In the early-diverging angiosperm species, *A. trichopoda* and *Nymphaea alba*, of the ANITA group (Qiu et al. 1999), the gene orders of the fragments are *rps19-rpl2-rpl23-trnL-ycf2-trnL* and *trnL-ycf2-trnL-rpl23-rpl2-trnH* (Goremykin et al. 2003b, 2004). These orders are retained in the early diverging monocot *Tofieldia thibetica* (Luo et al. 2016) and *Ceratophyllum demersum* in the Ceratophyllaceae (Moore et al. 2007). In the early diverging eudicot *Euptelea pleiosperma* (Sun et al. 2016), the only change in the gene order is a new insertion of a fragment of *rps19*. In the magnoliids, the same gene order for both fragments is retained in the sequenced species of Choranthaceae (Hansen et al. 2007), Piperales (Cai et al. 2006), and Magnoliales (Zhu et al. 2016), but a new copy of *trnH* has been inserted between *rps19* and *rpl2* in the IRa fragment of *Drimys granadensis* in the Canellales (Cai et al. 2006) and the copy of *rpl2* has been lost between *rps19* and *rps23* in the IRa region of *Endiandra*, *Beilschmiedia*, *Cryptocarya*, and *Eusideroxylon* species in Lauraceae, and in




Fig. 5.—Model of the origin and variation of the fragments flanked by *rps19* and *trnL* in IRa and *trnL* and *trnH* in IRb among plastomes of angiosperms. The pink asterisks indicate the varied gene loci.

IRb of *Calycanthus* in Calycanthaceae (Goremykin et al. 2003a). Nevertheless, our comparative genomic analysis concluded that the regions encompassing the *ycf2* and the adjoined *trnH*-GUG or *trnL*-CAA gene in the plastomes of *C. henryi* and other early-diverging angiosperms are the retained IRs, corresponding to either IRa or IRb in the basal and core groups of Lauraceae.

Supplementary Material

Supplementary data are available at *Genome Biology and Evolution* online.

Acknowledgments

The authors would like to acknowledge Jing Yang, Juan-Hong Zhang, Zheng-Shan He, Chun-Yan Lin, and Ji-Xiong Yang at the Germplasm Bank of Wild Species, Kunming Institute of Botany, Chinese Academy Sciences, for sequencing technology. They sincerely thank the anonymous referees and Prof. Giovanni Vendramin for their critical and invaluable comments that greatly improved our manuscript. This work was supported by the National Natural Science Foundation of China (No. 31600531), a grant of the

Large-scale Scientific Facilities, CAS (No.2017-LSF-GBOWS-02), the CAS “Light of West China” Program (Y7XB061B01), the 1000 Talents Program (WQ20110491035), and the project of the Southeast Asia Biodiversity Research Institute, CAS.

Literature Cited

- Bankovich A, et al. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol.* 19(5):455–477.
- Barrett CF, et al. 2014. Investigating the path of plastid genome degradation in an early-transitional clade of heterotrophic orchids, and implications for heterotrophic angiosperms. *Mol Biol Evol.* 31(12):3095–3112.
- Bock R. 2007. Plastid biotechnology: prospects for herbicide and insect resistance, metabolic engineering and molecular farming. *Curr Opin Biotechnol.* 18(2):100–106.
- Braukmann TWA, Kuzmina M, Stefanovic S. 2009. Loss of all plastid *ndh* genes in Gnetales and conifers: extent and evolutionary significance for the seed plant phylogeny. *Curr Genet.* 55(3):323–337.
- Byng JW, et al. 2016. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. *Bot J Linn Soc.* 181(1):1–20.
- Cai Z, et al. 2006. Complete plastid genome sequences of *Drimys*, *Liriodendron*, and *Piper*: implications for the phylogenetic relationships of magnoliids. *BMC Evol Biol.* 6:77.

- Cai ZQ, et al. 2008. Extensive reorganization of the plastid genome of *Trifolium subterraneum* (Fabaceae) is associated with numerous repeated sequences and novel DNA insertions. *J Mol Evol.* 67(6):696–704.
- Casano LM, Zapata JM, Martin M, Sabater B. 2000. Chlororespiration and poisoning of cyclic electron transport. Plastocyanin as electron transporter between thylakoid NADH dehydrogenase and peroxidase. *J Biol Chem.* 275(2):942–948.
- Chanderbali AS, van der Werff H, Renner SS. 2001. Phylogeny and historical biogeography of Lauraceae: evidence from the chloroplast and nuclear genomes. *Ann Missouri Bot Gard.* 88(1):104–134.
- Chang CC, et al. 2006. The chloroplast genome of *Phalaenopsis aphrodite* (Orchidaceae): comparative analysis of evolutionary rate with that of grasses and its phylogenetic implications. *Mol Biol Evol.* 23(2):279–291.
- Darling ACE, Mau B, Blattner FR, Perna NT. 2004. Mauve: multiple alignment of conserved genomic sequence with rearrangements. *Genome Res.* 14(7):1394–1403.
- Darriba D, Taboada GL, Doallo R, Posada D. 2012. jModelTest 2: more models, new heuristics and parallel computing. *Nat Methods.* 9(8):772–772.
- dePamphilis CW, Palmer JD. 1990. Loss of photosynthetic and chlororespiratory genes from the plastid genome of a parasitic flowering plant. *Nature* 348(6299):337–339.
- Doyle JJ, Dickson EE. 1987. Preservation of plant-samples for DNA restriction endonuclease analysis. *Taxon* 36(4):715–722.
- Doyle JJ, Doyle JL, Palmer JD. 1995. Multiple independent losses of 2 genes and one intron from legume chloroplast genomes. *Syst Bot.* 20(3):272–294.
- Gao L, Su Y-J, Wang T. 2010. Plastid genome sequencing, comparative genomics, and phylogenomics: current status and prospects. *J Syst Evol.* 48(2):77–93.
- Goremykin V, Hirsch-Ernst K, Wolf S, Hellwig F. 2003a. The chloroplast genome of the “basal” angiosperm *Calycanthus fertilis* – structural and phylogenetic analyses. *Plant Syst Evol.* 242(1):119–135.
- Goremykin VV, Hirsch-Ernst KI, Wolf S, Hellwig FH. 2003. Analysis of the *Amborella trichopoda* chloroplast genome sequence suggests that amborella is not a basal angiosperm. *Mol Biol Evol.* 20(9):1499–1505. [pii]
- Goremykin VV, Hirsch-Ernst KI, Wolf S, Hellwig FH. 2004. The chloroplast genome of *Nymphaea alba*: whole-genome analyses and the problem of identifying the most basal angiosperm. *Mol Biol Evol.* 21(7):1445–1454. [pii]
- Goremykin VV, Holland B, Hirsch-Ernst KI, Hellwig FH. 2005. Analysis of *Acorus calamus* chloroplast genome and its phylogenetic implications. *Mol Biol Evol.* 22(9):1813–1822.
- Guindon S, Gascuel O, Rannala B. 2003. A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Syst Biol.* 52(5):696–704.
- Guisinger M, Chumley T, Kuehl J, Boore J, Jansen R. 2010. Implications of the plastid genome sequence of *Typha* (Typhaceae, Poales) for understanding genome evolution in Poaceae. *J Mol Evol.* 70(2):149–166.
- Guisinger MM, Kuehl JV, Boore JL, Jansen RK. 2011. Extreme reconfiguration of plastid genomes in the angiosperm family Geraniaceae: rearrangements, repeats, and codon usage. *Mol Biol Evol.* 28(1):583–600.
- Hansen DR, et al. 2007. Phylogenetic and evolutionary implications of complete chloroplast genome sequences of four early-diverging angiosperms: *Buxus* (Buxaceae), *Chloranthus* (Chloranthaceae), *Dioscorea* (Dioscoreaceae), and *Illicium* (Schisandraceae). *Mol Phylogenet Evol.* 45(2):547–563.
- Hirao T, Watanabe A, Kurita M, Kondo T, Takata K. 2008. Complete nucleotide sequence of the *Cryptomeria japonica* D. Don. chloroplast genome and comparative chloroplast genomics: diversified genomic structure of coniferous species. *BMC Plant Biol.* 8:70.
- Hu B, et al. 2015. GSDS 2.0: an upgraded gene feature visualization server. *Bioinformatics* 31(8):1296–1297.
- Karol KG, et al. 2010. Complete plastome sequences of *Equisetum arvense* and *Isoetes flaccida*: implications for phylogeny and plastid genome evolution of early land plant lineages. *BMC Evol Biol.* 10(1):321.
- Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol.* 30(4):772–780.
- Kearse M, et al. 2012. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28(12):1647–1649.
- Kim HT, et al. 2015. Seven new complete plastome sequences reveal rampant independent loss of the *ndh* gene family across orchids and associated instability of the inverted repeat/small single-copy region boundaries. *PLoS ONE.* 10(11):e0142215.
- Kim YD, Jansen RK. 1994. Characterization and phylogenetic distribution of a chloroplast DNA rearrangement in the Berberidaceae. *Plant Syst Evol.* 193(1–4):107–114.
- Lavin M, Doyle JJ, Palmer JD. 1990. Evolutionary significance of the loss of the chloroplast-DNA inverted repeat in the leguminosae subfamily Papilionoideae. *Evolution* 44(2):390–402.
- Liu C, et al. 2012. CpGAVAS, an integrated web server for the annotation, visualization, analysis, and GenBank submission of completely sequenced chloroplast genome sequences. *BMC Genomics.* 13:715.
- Lohse M, Drechsel O, Kahlau S, Bock R. 2013. OrganellarGenomeDRAW – a suite of tools for generating physical maps of plastid and mitochondrial genomes and visualizing expression data sets. *Nucleic Acids Res.* 41(Web Server issue):W575–W581.
- Luo Y, et al. 2016. Plastid phylogenomic analyses resolve Tofieldiaceae as the root of the early diverging monocot order Alismatales. *Genome Biol Evol.* 8(3):932–945.
- McCoy SR, Kuehl JV, Boore JL, Raubeson LA. 2008. The complete plastid genome sequence of *Welwitschia mirabilis*: an unusually compact plastome with accelerated divergence rates. *BMC Evol Biol.* 8:130.
- McNeal JR, Kuehl JV, Boore JL, de Pamphilis CW. 2007. Complete plastid genome sequences suggest strong selection for retention of photosynthetic genes in the parasitic plant genus *Cuscuta*. *BMC Plant Biol.* 7:57.
- Moore MJ, Bell CD, Soltis PS, Soltis DE. 2007. Using plastid genome-scale data to resolve enigmatic relationships among basal angiosperms. *Proc Natl Acad Sci U S A.* 104(49):19363–19368.
- Naumann J, et al. 2016. Detecting and characterizing the highly divergent plastid genome of the nonphotosynthetic parasitic plant *Hydnora visseri* (Hydnoraceae). *Genome Biol Evol.* 8(2):345–363.
- Palmer JD, Osorio B, Aldrich J, Thompson WF. 1987. Chloroplast DNA evolution among legumes – loss of a large inverted repeat occurred prior to other sequence rearrangements. *Curr Genet.* 11(4):275–286.
- Plunkett GM, Downie SR. 2000. Expansion and contraction of the chloroplast inverted repeat in Apiaceae subfamily Apioideae. *Syst Bot.* 25(4):648–667.
- Qiu YL, et al. 1999. The earliest angiosperms: evidence from mitochondrial, plastid and nuclear genomes. *Nature* 402(6760):404–407.
- Raubeson LA, Jansen RK. 1992. A rare chloroplast-DNA structural mutation is shared by all conifers. *Biochem Syst Ecol.* 20(1):17–24.
- Renner SS. 1999. Circumscription and phylogeny of the Laurales: evidence from molecular and morphological data. *Am J Bot.* 86(9):1301–1315.
- Rohwer JG, Rudolph B. 2005. Jumping genera: the phylogenetic positions of *Cassytha*, *Hypodaphnis*, and *Neocinnamomum* (Lauraceae) based on different analyses of *trnK* intron sequences. *Ann Missouri Bot Gard.* 92(2):153–178.
- Ronquist F, Huelsenbeck JP. 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19(12):1572–1574.

- Roper JM, et al. 2007. The complete plastid genome sequence of *Angiopteris evecta* (G. Forst.) Hoffm. (Marattiaceae). *Am Fern J.* 97(2):95–106.
- Sanderson MJ, et al. 2015. Exceptional reduction of the plastid genome of saguaro cactus (*Carnegiea gigantea*): loss of the *ndh* gene suite and inverted repeat. *Am J Bot.* 102(7):1115–1127.
- Shikanai T. 2014. Central role of cyclic electron transport around photosystem I in the regulation of photosynthesis. *Curr Opin Biotechnol.* 26:25–30.
- Song Y, et al. 2015. Comparative analysis of complete chloroplast genome sequences of two tropical trees *Machilus yunnanensis* and *Machilus balansae* in the family Lauraceae. *Front Plant Sci.* 6:662.
- Song Y, Yao X, Tan YH, Gan Y, Corlett RT. 2016. Complete chloroplast genome sequence of the avocado: gene organization, comparative analysis, and phylogenetic relationships with other Lauraceae. *Can J For Res.* 46(11):1293–1301.
- Sun Y, et al. 2016. Phylogenomic and structural analyses of 18 complete plastomes across nearly all families of early-diverging eudicots, including an angiosperm-wide analysis of IR gene content evolution. *Mol Phylogenet Evol.* 96:93–101.
- Sun YX, et al. 2013. Complete plastid genome sequencing of trochodendraceae reveals a significant expansion of the inverted repeat and suggests a paleogene divergence between the two extant species. *PLoS ONE.* 8(4):e60429.
- Tamura K, et al. 2011. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol.* 28(10):2731–2739.
- Turmel M, Otis C, Lemieux C. 2006. The chloroplast genome sequence of *Chara vulgaris* sheds new light into the closest green algal relatives of land plants. *Mol Biol Evol.* 23(6):1324–1338.
- Wakasugi T, et al. 1994. Loss of all *ndh* genes as determined by sequencing the entire chloroplast genome of the black pine *Pinus thunbergii*. *Proc Natl Acad Sci U S A.* 91(21):9794–9798.
- Wang P, et al. 2006. Chloroplastic NAD(P)H dehydrogenase in tobacco leaves functions in alleviation of oxidative damage caused by temperature stress. *Plant Physiol.* 141(2):465–474.
- Wang R-J, et al. 2008. Dynamics and evolution of the inverted repeat-large single copy junctions in the chloroplast genomes of monocots. *BMC Evol Biol.* 8(1):36.
- Wang W, Messing J. 2011. High-throughput sequencing of three Lemnoideae (duckweeds) chloroplast genomes from total DNA. *PLoS ONE.* 6(9):e24670.
- Wick RR, Schultz MB, Zobel J, Holt KE. 2015. Bandage: interactive visualization of de novo genome assemblies. *Bioinformatics* 31(20):3350–3352.
- Wicke S, et al. 2016. Mechanistic model of evolutionary rate variation en route to a nonphotosynthetic lifestyle in plants. *Proc Nat Acad Sci U S A.* 113(32):9045–9050.
- Wicke S, Schneeweiss GM, dePamphilis CW, Müller KF, Quandt D. 2011. The evolution of the plastid chromosome in land plants: gene content, gene order, gene function. *Plant Mol Biol.* 76(3–5):273–297.
- Wickett NJ, Forrest LL, Budke JM, Shaw B, Goffinet B. 2011. Frequent pseudogenization and loss of the plastid-encoded sulfate-transport gene *cysA* throughout the evolution of liverworts. *Am J Bot.* 98(8):1263–1275.
- Wickett NJ, et al. 2008. Functional gene losses occur with minimal size reduction in the plastid genome of the parasitic liverwort *Aneura mirabilis*. *Mol Biol Evol.* 25(2):393–401.
- Wu CC, Chu FH, Ho CK, Sung CH, Chang SH. 2017. Comparative analysis of the complete chloroplast genomic sequence and chemical components of *Cinnamomum micranthum* and *Cinnamomum kanehirae*. *Holzforchung* 71(3):189–197.
- Wu CS, Lai YT, Lin CP, Wang YN, Chaw SM. 2009. Evolution of reduced and compact chloroplast genomes (cpDNAs) in gnetophytes: selection toward a lower-cost strategy. *Mol Phylogenet Evol.* 52(1):115–124.
- Wu CS, Lin CP, Hsu CY, Wang RJ, Chaw SM. 2011. Comparative chloroplast genomes of Pinaceae: insights into the mechanism of diversified genomic organizations. *Genome Biol Evol.* 3:309–319.
- Wu CS, Wang YN, Hsu CY, Lin CP, Chaw SM. 2011. Loss of different inverted repeat copies from the chloroplast genomes of Pinaceae and cupressophytes and influence of heterotachy on the evaluation of gymnosperm phylogeny. *Genome Biol Evol.* 3:1284–1295.
- Yang JB, Li DZ, Li HT. 2014. Highly effective sequencing whole chloroplast genomes of angiosperms by nine novel universal primer pairs. *Mol Ecol Resour.* 14(5):1024–1031.
- Yi X, Gao L, Wang B, Su YJ, Wang T. 2013. The complete chloroplast genome sequence of *Cephalotaxus oliveri* (Cephalotaxaceae): evolutionary comparison of cephalotaxus chloroplast DNAs and insights into the loss of inverted repeat copies in gymnosperms. *Genome Biol Evol.* 5(4):688–698.
- Zhu AD, Guo WH, Gupta S, Fan WS, Mower JP. 2016. Evolutionary dynamics of the plastid inverted repeat: the effects of expansion, contraction, and loss on substitution rates. *New Phytol.* 209(4):1747–1756.

Associate editor: Bill Martin