

Review

A Horizon Scan of Emerging Global Biological Conservation Issues for 2020

William J. Sutherland,^{1,*} Maria P. Dias,^{2,3} Lynn V. Dicks,^{1,4} Helen Doran,⁵ Abigail C. Entwistle,⁶ Erica Fleishman,⁷ David W. Gibbons,^{8,9} Rosie Hails,¹⁰ Alice C. Hughes,¹¹ Jonathan Hughes,¹² Ruth Kelman,¹³ Xavier Le Roux,^{14,15} Becky LeAnstey,¹⁶ Fiona A. Lickorish,¹⁷ Luke Maggs,¹⁸ James W. Pearce-Higgins,^{1,19} Lloyd S. Peck,²⁰ Nathalie Pettorelli,²¹ Jules Pretty,²² Mark D. Spalding,²³ Femke H. Tonneijck,²⁴ Jonathan Wentworth,²⁵ and Ann Thornton¹

In this horizon scan, we highlight 15 emerging issues of potential relevance to global conservation in 2020. Seven relate to potentially extensive changes in vegetation or ecological systems. These changes are either relatively new, for example, conversion of kelp forests to simpler macroalgal systems, or may occur in the future, for example, as a result of the derivation of nanocellulose from wood or the rapid expansion of small hydropower schemes. Other topics highlight potential changes in national legislation that may have global effect on international agreements. Our panel of 23 scientists and practitioners selected these issues using a modified version of the Delphi technique from a long-list of 89 potential topics.

Horizon Scanning for Conservation

This 11th annual horizon scan identified novel issues that may have substantive positive or negative effects on global biological conservation. We do not aim to predict outcomes, but to highlight issues to which societies may wish or need to respond to in the future on the basis of improved knowledge. Here, we present the 15 topics identified by our panel, which comprised 23 scientists, conservation practitioners, and experts in foresight research and horizon scanning. These topics, about which we believe relatively little is known among those working in conservation, are diverse. They include effects on wildlife of a range-expanding, invasive tick species; implications of new legislation to promote sourcing of energy from wood; and the application of a genetically modified fungus to kill malaria-carrying mosquitoes. Many of the issues we identified in previous scans have been realized or become better understood [1].

Identification of Issues

Our annual horizon-scanning methods have been consistent (Figure 1). We apply a modified version of the Delphi technique, which facilitates a process that is repeatable, inclusive, and transparent [2,3]. This year's 23 core participants had diverse subject-matter expertise, including, but not limited to, agriculture and land use, microbiology, conservation practice and technology, sustainability, environmental management, policy, economics, research programming, science communication, and professional horizon scanning.

Members of our team used different methods to identify and select issues, including, but not limited to, consulting with colleagues in person or by email, soliciting issues via Twitter and other forms of social media or through established networks, and tracking via curation tools, such as *pearltrees*¹. We engaged approximately 830 people. If face-to-face meetings were held, then we counted all participants. If messages were sent by email or social media, we counted only those who responded. In addition, we noted issues that we encountered throughout the year in the popular and social media, scientific journals, seminars, and other professional presentations, and even casual conversations in which we engaged or that we overheard.

We assessed the suitability of each issue for inclusion on the basis of criteria established during the first horizon scan [4]: issues must be novel or represent novelty through a step-change in impact, have potentially substantive positive or negative impacts on conservation of biological diversity at a global or regional level, and appear likely to have greater impact in the future.

Highlights

We present 15 issues that emerged from our 11th annual horizon scan of global biological conservation.

Issues were identified by a diverse group of scientists and practitioners.

The group scored the long-list of 89 issues using a Delphi-like process. Issues were ranked according to novelty, relevance, and potential impact on biological conservation.

The top-38 ranked issues were debated at a workshop in Cambridge, UK, with each topic scored immediately following the discussion.

Emerging themes include changes in national legislation that could affect international agreements; recognising ecocide as an international crime; and the global decline in kelp forests.

Other issues include the increasing use of traditional Asian medicine; the spread of the Asian long-horned tick; use of artificial wombs in mammalian conservation; and the impact that the shrinking ozone hole may have on Antarctic sea ice.

¹Conservation Science Group, Department of Zoology, Cambridge University, The David Attenborough Building, Pembroke Street, Cambridge CB2 3QZ, UK

²BirdLife International, The David Attenborough Building, Pembroke Street, Cambridge CB2 3QZ, UK

³MARE Marine and Environmental Sciences Centre, ISPA, Instituto






Figure 1. Our Process for Identifying and Evaluating Issues during the 2020 Scan.

For this scan, we compiled a long-list of 89 issues and grouped them on the basis of theme. Where two closely related issues were submitted by different participants, we combined the issues and assessed them as one. Participants independently and confidentially scored each issue from 0 to 1000

(low–high) on the basis of its potential effects on biological conservation, and noted if they were previously aware of the issue. Participants often added notes or queries to their scoresheet (e.g., suggesting issues that could be combined or topics that were well known). These notes and queries informed discussion at the expert meeting (see later). Fatigue can result in attention waning throughout a process of scoring issues (e.g., [5]). To counter this potential bias, the long-list was ordered in three different ways, and participants randomly assigned to each list. Each participant's scores were converted to ranks (1–89). The most highly ranked 38 issues (due to a four-way tie for 35th place) were retained for further discussion and a second round of assessment (described later).

Two weeks before the expert meeting, each topic was assigned to two participants for further investigation to ensure detailed preparation. The person who submitted a given topic was not assigned that topic. Therefore, investigators were unlikely to be experts on the topic, but could collate relevant information to allow assessment of its potential impacts. Investigators focused on the novelty, likelihood of occurrence, magnitudes of impact, and relevance to biological conservation of the issues.

At an expert meeting in Cambridge (UK) in September 2019, we discussed each issue in relation to the above criteria. During discussions, we clarified technical details and levels of awareness, and presented further sources of evidence. After the discussion of each issue, all participants rescored that issue as in the first round. These scores again were converted to ranks, from which we calculated a median rank. Here, we present the 15 issues with highest median ranks. The issues are in thematic rather than rank order.

The 2020 Issues

Land-Use Change in Response to Derivation of Nanocellulose from Wood

Innovations in materials science, particularly in nanocomposites, are beginning to create novel opportunities for manufacturing that may increase demand for wood. Cellulose is a strong, stiff polymer produced by plants (particularly trees), animals, and bacteria. When cut into nanocellulose, its key properties change [6], making it suitable as a feedstock for industrial processes. Nanocellulose is used to produce a range of products, including construction materials, packaging for clothing, and consumer products, such as transparent wood-based packaging. The global nanocellulose market is growing by 18% per annum and is estimated to reach US\$660 million by 2023 [57], thus increasing demand for wood. In response, tree planting may increase, temporarily boosting carbon stocks and reducing reliance on fossil fuels. Nanocellulose products could replace plastics, so the volume of plastic waste may also decrease. However, unregulated demand for wood for the nanocellulose market could accelerate global forest loss. Biological diversity will also be lost if monoculture tree plantations replace natural ecosystems, or if stands of mature trees are clear harvested.

Policy Incentives for Derivation of Energy from Wood

Demand for wood may increase substantially in response to the combination of the European Union (EU) Renewable Energy Directive of 2018, which treats wood, even from biologically diverse forests, as a renewable energy source, and the EU pledge to double the 2015 renewable energy levels in Europe by 2030 to meet commitments in the 2015 Paris Agreement. As a result, EU demand may no longer be satisfied by current practices of deriving wood from forests in the USA and Canada and promoting the intensification of European forestry [7]. Such commercialisation of forest biomass may accelerate the loss of primary forest and exacerbate climate change. The latter changes could be amplified if similar policies are adopted by other countries following the lead of the EU. A lawsuit is now challenging the inclusion by the EU of forest biomass as a source of renewable energy.

Manipulating Floral Species Composition to Improve Bee Health

Variation in the nutritional content of nectar and pollen among plant species influences how bee species forage, but little is known about differences in nutritional requirements among species or populations of bees. Land uses that reduce the species richness and abundance of flowering plants are likely to interact with other factors, such as pathogens, to drive bee declines. Two recent studies conducted in the USA suggested that pollen from sunflowers *Helianthus* spp. and closely related

Asteraceae species reduces the prevalence of parasite infection in bees. Laboratory studies indicated a reduction in the load of the gut parasite *Crithidia bombi* in bumblebees *Bombus pensylvanicus*, and a field survey indicated lower levels of infection in areas with high cover of planted sunflowers, even though sunflower pollen has relatively low nutritional value compared with other pollens [8,9]. *C. bombi* also reduces reproductive success in colonies of the European bumblebee *Bombus terrestris* [10]. This new knowledge could be used to alter crop choices or planting regimes for bees before impacts on wild bee populations are fully understood. The latter actions may disrupt host–parasite dynamics in wild bees [11], and could also reduce nutritional availability relative to native wildflowers.

Asian Long-Horned Tick Reaches the Americas

The non-native Asian long-horned tick *Haemaphysalis longicornis* is well established in Australia and New Zealand, where it is associated with economically important cattle pathogens, including *Theileria orientalis*. First detected in the USA in 2017, the tick has now spread to nine states. Coincident infection of three unrelated cattle herds in Virginia, USA by the virulent, pathogenic Ikeda genotype of *T. orientalis* led to seven cattle deaths [12]. The tick has a wide climatic tolerance, with potential to colonise the eastern and western seabords of the USA and Canada and progress south into Central America [13] and to extend across South America. Invasion by both a novel tick and associated pathogens would represent a major emerging disease for the Americas. If the potential effects on animal health and livestock farming are of sufficient magnitude to affect regional and national economies by reducing milk or meat production, land use may change across extensive areas [12]. *H. longicornis* also has been associated with mortality of cattle in New Zealand and is carried by diverse host mammals and birds [14]. Therefore, its introduction to the Americas with at least one associated pathogen may also have population-level effects on native wildlife.

Global Declines of Kelp Forest

Kelps are an order of brown algae (Laminariales) with high primary productivity that occur on ~25% of coastlines worldwide and function as complex habitats for many other species. Declines in kelp abundance have been reported widely, albeit with substantial regional variability [15]. Kelps long have been considered resilient to environmental stress. However, this resilience may be waning in response to diversification of potential drivers of decline, which include increases in sea temperatures caused by anthropogenic climate change, non-native invasive species, eutrophication, and harvesting [15–17]. The declines may result in fundamental shifts from complex kelp forests to simpler macroalgal turf systems [17]. Future kelp forest declines in response to accelerating climate change would have significant consequences for biological diversity and ecosystem processes [16]. The ecological benefits to humans that are supported by kelp forests, including commercial fisheries and shoreline protection, are valued at billions of dollars annually [16].

Atmospheric Circulation and the Shrinking Antarctic Ozone Hole May Affect Extent of Polar Ice

Rising sea levels, in large part caused by melting of the polar ice caps and thermal expansion of water, affect human societies, land use, and coastal ecosystems worldwide. Understanding of how the extent of Antarctic coastal and sea ice responds to interactions with stratospheric ozone, atmospheric circulation, and storms and waves is evolving rapidly (e.g., [18]). Reduced chlorofluorocarbon emissions since the Montreal Protocol of 1987 have led to less ozone depletion during polar winters and, as a result, a reduction in the size of the ozone holes. However, in the Antarctic, the shrinking of the ozone hole may weaken the north–south movement of the westerly wind belt that circles the continent [19], the Southern Annular Mode (SAM), counteracting the general strengthening of the SAM as concentrations of greenhouse gases increase. Variation in the movement of the SAM alters Antarctic temperature and storm patterns [20]. For example, when the SAM is at its southernmost position, westerly winds strengthen over Antarctica and sea-surface and air temperatures decrease. Decreases in the size of the ozone hole are among the factors implicated in changes in wind and other weather patterns that will likely contribute to decreases in the extent of Antarctic ice and increases in global sea levels.

Effects of Small Hydropower Systems on Riverine Ecosystems

The cumulative environmental effects of >80 000 small hydropower dams built on small upland streams have received less attention than those of large dams. There are now over 11 small dams for every large dam [21], and efforts are underway in the Himalayas, other mountain ranges in Asia, and the Andes [22] to increase their use to empower local communities. Such small run-of-river schemes are associated with a smaller footprint of land-cover conversion compared with large storage schemes. However, guidance to decrease negative impacts on biological diversity often is not provided. Moreover, impact assessments are rarely required for small individual dams, although they could be associated with considerable cumulative impacts on particular watersheds and species. For example, although few studies of their local and downstream ecological effects have been conducted, small dams can be associated with altered hydrological and sediment flow regimes. These alterations may cause sediment scarcity downstream, limit the dispersal of organisms, serve as barriers to migratory fish, and reduce oxygen concentrations and increase water temperatures, decreasing habitat quality for some endemic fish species [23].

Large Recirculating Aquaculture Systems

Intensive aquaculture is associated with high levels of water use, local environmental pollution [24], and loss of coastal ecosystems. Recirculating aquaculture systems (RAS) circulate water around tanks that hold cultured species, treat the water to maintain high quality, and reduce water flow through the system. Such enclosed systems usually have fewer direct environmental effects compared with traditional aquaculture. Small RAS have been successful, notably in freshwater ecosystems [25], but there is a growing trend in the development of large saltwater systems (e.g., for salmon production) capable of producing tens of thousands of tonnes of fish annually (e.g., [26]). Intensive RAS have 1–3% of the water demand of throughflow aquaculture [25]. Recent technological advances linked RAS to aquaponics systems that utilise the high-nutrient effluents. However, implementation of RAS is constrained by the need for high capital investment, with payback periods of ~8 years or more. Establishment of extensive RAS could increase the sustainability of food production from aquaculture and may reduce the risks from pollution and parasite release often associated with throughflow aquaculture. Other challenges, such as impacts from the sources of aquaculture feed, and energy requirements, have not yet been addressed.

Genetically Modified Fungus Kills Malaria-Carrying Mosquitoes

The evolution of insecticide resistance by mosquito species that serve as malaria vectors means that development of novel approaches to limit the spread of malaria-carrying mosquitos is becoming necessary to reduce prevalence of the disease [27]. Entomopathogenic fungi are promising in this regard, but their use is limited by the time lag in host mortality and the high dose required for infection. The fungus *Metarhizium pingshaense*, which naturally infects a mosquito species capable of carrying malaria, was genetically modified recently to produce a toxin derived from spider venom. The modified pathogen killed mosquitoes faster and at lower spore doses compared with the unmodified fungus. Its mode of action differs from that of pyrethroids, enabling the two control agents to act synergistically [28]. In a demonstration village that was isolated for bioinsecticide application, the modified fungus led to increased fungal lethality and likelihood of mosquitoes being eliminated locally. Although fungal insecticides are not new [29], successful translation to semifield conditions, with high mortality of the mosquito population, is novel [27]. Reduced use of insecticides, particularly in wetlands close to urban centres, could have ecological benefits. However, any change in the host range of the modified pathogen could affect nontarget organisms.

Use of Artificial Wombs and Ectogenesis in Mammalian Conservation

Biobags are artificial wombs that allow partial ectogenesis (foetal development outside the body of the mother). Lamb foetuses that had partially developed *in utero* were successfully transferred into biobags to continue their gestation to full term *ex utero* [30]. Biobags contain the foetus, a pumpless oxygenator circuit, and a synthetic amniotic fluid, and are designed for human neonatal intensive care. Although human testing is several years off, biobags could be used by parents to overcome

troubled pregnancies or to aid premature babies [31]. Artificial wombs also could assist conservation breeding programs for threatened mammals, particularly if the technology matures to support complete ectogenesis (embryos created with *in vitro* fertilisation and gestated entirely within an artificial womb), and the rate of production of mammalian offspring increase markedly. Complete ectogenesis will raise numerous practical, ethical, and financial challenges [32]. These challenges include developmental or behavioural problems for embryos not biologically attached to a mother. Additionally, understanding of the role of the placental environment on gene activation in embryos and its effects on immunity in offspring is limited.

International Growth of Traditional Asian medicine

Traditional medicine, largely centred on ancient Asian medicine, was included for the first time in the International Classification of Disease [33], endorsed by the World Health Assembly, in May 2019. Inclusion has been viewed as an endorsement of traditional medicine and may accelerate already increasing patterns of its use [34]. Traditional medicine applies diagnostics and treatments that often have few similarities to western medicine, although efforts to expand the use of randomised controlled trials are increasing [34]. The Government of China is investing in the promotion of traditional Asian medicine through methods including health tourism and international market expansion. China has established some 25 traditional Asian medicine institutes in a range of cities, and more will be launched as a major component of the Belt and Road Initiative [34]. Sales of traditional Asian medicine products, such as herbal medicines, are growing: sales in Belt and Road countries grew by 54% in 2016 and 2017, reaching US\$295 million annually [34]. The growth of traditional Asian medicine will increase demand for ingredients that include some plant and animal species already endangered by harvest for international trade [35]. Increased connectivity across the Belt and Road route may increase access to formerly inaccessible wildlife populations and increase trafficking of wildlife for use in medicine [36].

Rise of Blockchain Companies with Hidden Owners

Blockchain, the distributed ledger technology, is revolutionising traditional corporate structures. The distributed consensus mechanism fundamental to blockchain technology ensures the security, integrity, and performance of the network. Although blockchain technology is secure in principle, applications running on top of them, such as self-enforcing contracts, can be subject to coding errors or security vulnerabilities just like any other software, and security systems will need to evolve to meet the needs of different applications. Companies that use blockchain need not adhere to a conventional management or financial infrastructure. In particular, self-enforcing agreements embedded in computer code may change how energy resources [37] and other natural assets are owned and managed. For example, in Berlin, two artists launched terra0, a blockchain experiment in which a forest autonomously sells its trees, harvests timber, and eventually uses the accumulated capital to buy itself and become a self-owned economic unit [38]. The rapid development in these self-enforcing contracts could enable companies to confirm resource streams and commit to future actions well beyond current political and regulatory timeframes and without the need for any physical or identifiable company. These resource transactions, which are secure, immutable, and verifiable, may strengthen environmental governance. They can also be used to reinforce entitlements to long-term resource extraction or even to substantiate indigenous land rights. Additionally, it may be necessary to clarify or amend existing laws, for example, to recognise the use of distributed ledgers as records of ownership [39].

Ecocide as an Internationally Recognised Crime

Currently, the International Criminal Court (ICC), governed by the Rome Statute, can prosecute individuals and states for ‘widespread, long-term and severe’ environmental destruction [40], but only in certain circumstances, such as during conflict or when the destruction has serious humanitarian consequences. However, as currently drafted, the Rome Statute contains no provisions to protect non-human inhabitants of a given territory or indigenous or cultural rights; neither does the Rome Statute cover environmental loss, damage, or destruction during peacetime. Legal scholars (e.g., [41]) argue that this international law must change to allow the crime of ecocide. Ecocide is defined as ‘the

extensive damage, destruction or loss of ecosystems of a given territory...to such an extent that peaceful enjoyment by the inhabitants of that territory has been severely diminished' [42]. Research to establish forensic standards for admissible evidence of ecocide is advancing [43]. Efforts to encourage the ICC to recognize ecocide focus on either the inclusion of ecocide under crimes against humanity or the establishment of ecocide as a distinct Crime Against Peace. Such changes to international law would enable individuals, states, and perhaps corporations (not possible under the current statute) to be prosecuted for extensive land-cover modification, pollution, and even contributing to climate change.

New United Nations Legal Principles to Reduce the Environmental Impact of Armed Conflict

In July 2019, the International Law Commission of the United Nations (UN) adopted draft principles on the protection of the environment in relation to armed conflicts. Its action represented a major step toward conclusion of a process to review the international law that was passed in 2013ⁱⁱ. This new international legislation, which would oblige states and other actors to protect the environment during periods of armed conflict, could have substantial effects on species and ecosystems worldwide, given the environmental impacts of modern warfare [44]. The detrimental effects of armed conflicts on species and ecosystems can be direct, through tactical military operations, or indirect, through their effects on institutions, human migration, and economies [45]. At the same time, degradation of ecosystems could cause armed conflict and human migration, as demonstrated in the Sahel, where the degradation of wetlands has resulted in conflict over resources [46]. Given that the most common links between armed conflicts and ecological responses are subsequent changes to institutions, societies, and economies rather than direct impacts of conflict [45], the set of principles of the new legislation applicable after armed conflictⁱⁱ are particularly relevant; draft principle 14 encourages parties to an armed conflict to address environmental restoration during the peace process.

New Regulations Jeopardise Net Neutrality

Conservation relies on the communication of knowledge to, and engagement with, the general public [47]. Digital tools, such as blogging and social media, provide authors who have diverse perspectives or agendas [48] with unprecedented access to the general public, often with no filter between author and audience. Easy and equal access to all websites and types of data underlies net neutrality (i.e., all internet data are treated equally by internet service providers). Without network neutrality, providers may block or restrict access to pages and content on the basis of their corporate policies or interests, allowing accurate information to be restricted or distorted. Such blocking already affects content in some countries, but network neutrality is now under a general threat: it was repealed in the USA (vote of the Federal Communications Commission, July 2018) [49], and other countries could follow suit. Loss of network neutrality could have major effects on conservation and climate regulation if access to accurate or sensitive information is denied or biased or, conversely, if guaranteeing such access is prioritised.

Concluding Remarks

The pace of data exchange and the volumes of information available continue to accelerate, with the quantity of data available on the internet doubling every 2 years. The challenges of horizon scanning include not only the process of searching for issues, but also understanding whether such information is sufficiently unknown to a given community to warrant inclusion. Some topics we initially considered, such as the effect of nocturnal harvesting of olives on roosting songbirds, subsequently were widely reported by the press and, consequently, we considered them too well known to include in this year's scan. However, this does not mean that the issues are any less relevant to policy-making and conservation.

Last year, we identified the importance of national government policy or economic decision making in driving global environmental impacts [50], a theme echoed by two of our issues this year, the potential impact of the new EU policy on the derivation of energy from wood and the repealing of network

neutrality in the USA. Another two of our issues this year are based on the potential for decisions by global institutions, such as the ICC and UN, to drive change, while a third highlighted the potential impact of decisions by the World Health Assembly. Whether these international instruments determine global environmental trajectories ultimately will depend on the extent to which countries become signatories and then incorporate decisions into national legislation. This challenge was exposed by the high-profile public protests over climate change during 2019, which highlighted the failure of national governments to realise their international commitments. With several major international meetings focused on the protection, restoration, and sustainable use of biological diversity, such as the International Union for Conservation of Nature (IUCN) World Conservation Congress and the 15th meeting of the Conference of the Parties to the Convention on Biological Diversity, scheduled for 2020, this tension between national decision making and international commitment is likely to persist. In particular, new conservation agendas, the European green deal, and indicators and targets to 2030 and beyond will focus attention on addressing future threats and opportunities, such as those we seek to identify through horizon scanning.

There is increasing awareness of the role that unconscious bias can have in decision making [51]. We recognise that the majority of the people participating in the horizon scan are residents of the UK, Europe, or the USA, which may create implicit biases in both the scope of the topics considered and the levels of awareness of the issues. We aim to incorporate a wider range of perspectives, particularly from the global south, into future horizon scans. Solicitation of topics could better reflect scientific knowledge and ways of knowing from individuals and societies in regions where biological diversity is frequently stated to be at the greatest level of risk. Nevertheless, we believe that the topics identified this year may affect global conservation, and we hope that this paper will prompt discussion and new research.

Strategic foresight can be defined as 'The systematic examination of potential threats, opportunities and likely future developments which are at the margins of current thinking and planning. [The research] may explore novel and unexpected issues, as well as persistent problems or trends. Overall, it is intended to improve the robustness of policies and the evidence base' [52]. Strategic foresight methods are relevant to experts in any discipline. Horizon scanning is one of many methods used in strategic foresight research, which also includes, but is not limited to, risk prioritisation, trend extrapolation, scenario development, backcasting, and stress testing [53]. Although no single method is applicable in all situations, horizon scanning provides the foundations on which all subsequent foresight research is based. The initial stages of horizon scanning are highly inclusive and, by definition, emphasise novelty, but are not intended to prioritise topics that are well known or proven. Horizon scanning aims to identify and explore new insights and evidence regardless of whether they are consistent with existing trends and developments [54]. The focus of horizon scanning is to provide an evidence base of current knowledge that suggests the potential for future change. Horizon scanning can be used to identify, assess, and understand gaps in knowledge, identify potential opportunities and risks, and inform research programming and resource allocation. As a tool for decision makers, horizon scanning aims to support strategic activities [55]. The insights identified through horizon scanning are intended to stimulate multiparty discussion and debate, leading to potentially collaborative solutions to complex issues [56].

Acknowledgements

This exercise was coordinated by the Cambridge Conservation Initiative, and funded by the Natural Environment Research Council and RSPB. We are grateful to everyone who submitted ideas to the exercise. We thank Stefano Basso (effects of small hydropower systems on riverine ecosystems), Jason Dinsdale (use of artificial wombs and ectogenesis in mammalian conservation and rise of block-chain companies with hidden owners), Sarah Durant (new United Nations legal principles to reduce the environmental impact of armed conflict), Lizzie Duthie (new regulations jeopardise net neutrality), Norah Eddy (global declines of kelp forest), Robert Jones (large recirculating aquaculture systems), and Catherine Julliot (policy incentives for derivation of energy from wood) for suggesting some issues that were included in the final scan. W.J.S. is funded by Arcadia.

Resources

ⁱwww.pearltrees.com

ⁱⁱ<http://legal.un.org/ilc/sessions/71/>

References

- Sutherland, W.J. et al. (2019) Ten years on: a review of the first global conservation horizon scan. *Trends Ecol. Evol.* 34, 139–153
- Sutherland, W.J. et al. (2011) Methods for collaboratively identifying research priorities and emerging issues in science and policy. *Methods Ecol. Evol.* 2, 238–247
- Mukherjee, N. et al. (2015) The Delphi technique in ecology and biological conservation: applications and guidelines. *Methods Ecol. Evol.* 6, 1097–1109
- Sutherland, W.J. et al. (2010) A horizon scan of global conservation issues for 2010. *Trends Ecol. Evol.* 25, 1–7
- Danziger, S. et al. (2011) Extraneous factors in judicial decisions. *Proc. Natl. Acad. Sci. U. S. A.* 108, 12001–12006
- Sharma, A. et al. (2019) Commercial application of cellulose nano-composites – a review. *Biotechnol. Rep.* 21, e00316
- Walker, S. et al. (2015) *An Analysis of UK Biomass Power Policy, US South Pellet Production and Impacts on Wood Fiber Markets, RISI*
- Giacomini, J. et al. (2018) Medicinal value of sunflower pollen against bee pathogens. *Sci. Rep.* 8, 10
- LoCasio, G. et al. (2019) Pollen from multiple sunflower cultivars and species reduces a common bumblebee gut pathogen. *R. Soc. Open Sci.* 6, 9
- Goulson, D. et al. (2018) The impacts of predators and parasites on wild bumblebee colonies. *Ecol. Entomol.* 43, 168–181
- Koch, H. et al. (2019) Flagellum removal by a nectar metabolite inhibits infectivity of a bumblebee parasite. *Curr. Biol.* 29, 1–7
- Oakes, V.J. et al. (2019) *Theileria orientalis* Ikeda genotype in cattle, Virginia, USA. *Emerg. Infect. Dis.* 25, 1653–1659
- Raghavan, R.K. et al. (2019) Potential spatial distribution of the newly introduced long-horned tick, *Haemaphysalis longicornis* in North America. *Sci. Rep.* 9, 498
- Heath, A.C.G. (2016) Biology, ecology and distribution of the tick, *Haemaphysalis longicornis* Neumann (Acari: Ixodidae) in New Zealand. *N. Z. Vet. J.* 64, 10–20
- Arafeh-Dalmau, N. et al. (2019) Extreme marine heatwaves alter kelp forest community near its equatorward distribution limit. *Front. Mar. Sci.* Published online August 23, 2019. <https://doi.org/10.3389/fmars.2019.00499>
- Krumhansl, K.A. et al. (2016) Global patterns of kelp forest change over the past half-century. *Proc. Natl. Acad. Sci. U. S. A.* 113, 13785–13790
- Filbee-Dexter, K. and Wernberg, T. (2018) Rise of turfs: a new battlefield for globally declining kelp forests. *Bioscience* 68, 64–76
- Parkinson, C.L. (2019) A 40-y record reveals gradual Antarctic sea ice increases followed by decreases at rates far exceeding the rates seen in the Arctic. *Proc. Natl. Acad. Sci. U. S. A.* 116, 11414–11423
- Thompson, D.W.J. et al. (2011) Signatures of the Antarctic ozone hole in Southern Hemisphere surface climate change. *Nat. Geosci.* 4, 741–749
- Seviour, W.J.M. et al. (2019) The Southern Ocean sea surface temperature response to ozone depletion: a multi-model comparison. *J. Climate* 32, 5107–5121
- Couto, T.B. and Olden, J.D. (2018) Global proliferation of small hydropower plants—science and policy. *Front. Ecol. Environ.* 16, 91–100
- Jumani, S. et al. (2018) Fish community responses to stream flow alterations and habitat modifications by small hydropower projects in the Western Ghats biodiversity hotspot, India. *Aquat. Conserv.* 28, 979–993
- Benejam, L. et al. (2016) Ecological impacts of small hydropower plants on headwater stream fish: from individual to community effects. *Ecol. Freshw. Fish* 25, 295–306
- Diana, J.S. (2009) Aquaculture production and biodiversity conservation. *Bioscience* 59, 27–38
- Bregnballe, J. (2015) *A Guide to Recirculation Aquaculture*, FAO and EUROFISH International Organisation
- White, C. (2017) *Atlantic Sapphire Building USD 350 Million Land-Based Salmon Farm in Miami*, SeafoodSource
- Lovett, B. et al. (2019) Transgenic *Metarhizium* rapidly kills mosquitoes in a malaria-endemic region of Burkina Faso. *Science* 31, 894–897
- Bilgo, E. et al. (2018) Transgenic *Metarhizium pingshaense* synergistically ameliorates pyrethroid-resistance in wild-caught, malaria-vector mosquitoes. *PLoS One* 13, e0203529
- Windley, M.J. et al. (2012) Spider-venom peptides as bioinsecticides. *Toxins (Basel)* 4, 191–227
- Partridge, E.A. et al. (2017) An extra-uterine system to physiologically support the extreme premature lamb. *Nat. Commun.* 8, 15112
- Romanis, E.C. (2018) Artificial womb technology and the frontiers of human reproduction: conceptual differences and potential implication. *J. Med. Ethics* 44, 751–755
- Harrop, F. (2019) Are we ready to confront ethics of artificial wombs? *Herald Net*
- WHO. (2018) *ICD-11 for Mortality and Morbidity Statistics (ICD-11 MMS) 2018 Version*, WHO
- Cyranoski, D. (2018) Why Chinese medicine is heading for clinics around the world. *Nature* 561, 448–450
- Byard, R.W. (2016) Traditional medicines and species extinction: another side to forensic wildlife investigation. *Forensic Sci. Med. Pat.* 12, 125–127
- Sutherland, W.J. et al. (2018) A 2018 horizon scan of emerging issues for global conservation and biological diversity. *Trends Ecol. Evol.* 33, 47–57
- Salmerón-Manzano, E. and Manzano-Agugliaro, F. (2019) The role of smart contracts in sustainability: worldwide research trends. *Sustainability* 11, 3049
- Cassauwers, T. (2018) How artists are bringing blockchain to their neck of the woods. *Ozymandias*
- European Bank (2018) *Smart Contracts, Legal Frameworks, and Proposed Guidelines for Lawmakers*, European Bank
- ICC (1998) *Rome Statute of the International Criminal Court*, ICC, (Last amended 2010).
- Mwanza, R. (2018) Enhancing accountability for environmental damage under international law: ecocide as a legal fulfilment of ecological integrity. *Melb. J. Int. Law* 19, 586–613

42. Higgins, P. (2015) *Eradicating Ecocide*, Shephard-Walwyn
43. Ahmed, N. (2017) Proof of ecocide: towards a forensic practice for the proposed international crime against the environment. *Archaeol. Environ. Forensic Sci.* 1, 139–147
44. Brito, J.C. et al. (2018) Armed conflicts and wildlife decline: challenges and recommendations for effective conservation policy in the Sahara-Sahel. *Conserv. Lett.* 11, e12446
45. Gaynor, K.M. et al. (2016) War and wildlife: linking armed conflict to conservation. *Front. Ecol. Environ.* 14, 533–542
46. Wetlands International. (2017) *Water Shocks: Wetlands and Human Migration in the Sahel*, Wetlands International
47. Thaler, A.D. et al. (2012) Digital environmentalism: tools and strategies for the evolving online ecosystem. In *Environmental Leadership: A Reference Handbook* (Rigling Gallagher, D. ed), pp. 364–372, Sage
48. Graf, H. (2016) OpenBook. *The Environment in the Age of the Internet*
49. Solomone, S. (2018) Regulate social media? It's a bit more complicated than that. *The Conversation*
50. Sutherland, W.J. et al. (2019) A horizon scan of emerging issues for global conservation in 2019. *Trends Ecol. Evol.* 34, 83–94
51. Newell, B.R. and Shanks, D.R. (2014) Unconscious influences on decision making: a critical review. *Behav. Brain. Sci.* 37, 1–19
52. DEFRA (2002) *Horizon Scanning and Futures*, DEFRA
53. Gov, U.K. (2017) *The Futures Toolkit: Tools for Futures Thinking and Foresight Across UK Government*, Government Office for Science
54. Pillkahn, U. (2008) *Using Trends and Scenarios as Tools for Strategy Development: Shaping the Future of Your Enterprise*, Wiley-VCH
55. Palomino, M.A. et al. (2012) Web-based horizon scanning: concepts and practice. *Foresight* 14, 355–373
56. Amanatidou, E. et al. (2012) On concepts and methods in horizon scanning: lessons from initiating policy dialogues on emerging issues. *Sci. Publ. Pol.* 39, 208–221
57. MarketsandMarkets, Nanocellulose Market by Type (Microfibrillated Cellulose, Cellulose Nanocrystals), Application (Pulp & Paper, Composites & Packaging, Biomedical & Pharmaceuticals, Electronics & Sensors), and Region (Europe, North America, APAC) – Global Forecast to 2023, MarketsandMarkets, 2019