



Litter decomposition above the treeline in alpine regions: A mini review

Monika Rawat ^a, Annika K. Jägerbrand ^b, Yang Bai ^c, Juha M. Alatalo ^{d,*}

^a School of Environment and Natural Resources, Doon University, Dehradun, 248001, India

^b Department of Environmental and Biosciences, Rydberg Laboratory of Applied Science (RLAS), School of Business, Engineering and Science, Halmstad University, P.O. Box 823, SE-301 18 Halmstad, Sweden

^c Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Mengla, Mengla, Yunnan, 666303, China

^d Environmental Science Center, Qatar University, PO Box 2713, Doha, Qatar

ARTICLE INFO

Keywords:

Carbon cycling
Decomposers
High altitude
Quality
Plant litter
Soil organic matter

ABSTRACT

Litter decomposition is a key driver of ecosystem processes and carbon cycling. Decomposition rate is influenced by numerous factors, such as temperature, humidity, litter properties, soil properties, and properties of soil fauna/microbial communities. The aim of this review was to summarize current knowledge on litter decomposition above the treeline in alpine regions worldwide and identify: I) factors that have been studied in great detail, II) factors that have been less intensively investigated, III) geographical regions that have been less well studied, and IV) factors with consistent or inconsistent effects on decomposition. The review showed inconsistent results for all factors covered by two or more studies regarding their effect on decomposition rate (positive, negative, no effect), usually a result of interactions between factors. Studies examining one or several factors in the physical environment (i.e., altitude, experimental warming, microclimate, snow cover and soil moisture) were most common, while studies on different aspects of resource quality were the second most common. The impacts of trophic interactions on soil microbes and fauna were less frequently studied. Europe and Asia were the best-represented regions, in terms of number of studies and geographical distribution, while there were no studies from Africa and very few from South America and Australia. North American studies were all from Colorado, and those from Asia were all from China. In order to obtain better global representation, there is a need for studies in Africa, South America, and Australia. There is also a need for more studies to explain the large variation in responses of litter decomposition rates to different influencing factors in alpine environments. Future research should focus on interactions between different factors and on experiments testing specific relationships, such as the potential interaction between temperature and soil moisture and its effect on litter decomposition above the treeline in alpine regions.

1. Introduction

Litter decomposition is an important ecological process for determining nutrient cycling and ecosystem functioning (Bryant et al., 1998; Gavazov, 2010). It is mainly influenced by three important factors: resource quality (e.g. plant spp composition), the physical environment (e.g. humidity, soil structure) and soil organisms (e.g. fungus, worms) (Hartig, 1878; Krishna and Mohan, 2017; Lindquist, 1941; Swift et al., 1979; Waksman et al., 1928). During the decomposition process, organic matter is broken down through leaching, fragmentation and chemical reduction (Chapin et al., 2011) building up soil organic matter (SOM) content (Gavazov, 2010) and regulate air carbon dioxide (CO₂) (Novara et al., 2015). As a result, litter decomposition is a major source for the

return of organic matter and nutrients release to the soil. This process enhances soil functions, control microenvironment conditions and influence microbial communities and its growth, mainly in the litter influenced upper soil horizon (Freschet et al., 2013; Zheng et al., 2018).

In alpine regions, it is estimated that more than 90% of the total carbon pool is found in SOM, and that live biomass contains only 3–5% of total carbon (Körner, 1999). This is due to the short life span of leaves (often one season) compared with the time needed for decomposition of leaves, which may take 2–10 years, depending on plant functional group (forbs, sedges, or evergreen shrubs), with forbs decomposing faster than evergreens (Seastedt et al., 2001). In addition, the decomposition process tends to be slower in colder conditions and this can cause nutrient limitation (Aerts, 2006). Thus, both total soil nitrogen (N) and

* Corresponding author.

E-mail address: jatalo@qu.edu.qa (J.M. Alatalo).

<https://doi.org/10.1016/j.actao.2021.103775>

Received 18 February 2021; Received in revised form 4 August 2021; Accepted 18 August 2021

Available online 26 August 2021

1146-609X/© 2021 The Authors.

Published by Elsevier Masson SAS. This is an open access article under the CC BY license

(<http://creativecommons.org/licenses/by/4.0/>).

plant-available N decrease with altitude (Huber et al., 2007). A previous global study with data from 33 experimental warming experiments in cold biomes found that direct climate effects and plant growth forms were the main contributors to litter decomposition rate, while within-species variation made a much smaller contribution (Cornelissen et al., 2007). Climate change could therefore affect decomposition indirectly, through causing changes in species composition in plant communities (Cornelissen et al., 2007; Mori et al., 2020). Similar conclusions were reached in a meta-study on data from 336 sites using the Teabag Index (Keuskamp et al., 2013), which found that litter quality and climate were the main controlling factors for initial decomposition rate (Djukic et al., 2018). Multiple other factors can also influence decomposition. These include e.g., grazing (Sun et al., 2018), soil fauna (Lindquist, 1941; Wang et al., 2009), animal dung deposition (Liang et al., 2019), soil fertilization (Song et al., 2019; Soudzilovskaia et al., 2007), soil microbial communities (Hartig, 1878; Lipson et al., 2000; Štursová et al., 2020; Waksman et al., 1928), snow cover (Baptist et al., 2010), soil moisture (Soudzilovskaia et al., 2007) and metal concentrations in soil (Sarnecki et al., 2020).

High-alpine regions (above the treeline) are sensitive to environmental changes (Alatalo et al., 2017a), so it is likely that climate change in coming decades will have an impact on alpine vegetation above the treeline (Schwager and Berg, 2019). This may have significant consequences for the decomposition process and for carbon and nutrient cycling in high-alpine regions.

In November 2019 we searched six literature databases (Web of Science, Pub Med, Science Direct, Pro-Quest, Scopus, and Google scholar), for “alpine” and “decomposition” in titles, abstracts and keywords. In addition, we checked reference lists and recent citations to find additional papers not identified by our initial search. Our selection criteria included: (i) articles published in peer-reviewed journals, and (ii) conducted above the treeline in alpine regions. The literature review

was restricted to studies published in English.

The aim of this review was to summarize current knowledge on litter decomposition above the treeline in alpine regions worldwide. The focus was on broader factors that may affect litter decomposition and potential evidence of their influence on litter decomposition. In addition to study location, we analyzed the direction (positive, negative, or no effect) of responses to the factors studied and the number of studies on different factors. We grouped relevant studies into the categories “physical environment” (altitude, snow cover, soil moisture, soil pH, temperature, etc.), “resource quality” (litter quality, lignin/N content, organic content, vegetation type, etc.), and “trophic interactions” (soil fauna, microorganisms, grazing). Specific objectives of the review were to identify: I) factors affecting litter decomposition that have been studied in great detail; II) factors that have been less intensively investigated, III) geographical regions that have been less well studied, and where future studies should be conducted to verify large-scale patterns; and IV) factors for which consistent or inconsistent effects on litter decomposition are reported in different studies. The overall objective was to identify factors and regions that should be prioritized in future research.

2. Factors studied and their impact on litter decomposition

2.1. Physical environment

2.1.1. Altitude

Conflicting effects of altitude on litter decomposition have been reported (Fig. 1). Three studies have found negative effects of altitude on decomposition rate (Andreyashkina, 2008; Couëteux et al., 2002; Schinner, 1982). Two have found positive effects (Andreyashkina and Peshkova, 2003; Wood, 1970) or different patterns (positive and no effect) of effect between two mountain ranges (Withington and Sanford,




Fig. 1. Network diagram showing the linkages between litter decomposition with physical environment, resource quality and trophic interactions in alpine ecosystems. The width of the lines reflects the number of papers showing that linkage. The direction of the effect is indicated positive (+), negative (−), no relationships (0) when the result from the study is unclear. Arrows indicating direct effects and indirect effects.

2007), while one has found no effect (Duboc et al., 2012). Several indirect explanations have been suggested for the positive effects of altitude. These include increased soil moisture content (Withington and Sanford, 2007; Wood, 1970), soil organisms (Wood, 1970), better aeration, higher warming of soils, and higher proportion of herbaceous plants (Andreyashkina and Peshkova, 2003). The negative effect of altitude on decomposition has been suggested to be caused indirectly by decreasing microbial activity with altitude (Schinner, 1982), and e.g., a decrease in temperature with altitude in the tropical Andes (Coûteaux et al., 2002).

2.1.2. Experimental warming, microclimate, snow cover, and soil moisture

Experimental warming in alpine ecosystems has been found to result in positive effects on litter decomposition (Aerts, 2006; Carbognani et al., 2014; Gavazov, 2010; Luo et al., 2010; Lv et al., 2020; Sjögersten and Wookey, 2004), or negative effects (Cornelissen et al., 2007; Hong et al., 2021; Liu et al., 2021; Sarneel et al., 2020) (Fig. 1). Microclimate has also been found to have positive effects (Chen et al., 2018; Shaw and Harte, 2001) or negative effects linked to the interaction with grazing pressure, soil moisture content, and changes in vegetation structure (Vaieretti et al., 2018) (Fig. 1). Similarly, studies on snow cover have found negative (Baptist et al., 2010), positive (O'Lear and Seastedt, 1994), or no effect (Coûteaux et al., 2002; Huang et al., 2016; Zeidler et al., 2014) on litter decomposition (Fig. 1). Snow cover also affects soil moisture. Drought is reported to have a negative influence on decomposition (Schinner, 1983).

The majority of studies on soil moisture (78%, or seven out nine studies) have found a positive effect on decomposition (Bryant et al., 1998; Liu et al., 2021; Lv et al., 2020; Sarneel et al., 2020; Sjögersten and Wookey, 2004; Withington and Sanford, 2007; Wood, 1970) (Fig. 1). However, one study has reported a negative impact (Schinner, 1983) and one no effect (Coûteaux et al., 2002). The positive effect of soil moisture on litter decomposition is hypothesized to be caused by a water deficit in upper soil and litter layers in mesic/dry tundra (Sjögersten and Wookey, 2004), and large mass loss due to leaching in the early stage of decomposition (Bryant et al., 1998). In fact, it has been suggested that decomposition is primarily constrained by soil moisture and secondarily by temperature (Prescott, 2010). Thus, warming can be expected to have contrasting effects on decomposition of litter on drier and wetter alpine soils, with drier soils being more negatively affected by warming accompanied by drying of soils, while higher temperature might have a greater effect at wetter sites (Liu et al., 2021; Sarneel et al., 2020). However, the negative effect reported for both drought and excess water suggests that too much or too little water can impair the decomposition process (Schinner, 1983).

Soil physico-chemical properties (pH and carbon/nitrogen (C/N) ratio) have been found to have positive effects on litter decomposition at high-alpine sites (Duboc et al., 2012; Schinner, 1983; Sjögersten and Wookey, 2004; Wang et al., 2020) (Fig. 1). The cause for this positive relationship has been hypothesized to be due to adaptation of microbial populations to low quality litter with increasing soil C:N ratio (Duboc et al., 2012). One study found a negative effect of soil metal concentrations and moisture on decomposition rate (Sarneel et al., 2020) (Fig. 1). In another study, soil C/N ratio had a positive effect during the first year of decomposition, while soil pH was important during the second year (Duboc et al., 2012). This might have been caused by indirect effects of soil properties on microbial communities. Experimental soil acidification, lowering the pH, has been found to have a negative effect on root decomposition in Tibet, causing an increase in below-ground biomass (Wang et al., 2020). In addition, other properties of the soil have been studied, e.g., it has been found that soil structural stability is increased by the presence of earthworms (Seeber et al., 2006).

The physical environment broadly influences litter decomposition in the global carbon and nutrient cycles (Murphy et al., 1998). Climate factors such as temperature, precipitation, and evapotranspiration control decomposition mechanisms to a large extent and have a

significant impact on carbon and nutrient cycling (Giweta, 2020; Zhang et al., 2008; Zhou et al., 2008). Thus, an improved understanding of how physical factors control/affect decomposition processes could lead to better predictions on how global warming may affect nutrient cycling and ecosystem functioning in the future (Song et al., 2014).

2.2. Resource quality

Studies on resource quality in alpine regions report contrasting effects on litter decomposition (Fig. 1). Positive responses of litter quality, i.e., litter N, lignin, C/N ratio, nitrogen/phosphorus (N/P) ratio, cellulose content, have been found (Carbognani et al., 2014; Drewnik, 2006; Hong et al., 2021; Wang et al., 2021). Negative responses (e.g., of lignin/nitrogen ratio and phenolics) (Shaw and Harte, 2001; Steltzer and Bowman, 2005; Wang et al., 2021) and negligible effects of litter chemistry (Soudzilovskaia et al., 2007) have also been reported. Leaf trait diversity can have a positive effect (Song et al., 2019). Species and vegetation type specific litter quality parameters have been found to influence decomposition in different ways (Andreyashkina, 2008; Andreyashkina and Peshkova, 2003; Carbognani et al., 2014; Sarneel et al., 2020; Soudzilovskaia et al., 2007; Wood, 1970; Zeidler et al., 2014; Zheng et al., 2020), with forbs decomposing at a faster rate than graminoids (Soudzilovskaia et al., 2007). In a study that included four species (three *Eucalyptus* spp., one *Poa* sp.), but no chemical analysis of the plant material, it was found that *Eucalyptus* litter decomposed faster than *Poa* litter at higher altitude, where *Eucalyptus* does not naturally occur (Wood, 1970). Experimental fertilization has also been found to result in contrasting responses, e.g., in one study fertilization significantly increased N and P concentrations in plant litter, but had no effect on decomposition (Soudzilovskaia et al., 2007). Similarly, no effect of fertilization was found in a study in the Rocky Mountains, Colorado (Bryant et al., 1998). However, a study in Switzerland found that soil fertilization increased decomposition rate (Arnold and Hirschel, 1997). A study in Tibet found that dung deposition had a positive effect on litter decomposition (Liang et al., 2019). In general, higher N and P concentrations are believed to have a positive effect on decomposition (Cornwell et al., 2008). However, it has also been suggested that other traits, such as toughness, lignin concentration, or cellulose concentration, might be of greater importance for decomposability than high N or P concentrations in litter (Soudzilovskaia et al., 2007).

Decomposition is strongly influenced by the physical environment, but litter quality aspects such as nitrogen, lignin, polyphenol, C/N ratio, and lignin/nitrogen ratio can also play an important role in influencing the decomposition process (Giweta, 2020; Krishna and Mohan, 2017; Zheng et al., 2020). Decomposing plant material passes through three different stages, i.e., nutrient release, net immobilization, and net release (Wang et al., 2021), which control the functioning of the forest ecosystem (Zhou et al., 2008). Thus, litter quality can serve as a predictor of decay rate and is an important component of the biogeochemical cycle (Giweta, 2020).

2.3. Trophic interactions

Studies on the effect of trophic interactions on decomposition have also reported conflicting results (Fig. 1). Most studies have found that soil fauna have a positive effect on decomposition (Kitz et al., 2015; Ma et al., 2019; O'Lear and Seastedt, 1994; Wang et al., 2018). One study found contrasting results for different species of macro-decomposers, with primary decomposers increasing decomposition and secondary decomposers having no or negative effects (Seeber et al., 2006). Contrary to expectations, that study also found that decomposers preferred to feed on low-quality litter with high lignin and tannin content (Seeber et al., 2006). Grazing has been found to have both positive (Luo et al., 2010; Sun et al., 2018) and negative (Vaieretti et al., 2018) effects on litter decomposition on alpine soils. The negative effect is reported to be caused by grazing interacting with vegetation, microclimate, and soil

moisture, with reduced grazing leading to taller vegetation, higher soil moisture content, and ultimately increased decomposition rates (Vaieretti et al., 2018).

Trophic interactions are crucial for decomposition processes in many ecosystems. Soil fauna in particular increase plant litter decomposition in ecosystems, but their distribution varies with climate conditions and habitats (Phillips et al., 2019). Furthermore, the results on earthworm global distribution tend to differ from what is normally found for aboveground taxa, suggesting that aboveground and belowground biodiversity may not follow the same broad distribution pattern (Phillips et al., 2019). Macrofauna, such as millipedes and earthworms, frequently act as detritivores, while soil microarthropods, such as collembolans and oribatid mites, impact litter decomposition by modifying the activity and composition of saprotrophic fungal communities (Schaefer et al., 2009; Zhou et al., 2020). How climate change will affect alpine soil fauna, and their contribution to alpine decomposition, is still unclear, as there are still only a few experimental climate change studies on soil fauna in alpine regions (Alatalo et al., 2015, 2017b; Hågvar and Klanderud, 2009; Roos et al., 2020).

2.4. Geographical distribution of studies

Regarding the geographical distribution of published studies on litter decomposition above the treeline included in this review, we found that the majority of the studies were carried out in Europe and Asia (Austria, Norway, Sweden, Italy, France, Russia, Switzerland, and China), and five studies were performed in North America (all USA) (Fig. 2). Two were from South America (Argentina and Venezuela) and one from Australia. The three global studies included compiled data from 18 sites with Arctic and Alpine climate change experiments. Of these, one study included data from 10 sites in Canada, China, Finland, Italy, Japan, Norway, Russia, Sweden, and USA (Cornelissen et al., 2007) and one included data from climate change experiments at one alpine site in the USA and three alpine tundra sites in Norway and Sweden (Aerts, 2006). The third meta-study was a conventional review paper on the potential impact of climate change on alpine plant litter decomposition (Gavazov, 2010). While it is likely that we overlooked some studies, it is unlikely that this would affect the general geographical pattern observed to any great degree.

3. Conclusions

This review showed that factors relating to the physical environment and resource quality have been the main focus in the majority of published studies on litter decomposition above the treeline in alpine areas, while the potential impacts of trophic interactions have received less attention. The review also revealed a severe lack of studies from Africa and South America. Surprisingly, all studies from North America were from Colorado, while all studies from Asia were from China. There is thus a need for studies in Africa and South America, and also in other regions of North America and Asia, to provide data across broader scales. Overall, Europe was well represented in terms of number of studies and different alpine regions. However, more recent studies have mainly originated from China. A noteworthy finding was that factors (altitude, experimental warming, litter quality, soil organisms, soil physico-chemical properties) covered by more than one study all showed inconsistent results regarding their effect on decomposition. Thus, there is a need for further studies on how different factors interact/influence the litter decomposition process at high-alpine sites. Future research should focus on interactions between different factors and conduct experiments to examine specific relationships, such as potential interactions between temperature and soil moisture and the effect on litter decomposition on soils with different moisture levels. There is also a need to evaluate the impact of trophic interactions between organisms and decomposition processes.




Fig. 2. Geographical distribution of studies on plant litter decomposition in alpine regions above treeline.

Author contributions

JMA conceived the study, MR and JMA conducted the literature review, MR and JMA were the main authors of the paper, AKJ and YB commented on the manuscript.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

This study was supported by a grant from Qatar Petroleum (QUEX-CAS-QP-RD-18/19).

References

- Aerts, R., 2006. The freezer defrosting: global warming and litter decomposition rates in cold biomes. *J. Ecol.* 94, 713–724. <https://doi.org/10.1111/j.1365-2745.2006.01142.x>.
- Alatalo, J.M., Jägerbrand, A.K., Chen, S., Molau, U., 2017a. Responses of lichen communities to 18 years of natural and experimental warming. *Ann. Bot.* 120, 159–170.
- Alatalo, J.M., Jägerbrand, A.K., Čuchta, P., 2015. Collembola at three alpine subarctic sites resistant to twenty years of experimental warming. *Sci. Rep.* 5, 18161. <https://doi.org/10.1038/srep18161>.
- Alatalo, J.M., Jägerbrand, A.K., Juhanson, J., Michelsen, A., Lūptāčik, P., 2017b. Impacts of twenty years of experimental warming on soil carbon, nitrogen, moisture and soil mites across alpine/subarctic tundra communities. *Sci. Rep.* 7, 44489. <https://doi.org/10.1038/srep44489>.
- Andreyashkina, N.I., 2008. Plant matter decomposition in the upper timberline ecotone of the Polar Urals. *Russ. J. Ecol.* 39, 490–494. <https://doi.org/10.1134/S1067413608070059>.
- Andreyashkina, N.I., Peshkova, N.V., 2003. On characteristics of production and destruction processes in lowland and mountain tundras of the extreme North. *Russ. J. Ecol.* 34, 98–103. <https://doi.org/10.1023/A:1023094913512>.
- Arnold, J.A., Hirschel, G., 1997. Does fertilizer application alter the effects of elevated CO₂ on Carex leaf litter quality and in situ decomposition in an alpine grassland? In: *Acta Oecologica, from Alpine Grassland to Tropical Forest: Biological Consequences of Elevated Atmospheric CO₂ (A Synthesis of Swiss Research)*, 18, pp. 201–206. [https://doi.org/10.1016/S1146-609X\(97\)80006-9](https://doi.org/10.1016/S1146-609X(97)80006-9).
- Baptist, F., Yoccoz, N.G., Choler, P., 2010. Direct and indirect control by snow cover over decomposition in alpine tundra along a snowmelt gradient. *Plant Soil* 328, 397–410. <https://doi.org/10.1007/s11104-009-0119-6>.
- Bryant, D.M., Holland, E.A., Seastedt, T.R., Walker, M.D., 1998. Analysis of litter decomposition in an alpine tundra. *Can. J. Bot.* 76, 1295–1304. <https://doi.org/10.1139/b98-117>.
- Carbognani, M., Petraglia, A., Tomaselli, M., 2014. Warming effects and plant trait control on the early-decomposition in alpine snowbeds. *Plant Soil* 376, 277–290. <https://doi.org/10.1007/s11104-013-1982-8>.
- Chapin III, F.S., Matson, P.A., Vitousek, P., 2011. *Principles of Terrestrial Ecosystem Ecology*. Springer Science & Business Media.
- Chen, Y., Liu, Y., Zhang, J., Yang, W., He, R., Deng, C., 2018. Microclimate exerts greater control over litter decomposition and enzyme activity than litter quality in an alpine forest-tundra ecotone. *Sci. Rep.* 8, 14998. <https://doi.org/10.1038/s41598-018-33186-4>.

- Cornelissen, J.H.C., Bodegom, P.M.V., Aerts, R., Callaghan, T.V., Logtestijn, R.S.P.V., Alatalo, J., Chapin, F.S., Gerdol, R., Gudmundsson, J., Gwynn-Jones, D., Hartley, A. E., Hik, D.S., Hofgaard, A., Jónsdóttir, I.S., Karlsson, S., Klein, J.A., Laundre, J., Magnusson, B., Michelsen, A., Molau, U., Onipchenko, V.G., Quedest, H.M., Sandvik, S.M., Schmidt, I.K., Shaver, G.R., Solheim, B., Soudzilovskaia, N.A., Stenström, A., Tolvanen, A., Totland, Ø., Wada, N., Welker, J.M., Zhao, X., 2007. Global negative vegetation feedback to climate warming responses of leaf litter decomposition rates in cold biomes. *Ecol. Lett.* 10, 619–627. <https://doi.org/10.1111/j.1461-0248.2007.01051.x>.
- Cornwell, W.K., Cornelissen, J.H.C., Amatangelo, K., Dorrepaal, E., Eviner, V.T., Godoy, O., Hobbie, S.E., Hoorens, B., Kurokawa, H., Pérez-Harguindeguy, N., Quedest, H.M., Santiago, L.S., Wardle, D.A., Wright, I.J., Aerts, R., Allison, S.D., Bodegom, P.V., Brovkin, V., Chatain, A., Callaghan, T.V., Díaz, S., Garnier, E., Gurvich, D.E., Kazakou, E., Klein, J.A., Read, J., Reich, P.B., Soudzilovskaia, N.A., Vaieretti, M.V., Westoby, M., 2008. Plant species traits are the predominant control on litter decomposition rates within biomes worldwide. *Ecol. Lett.* 11, 1065–1071. <https://doi.org/10.1111/j.1461-0248.2008.01219.x>.
- Coûteaux, M.M., Sarmiento, L., Bottner, P., Acevedo, D., Thiéry, J.M., 2002. Decomposition of standard plant material along an altitudinal transect (65–3968 m) in the tropical Andes. *Soil Biol. Biochem.* 34, 69–78.
- Djukic, I., Kepfer-Rojas, S., Schmidt, I.K., Larsen, K.S., Beier, C., Berg, B., Verheyen, K., 2018. Early stage litter decomposition across biomes. *Sci. Total Environ.* 628–629, 1369–1394. <https://doi.org/10.1016/j.scitotenv.2018.01.012>.
- Drewnik, M., 2006. The effect of environmental conditions on the decomposition rate of cellulose in mountain soils. *Geoderma* 132, 116–130. <https://doi.org/10.1016/j.geoderma.2005.04.023>.
- Duboc, O., Zehetner, F., Djukic, I., Tatzber, M., Berger, T.W., Gerzabek, M.H., 2012. Decomposition of European beech and Black pine foliar litter along an Alpine elevation gradient: mass loss and molecular characteristics. *Geoderma* 189–190, 522–531. <https://doi.org/10.1016/j.geoderma.2012.06.018>.
- Freschet, G.T., Cornwell, W.K., Wardle, D.A., Elumeeva, T.G., Liu, W., Jackson, B.G., Onipchenko, V.G., Soudzilovskaia, N.A., Tao, J., Cornelissen, J.H.C., 2013. Linking litter decomposition of above- and below-ground organs to plant–soil feedbacks worldwide. *J. Ecol.* 101, 943–952. <https://doi.org/10.1111/1365-2745.12092>.
- Gavazov, K.S., 2010. Dynamics of alpine plant litter decomposition in a changing climate. *Plant Soil* 337, 19–32. <https://doi.org/10.1007/s11040-010-0477-0>.
- Giweta, M., 2020. Role of litter production and its decomposition, and factors affecting the processes in a tropical forest ecosystem: a review. *J. Ecol. Environ.* 44, 11. <https://doi.org/10.1186/s41610-020-0151-2>.
- Håggvar, S., Klanderud, K., 2009. Effect of simulated environmental change on alpine soil arthropods. *Global Change Biol.* 15, 2972–2980. <https://doi.org/10.1111/j.1365-2486.2009.01926.x>.
- Hartig, R., 1878. Die Zersetzungserscheinungen des Holzes der Nadelholzbäume und der Eiche in forstlicher, botanischer und chemischer Richtung. J. Springer.
- Hong, J., Lu, X., Ma, X., Wang, X., 2021. Five-year study on the effects of warming and plant litter quality on litter decomposition rate in a Tibetan alpine grassland. *Sci. Total Environ.* 750, 142306. <https://doi.org/10.1016/j.scitotenv.2020.142306>.
- Huang, C., Wu, F., Yang, W., Tan, B., He, W., Zhang, J., 2016. Effects of snow thickness on the abundance of archaeal and bacterial amoA genes and gene transcripts during dwarf bamboo litter decomposition in an alpine forest on the eastern Tibetan Plateau. *Russ. J. Ecol.* 47, 419–429. <https://doi.org/10.1134/S106741361604010X>.
- Huber, E., Wanek, W., Gottfried, M., Pauli, H., Schweiger, P., Arndt, S.K., Reiter, K., Richter, A., 2007. Shift in soil–plant nitrogen dynamics of an alpine–nival ecotone. *Plant Soil* 301, 65–76. <https://doi.org/10.1007/s10047-007-9422-2>.
- Keuskamp, J.A., Dingemans, B.J.J., Lehtinen, T., Sarneel, J.M., Hefting, M.M., 2013. Tea Bag Index: a novel approach to collect uniform decomposition data across ecosystems. *Methods Ecol. Evol.* 4, 1070–1075. <https://doi.org/10.1111/2041-210X.12097>.
- Kitz, F., Steinwandter, M., Traugott, M., Seeber, J., 2015. Increased decomposer diversity accelerates and potentially stabilises litter decomposition. *Soil Biol. Biochem.* 83, 138–141. <https://doi.org/10.1016/j.soilbio.2015.01.026>.
- Körner, C., 1999. *Alpine Plant Life: Functional Plant Ecology of High Mountain Ecosystems*. Springer, Berlin.
- Krishna, M.P., Mohan, M., 2017. Litter decomposition in forest ecosystems: a review. *Energy Ecol. Environ.* 2, 236–249. <https://doi.org/10.1007/s40974-017-0064-9>.
- Liang, D., Lamb, E.G., Zhang, S., 2019. Yak dung deposition affects litter mixing effects on mass loss in Tibetan alpine grassland. *Rangel. Ecol. Manag.* 72, 405–410. <https://doi.org/10.1016/j.rama.2018.11.004>.
- Lindquist, B., 1941. Undersökningar över några skandinaviska dagmaskarters betydelse för lövförmåns omvandling och för muldjordens struktur i svensk skogsmark. *Sven. Skogsv.- Foren Tidskr* 39, 179–242.
- Lipson, D.A., Schmidt, S.K., Monson, R.K., 2000. Carbon availability and temperature control the post-snowmelt decline in alpine soil microbial biomass. *Soil Biol. Biochem.* 32, 441–448. [https://doi.org/10.1016/S0038-0717\(99\)00068-1](https://doi.org/10.1016/S0038-0717(99)00068-1).
- Liu, H., Lin, L., Wang, H., Zhang, Z., Shangguan, Z., Feng, X., He, J.-S., 2021. Simulating warmer and drier climate increases root production but decreases root decomposition in an alpine grassland on the Tibetan plateau. *Plant Soil* 458, 59–73. <https://doi.org/10.1007/s11040-020-04551-y>.
- Luo, C., Xu, G., Chao, Z., Wang, S., Lin, X., Hu, Y., Zhang, Z., Duan, J., Chang, X., Su, A., Li, Y., Zhao, X., Du, M., Tang, Y., Kimball, B., 2010. Effect of warming and grazing on litter mass loss and temperature sensitivity of litter and dung mass loss on the Tibetan plateau. *Global Change Biol.* 16, 1606–1617. <https://doi.org/10.1111/j.1365-2486.2009.02026.x>.
- Lv, W., Zhang, L., Niu, H., Li, B., Wang, Q., Zhou, Y., Wang, Y., Jiang, L., Liu, P., Hong, H., Jia, S., Luo, C., Tsechoe, D., Zhou, H., Wang, S., 2020. Non-linear temperature sensitivity of litter component decomposition under warming gradient with precipitation addition on the Tibetan plateau. *Plant Soil* 448, 335–351. <https://doi.org/10.1007/s11040-020-04431-5>.
- Ma, C., Yin, X., Wang, H., 2019. Soil fauna effect on *Dryas octopetala* litter decomposition in an Alpine tundra of the Changbai Mountains, China. *Alpine Bot.* 129, 53–62. <https://doi.org/10.1007/s00035-018-0215-4>.
- Mori, A.S., Cornelissen, J.H.C., Fujii, S., Okada, K., Isbell, F., 2020. A meta-analysis on decomposition quantifies afterlife effects of plant diversity as a global change driver. *Nat. Commun.* 11, 1–9. <https://doi.org/10.1038/s41467-020-18296-w>.
- Murphy, K.L., Klopatek, J.M., Klopatek, C.C., 1998. The effects of litter quality and climate on decomposition along an elevational gradient. *Ecol. Appl.* 8, 1061–1071. <https://doi.org/10.2307/2640961>.
- Novara, A., Rühl, J., La Mantia, T., Gristina, L., La Bella, S., Tuttolomondo, T., 2015. Litter contribution to soil organic carbon in the processes of agriculture abandonment. *Solid Earth* 6. <https://doi.org/10.5194/se-6-425-2015>.
- O’Lear, H.A., Seastedt, T.R., 1994. Landscape patterns of litter decomposition in alpine tundra. *Oecologia* 99, 95–101. <https://doi.org/10.1007/BF00317088>.
- Phillips, H.R.P., Guerra, C.A., Bartz, M.L.C., Briones, M.J.I., Brown, G., Crowther, T.W., Ferlian, O., Gongalsky, K.B., Hoogen, J., van den, Krebs, J., Orgiazzi, A., Routh, D., Schwarz, B., Bach, E.M., Bennett, J.M., Brose, U., Decaens, T., König-Ries, B., Loreau, M., Mathieu, J., Mulder, C., Putten, W.H. van der, Ramirez, K.S., Rillig, M.C., Russell, D., Rutgers, M., Thakur, M.P., Vries, F.T. de, Wall, D.H., Wardle, D.A., Arai, M., Ayuke, F.O., Baker, G.H., Beauséjour, R., Bedard, J.C., Birkhofer, K., Blanchart, E., Blossey, B., Bolger, T., Bradley, R.L., Callahan, M.A., Capowiez, Y., Caulfield, M.E., Choi, A., Crotty, F.V., Crumsey, J.M., Dávalos, A., Cosin, D.J.D., Dominguez, A., Duhour, A.E., Eekeren, N. van, Emmerling, C., Falco, L.B., Fernández, R., Fonte, S.J., Fragoso, C., Franco, A.L.C., Fugère, M., Fusilero, A.T., Gholami, S., Gundale, M.J., López, M.G., Hackenberger, D.K., Hernández, L.M., Hishi, T., Holdsworth, A.R., Holmstrup, M., Hofensperger, K.N., Lwanga, E.H., Huhta, V., Hurisso, T.T., Iannone, B.V., Iordache, M., Joschko, M., Kaneko, N., Kanianska, R., Keith, A.M., Kelly, C.A., Kernecker, M.L., Klaminder, J., Koné, A.W., Kooch, Y., Kukkonen, S.T., Lalthanzara, H., Lammel, D.R., Lebedev, I.M., Li, Y., Lidon, J.B.J., Lincoln, N.K., Loss, S.R., Marichal, R., Matula, R., Moos, J.H., Moreno, G., Morón-Ríos, A., Muys, B., Neirynck, J., Norgrove, L., Novo, M., Nuutinen, V., Nuzzo, V., Rahman, M., Pansu, J., Paudel, S., Pérès, G., Pérez-Camacho, L., Piñeiro, R., Ponge, J.-F., Rashid, M.I., Rebollo, S., Rodeiro-Iglesias, J., Rodríguez, M.A., Roth, A.M., Rousseau, G.X., Rozen, A., Sayad, E., Schaik, L. van, Scharenbroch, B.C., Schirrmann, M., Schmidt, O., Schröder, J., Seeber, J., Shashkov, M.P., Singh, J., Smith, S.M., Steinwandter, M., Talavera, J.A., Trigo, D., Tsukamoto, J., Valença, A.W., de Vanek, S.J., Virto, I., Wackett, A.A., Warren, M.W., Wehr, N.H., Whalen, J.K., Wironen, M.B., Wolters, V., Zenkova, I.V., Zhang, W., Cameron, E.K., Eisenhauer, N., 2019. Global distribution of earthworm diversity. *Science* 366, 480–485. <https://doi.org/10.1126/science.aax4851>.
- Prescott, C.E., 2010. Litter decomposition: what controls it and how can we alter it to sequester more carbon in forest soils? *Biogeochemistry* 101, 133–149. <https://doi.org/10.1007/s10533-010-9439-0>.
- Roos, R.E., Birkemoe, T., Asplund, J., L’uptáček, P., Raschmanová, N., Alatalo, J.M., Olsen, S.L., Klanderud, K., 2020. Legacy effects of experimental environmental change on soil micro-arthropod communities. *Ecosphere* 11, e03030.
- Sarneel, J.M., Sundqvist, M.K., Molau, U., Björkman, M.P., Alatalo, J.M., 2020. Decomposition rate and stabilization across six tundra vegetation types exposed to >20 years of warming. *Sci. Total Environ.* 724, 138304. <https://doi.org/10.1016/j.scitotenv.2020.138304>.
- Schaefer, M., Migge-Kleian, S., Scheu, S., 2009. The role of soil fauna for decomposition of plant residues. *Functioning and Management of European Beech Ecosystems*, pp. 207–230. <https://doi.org/10.1007/b82392.13>.
- Schinner, F., 1983. Litter decomposition, CO₂-release and enzyme activities in a snowed and on a windswept ridge in an alpine environment. *Oecologia* 59, 288–291. <https://doi.org/10.1007/BF00378850>.
- Schinner, F., 1982. Soil microbial activities and litter decomposition related to altitude. *Plant Soil* 65, 87–94. <https://doi.org/10.1007/BF02376806>.
- Schwager, P., Berg, C., 2019. Global warming threatens conservation status of alpine EU habitat types in the European Eastern Alps. *Reg. Environ. Change* 19, 2411–2421. <https://doi.org/10.1007/s10113-019-01554-z>.
- Seastedt, T.R., Walker, M.D., Bryant, D.M., 2001. *Controls on decomposition processes in alpine tundra. Structure and Function of an Alpine Ecosystem*. Oxford University Press, Oxford, pp. 222–236.
- Seeber, J., Scheu, S., Meyer, E., 2006. Effects of macro-decomposers on litter decomposition and soil properties in alpine pastureland: a mesocosm experiment. *Appl. Soil Ecol.* 34, 168–175. <https://doi.org/10.1016/j.apsoil.2006.02.004>.
- Shaw, M.R., Harte, J., 2001. Control of litter decomposition in a subalpine meadow–sagebrush steppe ecotone under climate change. *Ecol. Appl.* 11, 1206–1223. [https://doi.org/10.1890/1051-0761.011\[1206:COLDIA\]2.0.CO;2](https://doi.org/10.1890/1051-0761.011[1206:COLDIA]2.0.CO;2).
- Sjögersten, S., Wookey, P.A., 2004. Decomposition of mountain birch leaf litter at the forest–tundra ecotone in the Fennoscandian mountains in relation to climate and soil conditions. *Plant Soil* 262, 215–227. <https://doi.org/10.1023/B:PLSO.0000037044.63113.fe>.
- Song, M.-H., Chen, J., Xu, X.-L., Yu, F.-H., Jiang, J., Zheng, L.-L., Cornelissen, J.H.C., 2019. Decreased community litter decomposition associated with nitrogen-induced convergence in leaf traits in an alpine meadow. *Soil Tillage Res.* 194, 104332. <https://doi.org/10.1016/j.still.2019.104332>.
- Song, P., Zhang, N.L., Ma, K.P., Guo, J.X., 2014. Impacts of global warming on litter decomposition. *Shengtai Xuebao Acta Ecol. Sin.* 34, 1327–1339. <https://doi.org/10.5846/stxb201210251479>.

- Soudzilovskaia, N.A., Onipchenko, V.G., Cornelissen, J.H., Aerts, R., 2007. Effects of fertilisation and irrigation on 'foliar afterlife' in alpine tundra. *J. Veg. Sci.* 18, 755–766. <https://doi.org/10.1111/j.1654-1103.2007.tb02591.x>.
- Steltzer, H., Bowman, W.D., 2005. Litter N retention over winter for a low and a high phenolic species in the alpine tundra. *Plant Soil* 275, 361–370. <https://doi.org/10.1007/s11004-005-3100-z>.
- Štursová, M., Šnajdr, J., Koukol, O., Tláškal, V., Cajthaml, T., Baldrian, P., 2020. Long-term decomposition of litter in the montane forest and the definition of fungal traits in the successional space. *Fungal Ecol* 100913.
- Sun, Y., He, X.Z., Hou, F., Wang, Z., Chang, S., 2018. Grazing increases litter decomposition rate but decreases nitrogen release rate in an alpine meadow. *Biogeosciences* 15, 4233–4243. <https://doi.org/10.5194/bg-15-4233-2018>.
- Swift, M.J., Heal, O.W., Anderson, Jonathan Michael, Anderson, J.M., 1979. *Decomposition in Terrestrial Ecosystems*. University of California Press.
- Vaieretti, M.V., Iamamoto, S., Pérez Harguindeguy, N., Cingolani, A.M., 2018. Livestock grazing affects microclimate conditions for decomposition process through changes in vegetation structure in mountain grasslands. *Acta Oecol.* 91, 101–107. <https://doi.org/10.1016/j.actao.2018.07.002>.
- Waksman, S.A., Tenney, F.G., Stevens, K.R., 1928. The role of microorganisms in the transformation of organic matter in forest soils. *Ecology* 9, 126–144. <https://doi.org/10.2307/1929350>.
- Wang, L., Zhang, J., He, R., Chen, Y., Yang, L., Zheng, H., Li, H., Xiao, J., Liu, Y., 2018. Impacts of soil fauna on lignin and cellulose degradation in litter decomposition across an alpine forest-tundra ecotone. *Eur. J. Soil Biol.* 87, 53–60. <https://doi.org/10.1016/j.ejsobi.2018.05.004>.
- Wang, Lifeng, Chen, Y., Zhou, Y., Zheng, H., Xu, Z., Tan, B., You, C., Zhang, L., Li, H., Guo, L., Wang, Lixia, Huang, Y., Zhang, J., Liu, Y., 2021. Litter chemical traits strongly drove the carbon fractions loss during decomposition across an alpine treeline ecotone. *Sci. Total Environ.* 753, 142287. <https://doi.org/10.1016/j.scitotenv.2020.142287>.
- Wang, P., Guo, J., Xu, X., Yan, X., Zhang, K., Qiu, Y., Zhao, Q., Huang, K., Luo, X., Yang, F., Guo, H., Hu, S., 2020. Soil acidification alters root morphology, increases root biomass but reduces root decomposition in an alpine grassland. *Environ. Pollut.* 265, 115016. <https://doi.org/10.1016/j.envpol.2020.115016>.
- Wang, S., Ruan, H., Wang, B., 2009. Effects of soil microarthropods on plant litter decomposition across an elevation gradient in the Wuyi Mountains. *Soil Biol. Biochem.*, Science Goes Underground in China 41, 891–897. <https://doi.org/10.1016/j.soilbio.2008.12.016>.
- Withington, C.L., Sanford, R.L., 2007. Decomposition rates of buried substrates increase with altitude in the forest-alpine tundra ecotone. *Soil Biol. Biochem.* 39, 68–75.
- Wood, T.G., 1970. Decomposition of plant litter in montane and alpine soils on Mt Kosciuszko, Australia. *Nature* 226, 561–562. <https://doi.org/10.1038/226561a0>.
- Zeidler, M., Duchoslav, M., Banaš, M., 2014. Effect of altered snow conditions on decomposition in three subalpine plant communities. *Open Life Sci.* 9, 811–822. <https://doi.org/10.2478/s11535-014-0312-3>.
- Zhang, D., Hui, D., Luo, Y., Zhou, G., 2008. Rates of litter decomposition in terrestrial ecosystems: global patterns and controlling factors. *J. Plant Ecol.* 1, 85–93. <https://doi.org/10.1093/jpe/rtn002>.
- Zheng, H., Chen, Y., Liu, Y., Hedenec, P., Peng, Y., Xu, Z., Tan, B., Zhang, L., Guo, L., Wang, L., 2020. Effects of litter quality diminish and effects of vegetation type develop during litter decomposition of two shrub species in an alpine treeline ecotone. *Ecosystems* 1–14. <https://doi.org/10.1007/s10021-020-00512-9>.
- Zheng, H., Chen, Y., Liu, Y., Zhang, J., Yang, W., Yang, L., Li, H., Wang, L., Wu, F., Guo, L., 2018. Litter quality drives the differentiation of microbial communities in the litter horizon across an alpine treeline ecotone in the eastern Tibetan Plateau. *Sci. Rep.* 8, 10029. <https://doi.org/10.1038/s41598-018-28150-1>.
- Zhou, G., Guan, L., Wei, X., Tang, X., Liu, S., Liu, J., Zhang, D., Yan, J., 2008. Factors influencing leaf litter decomposition: an intersite decomposition experiment across China. *Plant Soil* 311, 61–72. <https://doi.org/10.1007/s11104-008-9658-5>.
- Zhou, S., Butenschoten, O., Barantal, S., Handa, I.T., Makkonen, M., Vos, V., Aerts, R., Berg, M.P., McKie, B., Ruijven, J.V., Hättenschwiler, S., Scheu, S., 2020. Decomposition of leaf litter mixtures across biomes: the role of litter identity, diversity and soil fauna. *J. Ecol.* 108, 2283–2297. <https://doi.org/10.1111/1365-2745.13452>.