

Mitochondrial DNA Part B Resources

ISSN: (Print) (Online) Journal homepage: <https://www.tandfonline.com/loi/tmdn20>

The complete chloroplast genomes of two *Pedicularis* species (Orobanchaceae) from Southwest China

Wei-Jia Wang, Rong Liu, You Wu, Hong Wang & Wen-Bin Yu

To cite this article: Wei-Jia Wang, Rong Liu, You Wu, Hong Wang & Wen-Bin Yu (2022) The complete chloroplast genomes of two *Pedicularis* species (Orobanchaceae) from Southwest China, Mitochondrial DNA Part B, 7:6, 971-973, DOI: [10.1080/23802359.2022.2080018](https://doi.org/10.1080/23802359.2022.2080018)

To link to this article: <https://doi.org/10.1080/23802359.2022.2080018>



© 2022 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.



Published online: 10 Jun 2022.



Submit your article to this journal [↗](#)



Article views: 243



View related articles [↗](#)




View Crossmark data [↗](#)

MITOGENOME ANNOUNCEMENT



The complete chloroplast genomes of two *Pedicularis* species (Orobanchaceae) from Southwest China

Wei-Jia Wang^{a,b}, Rong Liu^{b,c}, You Wu^{a,b}, Hong Wang^a and Wen-Bin Yu^{c,d,e} 

^aKey Laboratory for Plant Diversity and Biogeography of East Asia, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, China; ^bUniversity of Chinese Academy of Sciences, Beijing, China; ^cCenter for Integrative Conservation, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Menglun, China; ^dCenter of Conservation Biology, Core Botanical Gardens, Chinese Academy of Sciences, Menglun, China; ^eSoutheast Asia Biodiversity Research Institute, Chinese Academy of Science, Nay Pyi Taw, Myanmar

ABSTRACT

We report the complete chloroplast genome (plastome) sequences of *Pedicularis cephalantha* (147,087 bp) and *P. nigra* (145,726 bp), endemic to southwestern China. Both plastomes have typical quadripartite structures with one large-single copy region, one small-single copy region, and two inverted repeat regions. Both plastome sequences contained 37 tRNA genes and eight rRNA genes, but they differed in the numbers of protein-coding genes: *P. cephalantha* had 76 functional genes and 12 pseudogenes while *P. nigra* had 74 functional genes and 13 pseudogenes. Phylogenetic analysis shows that *P. cephalantha* and *P. nigra* are closely related, then sister to *P. oederi* in the family Orobanchaceae.

ARTICLE HISTORY

Received 25 January 2022
Accepted 15 May 2022

KEYWORDS

Chloroplast genome;
Orobanchaceae; *Pedicularis*;
plastome; *Pedicularis*
cephalantha;
Pedicularis nigra

Pedicularis L. (Orobanchaceae) is the largest genus of hemiparasitic plants (c. 500 species; Yu et al. 2015) and shows remarkable diversification in corolla architecture and floral presentation. Approximately, 360 species are native to China with two-thirds endemic to the Himalaya-Hengduan Mountains region (Yang et al. 1998; Yu et al. 2015). Most previous molecular phylogenies of *Pedicularis* were based primarily on short DNA markers (Ree 2005; Yang and Wang 2007; Robart et al. 2015; Yu et al. 2015). However, chloroplast genome sequences provide additional information for clarifying molecular genetics, species delimitation, and phylogenetic reconstruction, so *Pedicularis* plastomes have been sequenced and used to build phylogenies of this genus (Li et al. 2021). In this study, we completed whole genome sequencing and *de novo* assembly of the chloroplast genomes of *Pedicularis cephalantha* Franch. ex Maxim. 1888 and *Pedicularis nigra* (Bonati) Vaniot ex Bonati 1921. We then reconstructed a plastome phylogeny within the genus by combining available plastome data of *Pedicularis* species in GenBank to determine the phylogenetic positions of both species. Voucher specimens of *P. cephalantha* (Liu R. et al., LR202070703) and *P. nigra* (Yu W.-B. et al., LIDZ1331) were collected from the Yulong Mountain, Lijiang, Yunnan, China (108°13'E, 27°00'N) and from Gaolushan, Yuxi, Yunnan, China (102°36'E, 24°50'N), respectively. Voucher specimens of the two species were deposited at the Herbarium of the Kunming Institute of Botany, Chinese Academy of Sciences

(KUN, <http://www.kun.ac.cn>, Prof. Zhu-Liang Yang, fungi@mail.kib.ac.cn, Yunnan, China).

Total genomic DNA was extracted from leaves of both species dried with silica gel using a modified CTAB method (Doyle and Doyle 1987). For library construction using the Kit protocol (NEBNext Ultra II DNA Library Prep Kit for Illumina), purified DNA was fragmented to 300–500 bp (Zeng et al. 2018). Illumina Hi-Seq 2500 was used to construct the library with 150 by pair-end reads. Around 21,775,108 reads (12,293,748 for *P. cephalantha* and 9,481,360 for *P. nigra*) were obtained. These raw data were used to assemble the complete plastome sequences *de novo* using the GetOrganelle toolkit (Jin et al. 2020). Read mappings showed coverage depths of ~455× in *P. cephalantha* and 805× in *P. nigra*. The chloroplast genomes were annotated using Geseq with the MPI-MP Reference Set (Tillich et al. 2017), then manually adjusted the start and stop codons of protein-coding genes using Geneious (Kearse et al. 2012). The newly assembled plastomes from *P. cephalantha* (NCBI accession: OL606628) and *P. nigra* (NCBI accession: OL544940) showed typical quadripartite structures with one large-single copy region, and one small-single copy region separated by two inverted repeat regions. Plastome sizes of *P. cephalantha* and *P. nigra* were 147,087 bp (GC content: 38.4%; LSC: 82,451 bp; SSC: 14,394 bp; IR: 25,121 bp) and 14,726 bp (GC content: 38.4%; LSC: 82,006 bp; SSC: 15,116 bp; IR: 24,302 bp), respectively. The total gene contents were 133 in *P. cephalantha*

CONTACT Hong Wang  wanghong@mail.kib.ac.cn  Key Laboratory for Plant Diversity and Biogeography of East Asia, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, China; Wen-Bin Yu  yuwenbin@xtbg.ac.cn  Center for Integrative Conservation, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Mengla, China

© 2022 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1. Phylogeny of 24 *Pedicularis* species including three varieties of *P. rex* based on chloroplast genomes using maximum-likelihood methods with bootstrap values on the branch. The bottom scale bar represents the number of substitutions per site.

and 132 in *P. nigra*. Both plastomes had 37 tRNA genes and eight rRNA genes, but the numbers of protein-coding genes differed: *P. cephalantha* had 76 functional genes and 12 pseudogenes while *P. nigra* had 74 functional genes and 13 pseudogenes.

To clarify the phylogenetic positions of these two species, we downloaded an additional 29 published plastomes of 22 *Pedicularis* species from GenBank. These complete plastome sequences with one IR region were aligned using MAFFT (Katoh and Standley 2013). The plastome of *Phtheiospermum japonicum* (Thunb.) Kanitz (Orobanchaceae) was selected as the outgroup. A maximum-likelihood tree was reconstructed using raxmlGUI (Edler et al. 2021) with GTRGAMMAI model by running 1000 bootstrap replicates. Our phylogenetic analyses showed that *P. cephalantha* and *P. nigra* formed a clade with 100% bootstrap values (Figure 1), representing Clade 3 as in the previous study (Yu et al. 2015). Our new clade is sister to *P. oederi* Vahl in the Clade 7. The complete chloroplast genomes and phylogenetic results in this study provide new insights into the phylogenetic backbone and evolutionary biology of *Pedicularis*.

Acknowledgements

We are grateful for the support of the Molecular Biology Experiment Center, Germplasm Bank of Wild Species in Southwest China, Kunming

Institute of Botany, Chinese Academy of Sciences, and the HPC Platform of the Public Technology Service Center, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences. We give particular thanks to Prof. Peter Bernhardt from the St. Louis University and Prof. Richard T. Corlett from the Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences for checking and correcting grammar and syntax.

Ethics statement

No specific permissions were required for these location and activities during fieldwork, and neither of the species studied are listed as endangered or protected in China's Catalogue of National Protected Key Wild Plants.

Author contributions

Hong Wang, Wen-Bin Yu, and Wei-Jia Wang designed and modified the experimental design. Wei-Jia Wang, Rong Liu, You Wu, and Wen-Bin Yu collected and analyzed the data. Wei-Jia Wang, Hong Wang, and Wen-Bin Yu interpreted the results and wrote the manuscript.

Disclosure statement

No potential conflict of interest is reported by the authors.

Funding

This study was supported by grants from the National Natural Science Foundation of China [Nos. 31870196, 32071670], the Large-Scale Scientific Facilities of the Chinese Academy of Sciences [No. 2017-LSFGBOWS-02], and the Open Research Project of “Cross-Cooperative Team” of the Germplasm Bank of Wild Species, Kunming Institute of Botany, Chinese Academy of Sciences, the CAS 135 Program [No. 2017XTBG-T03].

ORCID

Wen-Bin Yu  <http://orcid.org/0000-0002-7643-2112>

Data availability statement

The data were collected without violation of the protection of human subjects, or other valid ethical, privacy, or security concerns. Two complete plastome sequences were deposited in GenBank with accession numbers OL606628 and OL544940 and are also available at Figshare (<https://doi.org/10.6084/m9.figshare.17097098.v2>). The associated BioProject, SRA, and Bio-Sample numbers are PRJNA780958, SRR16960915–SRR16962222, and SAMN23224351–SAMN23224352, respectively.

References

- Doyle JJ, Doyle JL. 1987. A rapid DNA isolation procedure for small quantities of fresh leaf tissue. *Phytochemistry*. 19(1):971–15.
- Edler D, Klein J, Antonelli A, Silvestro D. 2021. RaxmlGUI 2.0: a graphical interface and toolkit for phylogenetic analyses using RAXML. *Methods Ecol Evol*. 12(2):373–377.
- Jin J-J, Yu W-B, Yang J-B, Song Y, dePamphilis CW, Yi T-S, Li D-Z. 2020. GetOrganelle: a fast and versatile toolkit for accurate de novo assembly of organelle genomes. *Genome Biol*. 21(1):241.
- Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol*. 30(4):772–780.
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, et al. 2012. Geneious basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics*. 28(12):1647–1649.
- Li X, Yang J-B, Wang H, Song Y, Corlett RT, Yao X, Li D-Z, Yu W-B. 2021. Plastid NDH pseudogenization and gene loss in a recently derived lineage from the largest hemiparasitic plant genus *Pedicularis* (Orobanchaceae). *Plant Cell Physiol*. 62(6):971–984.
- Ree RH. 2005. Phylogeny and the evolution of floral diversity in *Pedicularis* (Orobanchaceae). *Int J Plant Sci*. 166(4):595–613.
- Robart BW, Gladys C, Frank T, Kilpatrick S. 2015. Phylogeny and biogeography of North American and Asian *Pedicularis* (Orobanchaceae). *Syst Bot*. 40(1):229–258.
- Tillich M, Lehwark P, Pellizzer T, Ulbricht-Jones ES, Fischer A, Bock R, Greiner S. 2017. GeSeq – versatile and accurate annotation of organelle genomes. *Nucleic Acids Res*. 45(W1):W6–W11.
- Yang F-S, Wang X-Q. 2007. Extensive length variation in the cpDNA trnT-trnF region of hemiparasitic *Pedicularis* and its phylogenetic implications. *Plant Syst Evol*. 264(3–4):251–264.
- Yang H-B, Holmgren NH, Mill RR. 1998. *Pedicularis* Linn. In: Wu Z-Y, Raven PH, editors. *Flora of China*. St. Louis, Beijing: Missouri Botanical Garden Press & Science Press; p. 97–209.
- Yu W-B, Liu M-L, Wang H, Mill RR, Ree RH, Yang J-B, Li D-Z. 2015. Towards a comprehensive phylogeny of the large temperate genus *Pedicularis* (Orobanchaceae), with an emphasis on species from the Himalaya-Hengduan Mountains. *BMC Plant Biol*. 15(1):176.
- Zeng C-X, Hollingsworth PM, Yang J, He Z-S, Zhang Z-R, Li D-Z, Yang J-B. 2018. Genome skimming herbarium specimens for DNA barcoding and phylogenomics. *Plant Methods*. 14(1):43.