

中国兰科二新记录种*

李剑武¹, 叶德平², 刘 强¹, 殷建涛¹

(1 中国科学院西双版纳热带植物园, 云南 勐仑 666303; 2 普洱市林业局, 云南 普洱 665000)

摘要: 报道了中国兰科2新记录种, 大花大苞兰 (*Sunipia grandiflora* (Rolfe) P. F. Hunt) 及小花石豆兰 (*Bulbophyllum parviflorum* C. S. P. Parish & Rehb. f.)。

关键词: 大花大苞兰; 小花石豆兰; 中国; 新记录; 兰科

中图分类号: Q 949

文献标识码: A

文章编号: 2095-0845(2013)02-128-03

Two New Records of Orchid from China*

LI Jian-Wu¹, YE De-Ping², LIU Qiang¹, YIN Jian-Tao¹

(1 Xishuangbanna Tropical Botanical Garden, the Chinese Academy of Sciences, Menglun 666303, China;

2 Department of Forest Ministry, Pu'er 665000, China)

Abstract: Two orchids, *Sunipia grandiflora* (Rolfe) P. F. Hunt and *Bulbophyllum parviflorum* C. S. P. Parish & Rehb. f., are reported as new records from China.

Key words: *Sunipia grandiflora*; *Bulbophyllum parviflorum*; China; New record; Orchidaceae

During our botanical trip to Xishuangbanna in southern Yunnan Province, China, between 2011 and 2012, several orchids were found, and recently researching showed two species are new records in China.

1. *Sunipia grandiflora* (Rolfe) P. F. Hunt, Kew Bull. **26** (1): 184. 1971. —*Ione grandiflora* Rolfe in Bull. Misc. Inform. Kew **1908**: 413. 1908.

大花大苞兰 (Fig. 1: A-F)

Rhizomes creeping, ca. 2 mm in diam. Pseudobulbs 1–3 cm apart along the rhizome, ovoid, 8–13 mm long, 8–11 mm in diam., each with 1-leaf on top. Leaf fleshy and leathery; petiole ca. 4 mm long; leaf blade erect, oblong, 4–5×0.8–1 cm, base cuneate, apex obtuse and retuse. Scape ca. 3 cm long; raceme few flowered; floral bracts ovate-lanceolate, ca. 6 mm. Pedicel and ovary ca. 1.2

cm long, flowers purple, sepals full open; dorsal sepal oblong, ca. 13×6 mm, 7-veined, apex acute, reflexed; lateral sepals oblong, ca. 13×8 mm, 5-veined, apex acute, slightly connate; petals subulate, 11×5 mm, fimbriate in middle part of edges; lip broad cordate-ovate, apex obtuse, centre greatly thickened and fleshy, 11×8 mm, margins erose-lacerate, base retuse and carinate. Column ca. 3.5 mm, stout, with a foot ca. 1 mm, apex with 4 dentations; 2 porrect, ca. 2.2 mm long; 2 incurved, ca. 1 mm long. Anther cap galeated; pollinia 4, hemispheric. Flowering from November to December.

China. Yunnan (云南): Xishuangbanna (西双版纳), Menghai (勐海), alt. 1 600 m, 2011–11–26, on tree, broad-leaved evergreen forests, LI Jian-wu (李剑武) 1250 (HITBC).

Distribution: China (Yunnan), Laos, Thailand.

* Foundation item: Special Grant for Herbarium Running Fees from the Chinese Academy of Sciences (08ZK111B02)

Received date: 2012-08-21, Accepted date: 2012-09-25

作者简介: 李剑武 (1983-) 男, 实验师, 主要从事植物分类工作。E-mail: ljw@xtbg.org.cn



Fig. 1 A–F: *Sunipia grandiflora* (Rolfe) P. F. Hunt. ; G: *Bulbophyllum parviflorum* C. S. P. Parish & Rehb. f.
A. Plant; B. Dorsal sepal; C. petal; D. Column; E. Lateral sepals; F. Lip; G. Inflorescence

Sunipia grandiflora is similar to *S. bicolor* Lindl. and *S. kachinensis* Seidenfaden, but differs from them by having purple and larger flowers (3-times larger to *S. bicolor*), entire dorsal sepal (dorsal sepal erose in *S. kachinensis*), lateral sepals slightly connate (lateral sepals connate almost along the whole length in *S. kachinensis*), lip cordate with single ridge at base (lip with 7 rounded and parallel ridges in *S. kachinensis*; lip fiddle-shaped in *S. bicolor*) (Chen et Vermeulen, 2009; Seidenfaden, 1981; Hunt, 1971).

2. *Bulbophyllum parviflorum* C. S. P. Parish & Rchb. f., Trans. Linn. Soc. London **30** (1): 152. 1874.

小花石豆兰 (Fig. 1: G)

Rhizomes creeping, 3–4 mm in diam., branched. Roots arising from nodes of rhizome. Pseudobulbs 2–7 cm apart from each other, ovoid to broadly conic,

ca. 9 mm, 12–15 mm in diam., with a terminal leaf. Leaf leathery, petiole ca. 2 cm; leaf blade narrowly oblong, 11–13 × 1.2–1.4 cm, base contracted into petiole, apex obtuse. Scapes from base of pseudobulbs, erect, 8–12 cm, slender; raceme 4–6 cm, densely with many small flowers; peduncle with 3 tubular sheaths; floral bracts ovate-lanceolate, apex acute, 1.2 mm. Pedicel and ovary ca. 2.5 mm. Flowers pale yellow or whitish, not fully opening; dorsal sepal ovate-lanceolate, 1-veined, apex acute, 3.5 × 1 mm; lateral sepals slightly hairy along the lower margin, ovate-lanceolate, 1-veined, apex acute, 4 × 1.5 mm, base adanate to column foot; petals oblong, 1.5 × 0.5 mm, apex mucronulate, edges ciliate; lip lingulate, 2 × 0.6 mm, fleshy, apex obtuse, edges ciliate, with longitudinal 2 crests in middle. Column ca. 1 mm long; column foot ca. 1 mm long; stelids narrowly lanceolate, ca. 0.3

mm; anther cap densely finely papillate, centrally with one globose projection. Flowering from December to January.

China. Yunnan (云南): Xishuangbanna (西双版纳), Mengla (勐腊), alt. 780 m, 2011–12–16, on tree, tropical rain forests, LI Jian-wu (李剑武) 1518 (HITBC).

Distribution: China (Yunnan) and Myanmar.

Bulbophyllum parviflorum is similar to *B. limbatum* Lindley, however, it is readily to differ it by having less than 3 mm long dorsal sepal (more than 5 mm long *B. limbatum*), white lateral sepals slightly hairy along the lower edge (yellows and minutely hairy along all edges in *B. limbatum*), green lip

with long ciliae (brownish yellow lip without hairs or ciliae in *B. limbatum*) (Seidenfaden, 1979).

Acknowledgements: We are grateful to Dr. X. H. Jin for his critical review of the manuscript and the literatures.

References:

- Chen SC, et Vermeulen JJ, 2009. *Sunipia* [A]. In: Wu ZY, Raven PH, Hong DY (ed.), *Flora of China* Vol. 25 [M]. Beijing: Science Press, 440–444
- Hunt PF, 1971. *Notes on asiatic orchids*: VI [J]. Kew: Kew Bulletin, 26 (1): 171–185
- Seidenfaden G, 1979. Orchid Genera in Thailand VIII. *Bulbophyllum* Thou [M]. Dansk Botanisk Arkiv, 3: 94–103
- Seidenfaden G, 1981. Contributions to the orchid flora of Thailand IX [J]. *Nordic Journal of Botany*, 1 (2): 192–217



《植物分类与资源学报》入选2012年度“中国最具国际影响力学术期刊”

2012年12月26日,中国学术期刊电子杂志社、中国科学文献计量评价研究中心与清华大学图书馆在京发布首份全面评价我国学术期刊国际影响力的名单——2012年度中国最具国际影响力学术期刊和中国国际影响力优秀学术期刊。由中国科学院昆明植物研究所与中国植物学会共同主办的期刊《植物分类与资源学报》入选“中国最具国际影响力学术期刊”。

中国学术期刊(光盘版)电子杂志社、中国科学文献计量评价研究中心和清华大学图书馆,以中国大陆正式出版的5025种学术期刊为研究对象,包括3533种科技期刊和1360种人文社科期刊,将Web of Science收录的国际学术文献作为统计源,以2011年度为统计年,首次研制发布了《中国学术期刊国际引证报告(2012)》,简称CAJ-IJCR(2012),并研究制定了《中国最具国际影响力学术期刊、中国国际影响力优秀学术期刊遴选办法》。

该办法先以CAJ-IJCR(2012)给出的期刊国际影响力指标进行综合排序,再由专家依综合排序按入选条件评审。期刊国际影响力指标为国际总他引频次和国际他引影响因子,两项指标综合排序的权重为2:1。科技期刊、人文社科期刊分别排序。

中国最具国际影响力学术期刊和中国国际影响力优秀期刊,是依据《中国学术期刊国际引证报告(2012版)》,按2011年度中国学术期刊被SCI期刊、SSCI期刊引用的总被引频次和影响因子排序,经综合计算,并经40多位期刊界专家审议,最终遴选出的Top 5%期刊和Top 5%~10%期刊。我国科技期刊备选3533种,由上述方法选出“中国最具国际影响力学术期刊”和“中国国际影响力优秀学术期刊”各175种;人文社科类备选680种,分别选出各34种。

报告表明,SCI收录中国科技期刊127种,但我国非SCI科技期刊异军突起。在175种“2012中国最具国际影响力学术期刊”科技期刊中(其中与生命科学相关的部分期刊见附表),SCI收录87种,非SCI收录88种,未进入Top 5%的SCI收录期刊34种,占SCI收录中国期刊的26.8%。在175种“2012中国国际影响力优秀学术期刊”科技期刊中,SCI收录17种,非SCI收录158种;未进入Top 10%的SCI收录期刊23种,占SCI收录中国期刊的18%。

[下转第148页]