

Hierarchical Scheme for LC-MSⁿ Identification of Chlorogenic AcidsMICHAEL N. CLIFFORD,^{*,†} KELLY L. JOHNSTON,[†] SUSAN KNIGHT,[†] AND
NIKOLAI KUHNERT[§]Centre for Nutrition and Food Safety, and Synthetic and Biological Organic
Chemistry Laboratory, School of Biomedical and Life Sciences, University of Surrey,
Guildford, Surrey GU2 7XH, United Kingdom

The fragmentation behavior of 18 chlorogenic acids that are not substituted at position 1 has been investigated using LC-MS⁴ applied to a methanolic coffee bean extract and commercial cider (hard cider). Using LC-MS³, it is possible to discriminate between each of the three isomers of *p*-coumaroylquinic acid, caffeoylquinic acid, feruloylquinic acid, and dicaffeoylquinic acid, and a hierarchical key has been prepared to facilitate this process when standards are not available. MS⁴ fragmentations further support these assignments, but were not essential in reaching them. The distinctive behavior of 4-acyl and 3-acyl chlorogenic acids compared with the 5-acyl chlorogenic acids is a key factor permitting these assignments. The fragmentation patterns are dependent upon the particular stereochemical relationships between the individual substituents on the quinic acid moiety. Fragmentation is facilitated by 1,2-acyl participation and proceeds through quinic acid conformers in which the relevant substituents transiently adopt a 1,3-*syn*-diaxial relationship. Selected ion monitoring at *m/z* 529 clearly indicated the presence in coffee of six caffeoylferuloylquinic acid isomers, whereas previously only two or three had been demonstrated. The hierarchical key permitted specific structures to be assigned to each of the six isomers. These assignments are internally consistent and consistent with the limited data previously available.

KEYWORDS: Caffeic acid; caffeoylferuloylquinic acids; caffeoylquinic acids; chlorogenic acids; cider; coffee beans; *p*-coumaric acid; *p*-coumaroylquinic acids; dicaffeoylquinic acids; ferulic acid; feruloylquinic acids; LC-MSⁿ; quinic acid

INTRODUCTION

Classically, chlorogenic acids (CGA) are a family of esters formed between certain *trans* cinnamic acids and (–)-quinic acid [1L-1(OH),3,4/5-tetrahydroxycyclohexanecarboxylic acid] (1–3). IUPAC (1) has established a preferred nomenclature for cyclitols, including quinic acid derivatives. This system will be used here, and data published using other systems have been altered to ensure consistency and clarity of discussion.

CGA are characteristic components of coffee beans and commercial coffee products, in which caffeoylquinic (CQA), *p*-coumaroylquinic (*p*CoQA), feruloylquinic (FQA), dicaffeoylquinic (diCQA), and caffeoylferuloylquinic acids (CFQA) have been reported. In coffee, esterification occurs at positions 3, 4, and 5 of the quinic acid moiety, but not at position 1 (2–6). Individual compounds are commonly referred to by a well-established (2, 3, 7, 8) series of structure-defining abbreviations, for example, 3-CQA, 4-CQA, and 5-CQA. The full set of abbreviations and structures are illustrated in **Figure 1**.

The probable existence of CFQA was first noted in 1965 (9). Two CFQA have been characterized by NMR (10, 11). Three CFQA isomers have been demonstrated chromatographically (12), but it has been suggested that green robusta coffee beans would contain six (2, 3). The objectives of this study are to investigate the extent to which individual CGA isomers could be distinguished by LC-MSⁿ and to seek convincing evidence for the presence of six CFQA isomers (XIII–XVIII) in green robusta coffee beans.


MATERIALS AND METHODS

Sample Preparation. Green robusta coffee beans were frozen overnight (–12 °C) and ground in a hammer mill to pass 0.7 mm; 500 mg was extracted (4 × 25 mL, 25 min each) with 70% v/v aqueous methanol using an HT-1043 solid–liquid continuous extraction system (Tecator, Bristol, U.K.) (13). The bulked extracts were treated with Carrez reagents (1 mL of reagent A plus 1 mL of reagent B) (14) to precipitate colloidal material, diluted to 100 mL with 70% v/v aqueous methanol, and filtered through a Whatman no. 1 filter paper. The methanol was removed by evaporation with nitrogen (N-Evap-111, Organomation Associates Inc., Berlin, MA), and the aqueous extracts were stored at –12 °C until required, thawed at room temperature, and used directly for LC-MS.

* Author to whom correspondence should be addressed (fax 44 14 83 57 69 78; e-mail m.clifford@surrey.ac.uk).

[†] Centre for Nutrition and Food Safety.

[§] Synthetic and Biological Organic Chemistry Laboratory.



Name and abbreviation	Number	R ₃	R ₄	R ₅
3- <i>O</i> -caffeoylquinic acid (3-CQA)	I	C	H	H
5- <i>O</i> -caffeoylquinic acid (5-CQA)	II	H	H	C
4- <i>O</i> -caffeoylquinic acid (4-CQA)	III	H	C	H
3- <i>O</i> - <i>p</i> -coumaroylquinic acid (3- <i>p</i> CoQA)	IV	<i>p</i> -Co	H	H
5- <i>O</i> - <i>p</i> -coumaroylquinic acid (5- <i>p</i> CoQA)	V	H	H	<i>p</i> -Co
4- <i>O</i> - <i>p</i> -coumaroylquinic acid (4- <i>p</i> CoQA)	VI	H	<i>p</i> -Co	H
3- <i>O</i> -feruloylquinic acid (3-FQA)	VII	F	H	H
5- <i>O</i> -feruloylquinic acid (5-FQA)	VIII	H	H	F
4- <i>O</i> -feruloylquinic acid (4-FQA)	IX	H	F	H
3,4-di- <i>O</i> -caffeoylquinic acid (3,4-diCQA)	X	C	C	H
3,5-di- <i>O</i> -caffeoylquinic acid (3,5-diCQA)	XI	C	H	C
4,5-di- <i>O</i> -caffeoylquinic acid (4,5-diCQA)	XII	H	C	C
CFQA-1 = 3- <i>O</i> -feruloyl, 4- <i>O</i> -caffeoylquinic acid ^a (3F,4CQA)	XIII	F	C	H ^b
CFQA-2 = 3- <i>O</i> -caffeoyl, 4- <i>O</i> -feruloylquinic acid ^a (3C,4FQA)	XIV	C	F	H
CFQA-3 = 3- <i>O</i> -feruloyl, 5- <i>O</i> -caffeoylquinic acid ^a (3F,5- <i>p</i> CoQA)	XV	F	H	C
CFQA-4 = 3- <i>O</i> -caffeoyl, 5- <i>O</i> -feruloylquinic acid ^a (3C,5FQA)	XVI	C	H	F
CFQA-5 = 4- <i>O</i> -feruloyl, 5- <i>O</i> -caffeoylquinic acid ^a (4F,5CQA)	XVII	H	F	C
CFQA-6 = 4- <i>O</i> -caffeoyl, 5- <i>O</i> -feruloylquinic acid ^a (4C,5FQA)	XVIII	H	C	F

Figure 1. Structure of chlorogenic acids and associated cinnamic acids (IUPAC numbering) (7). ^aStructure assigned during this study.

LC-MSⁿ. The LC equipment (ThermoFinnigan, San Jose, CA) comprised a Surveyor MS pump, an autosampler with a 20 μ L loop, and a PDA detector with a light-pipe flow cell (recording at 320, 280, and 254 nm and scanning from 200 to 600 nm). This was interfaced with an LCQ Deca XP Plus mass spectrometer fitted with an ESI source (ThermoFinnigan) and operating in zoom scan mode for the accurate determination of parent ion *m/z* and in data-dependent, turboscan, MSⁿ mode to obtain fragment ion *m/z*. MS operating conditions (negative ion) had been optimized using 5-CQA (II) with a collision energy of 35%, an ionization voltage of 3.5 kV, a capillary temperature of 350 °C, a sheath gas flow rate of 65 arbitrary units, and an auxiliary gas flow rate of 10 arbitrary units.

CGA separation was achieved on a 150 \times 3 mm column containing Luna 5 μ phenylhexyl packing (Phenomenex, Macclesfield, U.K.). Solvent A was water/acetonitrile/glacial acetic acid (980:20:5 v/v, pH 2.68); solvent B was acetonitrile/glacial acetic acid (1000:5 v/v). Solvents were delivered at a total flow rate of 300 μ L min⁻¹. The gradient profile was 4% B to 33% B linearly in 90 min, a linear increase

to 100% B at 95 min, followed by 5 min isocratic, and a return to 4% B at 105 min, followed by 5 min isocratic to re-equilibrate.

Chlorogenic acid (5-CQA) (I) was obtained from Sigma Chemical Co. (Poole, Dorset, U.K.). A traditional commercial cider (hard cider) was used without further treatment as a convenient source of *p*CoQA (8). NMR-characterized diCQA (X–XII), 5-*p*CoQA (V), and 5-FQA (VIII) were available from previous studies (15). Tetramethylammonium hydroxide (30% methanolic solution) was obtained from Sigma Chemical Co. All other reagents were standard commercial items of an appropriate grade.

RESULTS AND DISCUSSION

General LC-MS Fragmentation. All data for CGA presented in this paper use the recommended IUPAC numbering system (1), and specimen structures are presented in Figure 1. When necessary, previously published data have been amended to ensure consistency and avoid ambiguity. The coffee extract

Table 1. Negative Ion MS⁴ Data for Monoacylchlorogenic Acids

compound	N	MS ¹		MS ²						MS ³					MS ⁴	
		parent ion	base peak m/z	secondary peaks						base peak m/z	secondary peaks				base peak m/z	
				m/z	intens ^a	m/z	intens	m/z	intens		m/z	intens	m/z	intens		
3-CQA (I)	3	353.7	191.5	179.5	49			135.7	10	85.6	127.1	93	172.3	60		
4-CQA (III)	3	353.5	173.5	179.5	68	191.6	20	135.7	14	93.3	111.2	47				
5-CQA (II)	3	353.3	191.5	179.5	5	161.4	2			85.7	127.1	93	172.2	75		
3-pCoQA (IV)	6	337.7	163.5							119.4	<i>b</i>					
4-pCoQA (VI)	6	337.5	173.5	163.6	7					93.5	111.0	60				
5-pCoQA (V)	6	337.7	191.6	163.3	5					85.6	127.0	97	172.2	95		
3-FQA (VII)	6	367.5	193.5	191.5	3					134.0	148.8	30			106.5	
4-FQA (IX)	3	367.4	173.5	191.8	87					93.3	111.5	44				
5-FQA (VIII)	3	367.6	191.6	173.6	6					85.6	127.1	100	172.2	90		

^a Intensity. ^b Signal too weak.Table 2. Negative Ion MS⁴ Data for Diacylchlorogenic Acids

compound	N	MS ¹		MS ²						MS ³					MS ⁴				
		parent ion	base peak m/z	secondary peaks						base peak m/z	secondary peaks				base peak m/z	secondary peaks			
				m/z	intens ^a	m/z	intens	m/z	intens		m/z	intens	m/z	intens		m/z	intens	m/z	intens
3,4-diCQA (X)	3	515.7	353.4	335.6	16	173.5	18			173.5	179.5	91	191.7	53	135.6	14	93.4	111.1	70
3,5-diCQA (XI)	3	515.2	353.5							191.5	179.5	53	173.5	8	135.6	12	85.5	127.0	95
4,5-diCQA (XII)	3	515.4	353.5							173.5	179.4	80	191.6	27	135.7	12	93.3	111.4	38
CFQA-1 (XIII)	3F,4CQA	6	529.8	353.2	366.9	89	335.4	73	349.4	48	173.6	179.4	85	191.5	27	135.8	11	134.1	148.9
CFQA-2 (XIV)	3C,4FQA	6	529.4	366.9	335.7	18	349.3	5			173.5	193.5	20				93.5	111.3	73
CFQA-3 (XV)	3F,5CQA	6	529.2	367.0	335.5	9					193.3	173.4	45				134.0	149.0	48
CFQA-4 (XVI)	3C,5FQA	6	529.3	353.2	366.8	45					191.5	179.5	56				127.2	172.4	68
CFQA-5 (XVII)	4F,5CQA	6	529.3	366.9							173.5	193.5	58				93.4	111.2	57
CFQA-6 (XVIII)	4C,5FQA	6	529.4	353.1	366.9	25					173.4	179.4	84	191.6	28	135.7	17	93.3	111.3


^a Intensity.

Figure 2. Specimen chromatogram (320 nm) of the coffee extract.

and cider (hard cider) sample were each examined on at least three separate occasions. Additional data were collected when the low concentration of the target molecule impaired the quality of the MS data (Tables 1 and 2). A specimen chromatogram (320 nm) for the coffee extract is shown in Figure 2. Because the content of *p*CoQA is very low in coffee and because 3-*p*-

CoQA (IV) elutes fused with 5-CQA (II), cider (hard cider) (Figure 3) was used as a convenient alternative source that gives superior mass spectra for this CGA subgroup.

5-CQA (II) was located on the chromatogram by comparison with a commercial standard. 3-CQA (I) and 4-CQA (III) were located using standards prepared from 5-CQA (II) by the




Figure 3. Specimen chromatogram (320 nm) of the cider (hard cider).

tetramethylammonium hydroxide interesterification method (15). 3,4-DiCQA (X), 3,5-diCQA (XI), 4,5-diCQA (XII), 5-*p*CoQA (V), and 5-FQA (VIII) were located by the use of NMR-characterized material isolated in previous studies (15). It was observed that on the phenylhexyl column packing, the monoacyl CGA eluted in the sequence 3-acyl, 5-acyl, and 4-acyl in contrast to the packings used previously in our laboratory, but which are no longer commercially available (12, 15–22). The elution sequence of the diCQA was unchanged (3,4-di, 3,5-di, and 4,5-diCQA).

When commercial standards were not available, peak identities were assigned primarily by means of their parent ion and supported by their UV spectrum and sequence of elution/retention time relative to 5-CQA using methods validated in our laboratory (8, 12, 15). As expected from numerous previous studies (2, 3, 8), selected ion monitoring (SIM) located three CQA (I–III), three *p*CoQA (IV–VI), three FQA (VII–IX), and three diCQA (X–XII). In addition, six putative CFQA, designated CFQA-1 to CFQA-6 (XIII–XVIII), were observed for the first time. Table 1 contains the summarized MSⁿ data for the monoacyl CGA and Table 2 the equivalent data for diacyl CGA.

In general, the base peaks were consistent both within and between analyses. However, when the mass spectrum for a particular substance included two ions of similar mean intensities, within-analysis experimental error dictated that in some individual MS scans one would be more intense and for other scans the reverse would be true. This phenomenon was encountered primarily when the signal intensity was lower, that is, with quantitatively minor components and/or higher order spectra. For example, the monoacyl CGA MS³ ions at $m/z \sim 85.6$ (Q₅) and at $m/z \sim 127.0$ (Q₇) are essentially co-equal in some spectra. However, in this particular case, the lower mass ion has been assigned consistently as the base peak because in the spectra of several compounds this was clearly the case. Fragment ions with intensities of <10% of the base peak have been reported only when they are needed for comparison. Selected diagnostic mass spectra are shown (Figures 5–9) to support the arguments advanced.

All monoacyl CGA gave the expected parent ion [monoacyl CGA – H]⁺. The fragment structures are presented in Figure 4. In the case of 3-*p*CoQA (IV) and 3-FQA (VII) the MS² and MS³ base peak ions were derived from the cinnamic acid moiety. For 3-*p*CoQA (IV), these ions are a [cinnamoyl – H]⁺ ion (A₃) and its decarboxylation product [cinnamoyl – CO₂ – H]⁺ (B₃), respectively. 3-FQA (VII) also yields a [cinnamoyl – H]⁺


ion (A₂), but the MS³ base peak is the “demethylated” ion (C) at $m/z \sim 134.0$. For the remaining seven monoacyl CGA the base peak ions were derived from the quinic acid moiety. Two distinct fragmentation pathways occur. One pathway gives [quinic acid – H]⁺ (Q₁) at MS² and a fragment (Q₅) at $m/z \sim 85.6$ in MS³; the other gives [quinic acid – H₂O – H]⁺ (Q₂) and an unassigned MS³ fragment at $m/z \sim 93.3$ (Figures 5 and 6). The MS¹ and MS² data are consistent with previous publications (8, 23); MS³ data have not been reported previously.

The diacyl CGA behaved similarly, giving the equivalent parent ion [diacyl CGA – H]⁺. All nine compounds examined lost either caffeic acid or ferulic acid, yielding a [diacyl CGA – cinnamoyl – H]⁺ ion as the MS² base peak. These ions are identical to the parent ions obtained from CQA and FQA, and it is clear that in general ions produced from diacyl CGA at MS⁽ⁿ⁺¹⁾ are identical to those produced from monoacyl CGA at MSⁿ—for example, compare Figures 5 and 7. CFQA-1 (XIII) and CFQA-3 (XV) closely resemble 3-FQA (VII), both yielding C, the demethylated (base peak), and the decarboxylated ions (B₁, B₂) at MS⁴. CFQA-4 (XVI) has an MS⁴ base peak (Q₇) at $m/z \sim 127.2$. For the remaining six diacyl CGA the MS³ base peak is one or the other of the quinic acid-derived ions that form the MS² base peaks of the monoacyl CGA. The base peaks reported for diCQA are consistent with previous publications (8, 23), but the fragment ions differ from those reported by Fang et al. (23). It must be noted that some of the mass assignments as printed in that paper appear to be in error. For example, [M – C₁ – H₂O]⁺ is given as m/z 353 when M = 516 and C₁ = 179 and has apparently been confused/transposed with [M – C₁]⁺ that is assigned as m/z 335. The term “chlorogenic acids” also is used confusingly when apparently “caffeic acid” is meant. Because there may also be less obvious errors, a more detailed comparison of their data and those reported here has not been attempted.

Key objectives of this investigation were to ascertain whether individual mono- or diacyl CGA could be identified unequivocally by their mass spectra and to seek to demonstrate the six CFQA that have been predicted as occurring in coffee beans.

Identification of Isomeric Monoacyl CGA. In zoom scan mode, the molecular masses predicted for the nine monoacyl CGA (I–IX) were consistently within 0.02 amu of the calculated accurate masses. In turboscan mode the mass predicted for 5-CQA (II) from the parent ion was 0.2 amu greater than the calculated accurate mass (353.10). The corresponding predictions for the remaining eight monoacyl CGA (I, III–IX) were higher by not more than 0.6 amu, and this level of imprecision is assumed for the fragment m/z values. It is clear from the data summarized in Table 1 that the parent ion clearly identifies the CGA subclass to which each monoacyl CGA belongs.

More importantly, it is easy to distinguish a 4-acyl CGA by its “dehydrated” MS² base peak Q₂ at $m/z \sim 173.5$ (Figure 5), supported by strong MS³ ions at $m/z \sim 93.3$ (unassigned) and Q₆ at $m/z \sim 111.2$ (Figure 6). In contrast, all three 5-acyl CGA (II, V, and VIII) and 3-CQA (I) produce an MS² base peak at $m/z \sim 191.5$ (Q₁), supported by strong MS³ ions at $m/z \sim 85.6$ (Q₅), ~ 127.0 (Q₇), and ~ 172.0 (Q₃). The distinctive MS² behavior of 4-CQA (III) has been reported previously (23), but the fragmentation could not be investigated further with the equipment available, and its structure–diagnostic value was not fully discussed. 3-*p*CoQA (IV) and 3-FQA (VII) are readily distinguished by their cinnamic acid-derived MS² base peaks (A₃ at $m/z \sim 163.5$ and A₂ at $m/z \sim 193.5$, respectively). 3-CQA (I) gives the same base peak as 5-CQA (II) but can be



Fragment	R	Cinnamic Acid	Accurate Mass
Q ₁			191.06
Q ₂			173.04
Q ₃			172.04
Q ₄	OH	Caffeic	335.08
	OCH ₃	Ferulic	349.08
Q ₅			85.03
Q ₆			111.04
Q ₇			127.04
A ₁	OH	Caffeic	179.04
A ₂	OCH ₃	Ferulic	193.04
A ₃	H	<i>p</i> -Coumaric	163.04
B ₁	OH	Caffeic	135.04
B ₂	OCH ₃	Ferulic	149.04
B ₃	H	<i>p</i> -Coumaric	119.04
C		Ferulic	134.04

Figure 4. Structure of quinic acid-derived and cinnamic acid-derived fragments.

distinguished from 5-CQA (**II**) by a comparatively intense caffeic acid-derived ion A₁ at $m/z \sim 179.5$ (mean intensity of 49% compared with 5%) that is reminiscent of A₃ produced from 3-*p*CoQA (**IV**) and A₂ produced from 3-FQA (**VII**). These data for 3-CQA (**I**) and 5-CQA (**II**) are identical to those reported by Fang et al. (23).

It is thus possible to discriminate between each of the nine monoacyl CGA (**I**–**IX**) on the basis of MS¹ and MS² spectra. The MS³ spectra provide confirmation of these assignments, but are not essential in making them. The potential diagnostic value of this distinctive MS behavior has not previously been recognized so far as we are aware (8). A hierarchical key to facilitate its exploitation is presented in Figure 10.

Identification of Isomeric DiCQA. The molecular masses predicted from the diCQA parent ions differed from the calculated accurate masses by not more than 0.03 amu in zoom scan mode and by not more than 0.5 amu in turboscan mode.

The *vic* diCQA [i.e., 3,4-diCQA (**X**) and 4,5-diCQA (**XII**)] give Q₂ as the MS³ base peak at $m/z \sim 173.5$ (Figure 7),

supported by strong MS⁴ ions at $m/z \sim 93.3$ (unassigned) and Q₆ at $m/z \sim 111.2$ (Table 2), as previously observed for 4-CQA (**III**), 4-*p*CoQA (**VI**), and 4-FQA (**IX**). These ions were not detectable in the spectrum of 3,5-diCQA (**XI**) (Figure 7), which gives Q₁ as an MS³ base peak at $m/z \sim 191.5$, supported by strong MS³ ions at $m/z \sim 85.6$ (Q₅), $m/z \sim 127.0$ (Q₇), and $m/z \sim 172.0$ (Q₃), as previously observed for 3-CQA (**I**) and 5-CQA (**II**). 3,5-DiCQA (**XI**) can thus be distinguished unequivocally from the *vic* diCQA (**X** and **XII**), and the ion at Q₂ at $m/z \sim 173.5$ in either an MS² or an MS³ spectrum is clearly diagnostic for substitution at position 4. There remains a need to determine which of the two *vic* diCQA has a substituent at position 3 and which at position 5.

The two *vic* diCQA isomers (**X** and **XII**) differ (Figure 7) with regard to the intensity of Q₄ the MS² “dehydrated” ion [CQA – H₂O – H⁺]⁺. In 3,4-diCQA (**X**) Q₄ is more intense (16% of base peak). In contrast, in 4,5-diCQA (**XII**) Q₄ is barely detectable (<5% of base peak). Similarly, the 3,4-isomer (**X**) (Figure 7; Table 2) produces Q₁ in MS³ and Q₆ in MS⁴ at approximately double the intensities of the 4,5-isomer (**XII**).




Figure 5. MS² spectra for isomeric caffeoylquinic acids.






Figure 6. MS³ spectra for isomeric caffeoylquinic acids.

Because this behavior was highly consistent over six separate determinations, it is sufficiently robust to discriminate between these two isomers.

As noted above, the MS² base peaks for the diCQA are identical to the parent ions for the CQA. The subsequent degradation of these ions will therefore be identical regardless of whether they have been derived from CQA or diCQA. Thus, by comparing the diCQA MS³ data (Figure 7; Table 2) with the CQA MS² data (Figure 5; Table 1) it is possible to define the precise structure of the ions responsible for the diCQA MS²

base peaks. With this information it becomes possible to specify which of the caffeoyl moieties is removed from the diCQA during MS¹ and which during MS².

Because the ion Q₂ at m/z ~173.5 is characteristic for an isomer substituted at position 4, the MS² base peak for the *vic* diCQA (X and XII) must be [4-CQA - H]⁺ rather than [3-CQA - H]⁺ or [5-CQA - H]⁺. It follows, therefore, that 3,4-diCQA (X) initially must lose the caffeoyl moiety at position 3, whereas 4,5-diCQA (XII) initially must lose that at position 5.

Figure 7. MS² and MS³ spectra for isomeric dicaffeoylquinic acids.Figure 8. MS² spectra for isomeric caffeoylferuloyl quinic acids.

Fragmentation of the MS² base peak for 3,5-diCQA (XI) yields (Figure 7) a comparatively intense [caffeoyl - H⁺]⁻ ion (A_1 = 50% of base peak). This is consistent with [3-CQA - H⁺]⁻ being the MS² base peak rather than [5-CQA - H⁺]⁻ where only a weak [caffeoyl - H⁺]⁻ ion (A_1 ~5% of base peak) would be expected (Table 1; Figure 5). These observations lead to the conclusion that the acylating residue at position 4 is the most difficult to remove, whereas that at position 5 is the easiest. This proposal is supported also by the observation (Figure 7; Table 2) that *vic* diCQA (X and XII)

do not give strong “dehydrated” ions (Q_4) at m/z ~335.5 that would otherwise be expected. These observations have been incorporated in the hierarchical key (Figure 10), and this has been used to interpret the spectra for the putative CFQA isomers (XIII–XVIII).

Characterization of Putative CFQA Isomers. Although previously only three CFQA isomers have been unequivocally characterized in any one coffee sample (12), in theory, one would expect six CFQA isomers (XIII–XVIII) to be present (12). SIM at m/z 529.0 ± 0.5 detected six peaks (Figure 8)




Figure 9. MS³ spectra of the isomeric caffeoylferuloylquinic acids.

with 325 nm λ_{max} that eluted between 57 and 65 min (Figure 2). These six compounds elute as three pairs with, in each case, the faster eluting component being present at a much smaller concentration than the slower eluting. The CFQA are minor components that, relative to unit concentration of 5-CQA (the major CGA in coffee), range from ≈ 0.004 unit (XV) to ≈ 0.08 unit (XVIII) assuming similar propensities to ionize.

In turboscan mode parent ions fall in the range m/z 529.2–529.9 compared with a calculated exact mass of 530.14. In zoom scan mode the predicted masses were within 0.1 amu of the calculated accurate masses. All six compounds show MS² base peaks (Figure 8) of either m/z 353.1–353.2 ([CQA – H⁺]⁺) or m/z 366.9–367.0 ([FQA – H⁺]⁺) and at MS³ (Figure 9) significant yields of A₁ ([caffeoyl – H⁺]⁺) or A₂ ([feruloyl – H⁺]⁺) consistent with these compounds being CFQA isomers.

The absence of the MS³ base peak Q₂ at m/z ~173.5 (Figure 9) clearly defines XV and XVI as lacking substitution at position 4. These isomers are, therefore, 3F,5CQA and 3C,5FQA. Similarly, the detection of Q₂ as the MS³ base peak clearly defines the four *vic* isomers as XIII, XIV, XVII, and XVIII.

The two forms of Q₄ at m/z 335.4–353.7 ([CQA – H₂O – H⁺]⁺) and m/z 349.3–349.4 ([FQA – H₂O – H⁺]⁺) are found in the MS² spectra (Figure 8) of XIII and XIV with intensities never less than 20% of the MS² base peaks. XVII and XVIII yield the same “dehydrated” ions with intensities never exceeding 6% of base peak. Therefore, by analogy with the behavior of the *vic* diCQA (Figure 7), it seems reasonable to conclude that XIII and XIV are the 3,4-substituted isomers and that XVII and XVIII are the 4,5-substituted isomers. Such assignments are also consistent with the known greater hydrophobicity of 4,5-substituted CGA compared with 3,4-substituted isomers (12, 15). It now remains to define which cinnamic acid is at which position.

Because XVIII yields an MS² base peak at m/z ~353.1 (Figure 8), this compound loses its feruloyl residue before its caffeoyl residue. Because during MS³ the base peak (Figure

9) at m/z ~173.5 (Q₂) is accompanied by an intense A₁ (70% of base peak) at m/z ~179.4 ([caffeoyl – H⁺]⁺), the MS² base peak must be [4-CQA – H⁺]⁺ rather than [4-FQA – H⁺]⁺. Accordingly, XVIII is 4C,5FQA.

It follows that if this assignment is correct, then XVII must be 4F,5CQA. A strong MS³ ion (Figure 9) at m/z ~193.5 (40% of the m/z 173.5 base peak) and the release of the caffeoyl residue at C5 before the feruloyl residue at C4 are consistent with these assignments. This order of elution is consistent also with previous studies (10, 11) in which 4F,5CQA (XVII) and 4C,5FQA (XVIII) had been isolated and characterized by NMR (but note that the original reports used non-IUPAC numbering).

XV yields an MS² base peak (Figure 8) at m/z ~367.0 indicating the loss of its caffeoyl residue before its feruloyl residue, suggesting that this compound is 3F,5CQA. This assignment is also supported by the observation of A₂ as the MS³ base peak at m/z ~193.3, a feature also of the MS² spectrum of 3-FQA (VII) and no other FQA isomer. In contrast, XV yields an MS² base peak at m/z ~353.2 indicating the loss of its feruloyl residue before its caffeoyl residue, implying consistently that this compound is 3C,5FQA (XVI). This isomer has a distinctive MS⁴ spectrum (base peak Q₇ at m/z ~127.2 and strong secondary ion Q₄ at m/z ~172.4), which are seen clearly also in the MS³ spectra of 3-CQA (I), 5-CQA (II), and 5-FQA (VIII) (Table 1).


XIV yields an MS² base peak at m/z ~366.9 indicating the loss of the caffeoyl residue prior to the feruloyl residue. This implies that the feruloyl residue is at the more stable location, which indicates that this compound is 3C,4FQA (XIV). If this assignment is correct, then XIII must be 3F,4CQA. However, XIII has distinctive behavior (Figure 8) that requires further comment. XIII gives an MS² base peak with a mean value ($N = 6$) of m/z 353.2 that is compatible with this assignment. However, it must be noted that it gives also a very strong MS² ion at m/z ~367.0 (mean intensity of 89% of base peak). In approximately half of the individual scans this ion is more

1. Parent ion m/z 337.5. ***p*-Coumaroylquinic acids. Go to 2**
 Parent ion m/z 353.5. **Caffeoylquinic acids. Go to 3**
 Parent ion m/z 367.5. **Feruloylquinic acids. Go to 4**
 Parent ion m/z 515.5. **Dicafeoylquinic acids. Go to 5**
 Parent ion m/z 529.5. **Caffeoylferuloylquinic acids. Go to 6**
2. MS² base peak m/z 163.5. **3-*p*-Coumaroylquinic acid (IV)**
 MS² base peak m/z 173.5. **4-*p*-Coumaroylquinic acid (VI)**
 MS² base peak m/z 191.5. **5-*p*-Coumaroylquinic acid (V)**
3. MS² base peak m/z 191.5, and relatively intense (*ca* 50% base peak) secondary ion at m/z 179.5. **3-Caffeoylquinic acid (I)**
 MS² base peak m/z 173.5. **4-Caffeoylquinic acid (III)**
 MS² base peak m/z 191.5, and weak or undetectable (<5% base peak) secondary ion at m/z 179.5. **5-Caffeoylquinic acid (II)**
4. MS² base peak m/z 193.5. **3-Feruloylquinic acid (VII)**
 MS² base peak m/z 173.5. **4-Feruloylquinic acid (IX)**
 MS² base peak m/z 191.5. **5-Feruloylquinic acid (VIII)**
5. MS² base peak m/z 353.5, MS³ base peak m/z 173.5, and comparatively intense (*ca* 20% of base peak) secondary ion at m/z 335.5. **3,4-Dicafeoylquinic acid (X)**
 MS² base peak m/z 353.5 and MS³ base peak m/z 191.5. **3,5-Dicafeoylquinic acid (XI)**
 MS² base peak m/z 353.5, MS³ base peak m/z 173.5, and comparatively intense (*ca* 20% of base peak) secondary ion at m/z 335.5. **4,5-Dicafeoylquinic acid (XII)**
6. MS³ base peak m/z 173.5. ***Vic*-Caffeoylferuloylquinic acids. Go to 7**
 MS³ base peak **not** m/z 173.5. **3,5-Caffeoylferuloylquinic acids Go to 10**
7. MS² secondary ions at m/z 335.5 or m/z 349.5 with intensities not less than *ca* 40% of base peak. **3,4-Caffeoylferuloylquinic acids. Go to 8**
 MS² secondary ions at m/z 335.5 or m/z 349.5 with intensities not more than *ca* 20% of base peak. **4,5-Caffeoylferuloylquinic acids. Go to 9**
8. MS² base peak at m/z 367.0 and MS³ secondary ion at m/z 193.5. **3-Caffeoyl-4-feruloylquinic acid (XIV)**
 MS² base peak at m/z 353.5 **or** m/z 367.0 with m/z 353.5 of near identical intensity, and MS³ secondary ion at m/z 179.5. **3-Feruloyl-4-cafeoylquinic acid (XIII)**
9. MS² base peak at m/z 367.0 and an intense (>50% of base peak) MS³ secondary ion at m/z 193.5. **4-Feruloyl-5-cafeoylquinic acid (XVII)**
 MS² base peak at m/z 353.5 and an intense (>50% of base peak) MS³ secondary ion at m/z 179.5. **4-Caffeoyl-5-feruloylquinic acid (XVIII)**
10. MS² base peak at m/z 367.0 and MS³ base peak at m/z 193.5. **3-Feruloyl-5-cafeoylquinic acid (XV)**
 MS² base peak at m/z 353.5 and an intense (>50% of base peak) MS³ secondary ion at m/z 179.5. **3-Caffeoyl-5-feruloylquinic acid (XVI)**


Figure 10. Hierarchical key for identification by LC-MSⁿ of mono- and diacylchlorogenic acids not substituted at position 1.

Scheme 1


Loss of caffeic acid in 5-CQA



Loss of caffeic acid in 3-CQA



Scheme 2



Scheme 3




Figure 11. Schemes of fragmentation.

intense than the ion at $m/z \sim 353.2$ and on these occasions is returned as the base peak. The distinctive MS⁴ ions at $m/z \sim 134.1$ (C) and $m/z \sim 148.9$ (B₂) seen previously in the MS³ spectrum of 3-FQA (VII) also suggest that XIII is indeed 3F,4CQA.

Such behavior is not seen in any other CFQA isomer, but it is consistent with the behavior of 3-FQA (VII) and 3F,5CQA (XV) discussed above. Such behavior, if it were to occur in the corresponding diCQA, could not be detected because both acylating residues have the same mass. However, closer examination of the MS³ spectra for 3,4-diCQA (X) reveals that in $\sim 50\%$ of the scans A₁ at $m/z \sim 179.5$ is more intense than the Q₂ at $m/z \sim 173.5$. This implies that such behavior is characteristic of acylation at C3 whichever cinnamic acid is involved and further strengthens the assignment of XIII as 3F,-4CQA.

Mechanisms of Fragmentation. The comparatively easy loss of the 5-acyl moiety, for example, from 5-CQA (II), can be rationalized by an abstraction of the proton from the C1 carboxyl while in an inverted chair conformation where the C1 carboxyl and the C5 caffeoyl have a 1,3-*syn*-diaxial arrangement (Figure 11, Scheme 1). The caffeoyl substituent in 3-CQA (I) has relative to the C1 hydroxyl a similar 1,3-*syn*-diaxial arrangement and can be lost by the same mechanism (Figure 11, Scheme 1). We expect the C5 hydroxyl to be lost more readily than the C1 hydroxyl due to a stronger hydroxyl-associated C—O bond α to COOH. The high pK of the hydroxyl ($pK > 15$) relative

to the carboxyl ($pK \approx 3.5$) explains why 3,5-diCQA (XI) loses the residue at C5 prior to the residue at C3. For loss of a substituent from C4 no suitable conformation can be found that would allow transfer of a proton required for the elimination step. In radical cation chemistry it has been argued that loss of water in a substituted cyclohexanol system occurs via abstraction of a C4 hydrogen in a boat conformation (24, 25). In the absence of a deuterated substrate we cannot exclude this stereochemical pathway for fragmentation. However, we believe that in radical anion chemistry any elimination pathway should lead to a more stable oxygen-centered anion rather than the carbon-centered anion that would be obtained as a result of such a *syn*-1,4 elimination. It should be noted, however, that although the rationale we have presented does not require such a *syn*-1,4 elimination pathway, all of our arguments would remain consistent and in complete agreement with the experimental observations if it were to occur.

It follows, therefore, that loss of water from the chlorogenic acids cannot involve the hydroxyls at C4 and C1, but could involve those at C3 or C5. From the foregoing, it is postulated that 4-CQA (III), for example, can lose water from either of two conformers (A or B) as illustrated in Figure 11 and Scheme 2. Although in the absence of the deuterated analogues it is not possible to establish which conformer is involved, we suggest that conformer A is preferred because it allows 1,2-acyl participation by the *trans*-vicinal acyl moiety on C4, thus facilitating the formation of a bicyclic oxonium radical anion

C by loss of OH from C5 and subsequent loss of the caffeoyl at C4. 1,2-Acyl participation is not only a powerful tool in synthetic chemistry, but common in mass spectrometric fragmentation (26, 27). In contrast, the loss of OH from C3 of conformer B would have to occur without 1,2-acyl participation and should be less favored. Further evidence for the importance of the 1,2-acyl participation hypothesis is given below.

Consistent with this proposal is the observation (Figure 5) that for 3-CQA (I) the ion Q_2 at $m/z \sim 173.5$ is only $\sim 5\%$ of the MS^2 base peak intensity; that is, water is not eliminated significantly during the fragmentation of this isomer in which the acyl residue is not *trans* relative to the vicinal hydroxyl and 1,2-acyl participation is thus not possible (Figure 11, Scheme 3). If 1,2-acyl participation were not involved in the fragmentation process, then 3-CQA (I) also should fragment via dehydration. In 5-CQA (II) the 5-acyl group is *trans* to the OH at C4. However, as argued before, loss of the C4 hydroxyl by 1,2-acyl participation is not observed because it would require abstraction of a *syn* proton that in the reactive conformation is unavailable in this particular isomer.

The increased intensity of the dehydrated peak in the MS^2 spectrum of 3,4-diCQA (X) as compared with 4,5-diCQA (XII) (Figure 7) further supports the above argument that loss of OH takes place from C5 preferentially by 1,2-acyl participation. Furthermore, loss of the caffeoyl group from C5 in 4,5-diCQA (XII), where 1,2-acyl participation can occur, is more facile than loss of caffeoyl from C3 in 3,4-diCQA (X), where 1,2-acyl participation is impossible.

The data obtained for the fragmentation of the diCQA indicate clearly that a caffeoyl residue is most easily removed from C5, somewhat less readily removed from C3, and not easily removed from C4. We have presented arguments based on the relative acidities of COOH and OH that perfectly explain these observations, but 1,2-acyl participation can be used as an alternative rationale. This increased stability of the C4 caffeoyl compared with the alternative 3- and 5-positions in 3,4-diCQA (X) and 4,5-diCQA (XII), respectively, where 1,2-acyl participation facilitates loss of 5-OH [in 3,4-diCQA (X)] or loss of 5-caffeoyl [in 4,5-diCQA (XII)], provides in summary a consistent rationale for all observed fragmentations.

However, comparisons of the fragmentation behavior of 3C,-4FQA (XIV) with 3F,4CQA (XIII) and of 3-CQA (I) with 3-FQA (VII) and 3-pCoQA (IV) suggest that for C3 the ease with which an acyl residue is removed depends also upon its identity. The fragmentation process has to be initiated by the capture of an electron by the aromatic side chain to produce a radical anion. If the caffeic ester is able to capture an electron more readily than the ferulic or *p*-coumaric ester, then the stereochemical relationships between the substituents might become less relevant than for the diCQA and CQA. For a caffeoyl residue, electron capture will give a radical anion that subsequently loses an H radical or proton. The *para* hydroxyl of a *vic* dihydroxyphenol such as caffeic acid has a bond dissociation enthalpy (BDE) of ~ 72 kcal mol⁻¹, lower than for isolated phenolic hydroxyls (28), and the anion/radical once formed is further stabilized by H-bonding to the adjacent hydroxyl (29, 30). The BDE for the *para* hydroxyl of a feruloyl or *p*-coumaroyl moiety is some 14 kcal mol⁻¹ higher and the anion/radical cannot hydrogen bond and is thus less stable. This stabilization available to the caffeoyl residue is sufficient, at least at C3, to account for its distinctive behavior compared with feruloyl or *p*-coumaroyl esters. That this effect is not apparent with 4-acyl-CGA suggests that the additional energy

required to remove a C4 caffeoyl moiety compared with a C3 caffeoyl moiety is somewhat greater than 14 kcal mol⁻¹.

Hierarchical Key to the Identification by LC-MSⁿ of Mono- and Diacyl CGA in Coffee. In conclusion, green robusta coffee beans contain six isomers of caffeoylferuloylquinic acid. Of greater importance, it is clear from the data in Tables 1 and 2, and the discussion above, that the 18 coffee chlorogenic acids examined (I–XVIII) give mass spectra that are amenable to logical interpretation, thus permitting their unequivocal discrimination. Such LC-MSⁿ procedures are thus potentially of much wider application because CGA are widespread in plant material and commercial standards are restricted to one or two individual compounds (8). The hierarchical key (Figure 10) is proposed to facilitate this process. All masses are approximate (± 0.3 amu). It must be noted however, that the key as presented is not intended to accommodate the 1-substituted CGA (as found in some Asteraceae) and has not been evaluated on CGA that contain other than the “normal” isomer of quinic acid (8, 31, 32).

LITERATURE CITED

- (1) IUPAC. Nomenclature of cyclitols. *Biochem. J.* **1976**, *153*, 23–31.
- (2) Clifford, M. N. Chlorogenic acids and other cinnamates—nature, occurrence, dietary burden, absorption and metabolism. *J. Sci. Food Agric.* **2000**, *80*, 1033–1042.
- (3) Clifford, M. N. Chlorogenic acids and other cinnamates—nature, occurrence and dietary burden. *J. Sci. Food Agric.* **1999**, *79*, 362–372.
- (4) Homma, S. Chemistry II. Non-volatile compounds, Part II. In *Coffee: Recent Developments*; Clarke, R. J., Vitzthum, O. G., Eds.; Blackwell Science: Oxford, U.K., 2001; pp 50–67.
- (5) Clifford, M. N. The nature of chlorogenic acids—are they advantageous compounds in coffee? In *Dix-septième Colloque Scientifique Internationale sur le Café*; ASIC: Paris, France, 1998; pp 79–91.
- (6) Clifford, M. N. Chlorogenic acids. In *Coffee I. Chemistry*; Clarke, R. J., Macrae, R., Eds.; Elsevier Applied Science: London, U.K., 1985; pp 153–202.
- (7) Clifford, M. N. The health effects of tea and tea components. Appendix 1. A nomenclature for phenols with special reference to tea. *Crit. Rev. Food Sci. Nutr.* **2001**, *41*, 393–397.
- (8) Clifford, M. N. The analysis and characterisation of chlorogenic acids and other cinnamates. In *Methods in Polyphenol Analysis*; Santos-Buelga, C., Williamson, G., Eds.; Royal Society of Chemistry: Cambridge, U.K., 2003; in press.
- (9) Corse, J.; Lundin, R. E.; Waiss, A. C. Identification of several components of isochlorogenic acid. *Phytochemistry* **1965**, *4*, 527–529.
- (10) Iwahashi, H.; Morishita, H.; Osaka, N.; Kido, R. 3-*O*-Feruloyl-4-*O*-caffeoylquinic acid from coffee beans. *Phytochemistry* **1985**, *24*, 630–632.
- (11) Morishita, H.; Iwahashi, H.; Kido, R. 3-*O*-Caffeoyl-4-*O*-feruloylquinic acid from green robusta coffee beans. *Phytochemistry* **1986**, *25*, 2679–2680.
- (12) Clifford, M. N.; Kellard, B.; Birch, G. G. Characterisation of caffeoylferuloylquinic acids by simultaneous isomerisation and transesterification with tetramethylammonium hydroxide. *Food Chem.* **1989**, *34*, 81–88.
- (13) Balyaya, K. J.; Clifford, M. N. Individual chlorogenic acids and caffeine contents in commercial grades of wet and dry processed Indian green robusta coffee beans. *J. Food Sci. Technol. Mysore* **1995**, *32*, 104–108.
- (14) Egan, H.; Kirk, R. S.; Sawyer, R. *Pearson's Chemical Analysis of Foods*; Churchill Livingstone: London, U.K., 1981.
- (15) Clifford, M. N.; Kellard, B.; Birch, G. G. Characterisation of chlorogenic acids by simultaneous isomerisation and transesterification with tetramethylammonium hydroxide. *Food Chem.* **1989**, *33*, 115–123.

- (16) Clifford, M. N.; Ramirez-Martinez, J. R. Phenols and caffeine in wet processed coffee beans and pulp. *Food Chem.* **1991**, *40*, 35–42.
- (17) Clifford, M. N.; Ramirez-Martinez, J. R. Chlorogenic acids and purine alkaloid content of Maté (*Ilex paraguariensis*) leaf and beverage. *Food Chem.* **1990**, *35*, 13–21.
- (18) Clifford, M. N.; Williams, T.; Bridson, D. Chlorogenic acids and caffeine as possible taxonomic criteria in *Coffea* and *Psilanthus*. *Phytochemistry* **1989**, *28*, 829–838.
- (19) Clifford, M. N.; Jarvis, T. The chlorogenic acids content of green robusta coffee beans as a possible index of geographic origin. *Food Chem.* **1988**, *29*, 291–298.
- (20) Clifford, M. N.; Shutler, S.; Thomas, G. A.; Ohiokpehai, O. The chlorogenic acids content of coffee substitutes. *Food Chem.* **1987**, *24*, 99–107.
- (21) Clifford, M. N.; Kazi, T. The influence of coffee bean maturity on the content of chlorogenic acids, caffeine and trigonelline. *Food Chem.* **1987**, *26*, 59–69.
- (22) Clifford, M. N. Coffee bean dicaffeoylquinic acids. *Phytochemistry* **1986**, *25*, 1767–1769.
- (23) Fang, N.; Yu, S.; Prior, R. L. LC/MS/MS characterization of phenolic constituents in dried plums. *J. Agric. Food Chem.* **2002**, *50*, 3579–3585.
- (24) Brion, C. I.; Hall, L. D. Studies of stereoisomerism by photo-sensitisation mass spectrometry. Fragmentation of *cis* and *trans* 4-*tert*-butylcyclohexanols. *J. Am. Chem. Soc.* **1966**, *88*, 3661–3662.
- (25) Green, M., M. Mass spectrometry—a sensitive probe of molecular geometry. *Pure Appl. Chem.* **1978**, *50*, 185–196.
- (26) Eames, J.; Kuhnert, N.; Warren, S. The scope and limitations of the [1,2]-alkylsulfonyl (SMe, SEt, and SCH₂Ph) and sulfanyl (SH) migration in the stereospecific synthesis of substituted tetrahydrofurans. *J. Chem. Soc., Perkin Trans. I* **2001**, 1504–1511.
- (27) Eames, J.; Kuhnert, N.; Warren, S. The scope and limitation of the [1,4]-Sbenzyl participation and debenzylolation in the stereochemically controlled synthesis of substituted thiolanes. *J. Chem. Soc., Perkin Trans. I* **2001**, 138–143.
- (28) Wright, J. S.; Johnson, E. R.; DiLabio, G. A. Predicting the activity of phenolic antioxidants: theoretical method, analysis of substituent effects, and application to major families of antioxidants. *J. Am. Chem. Soc.* **2001**, *123*, 1173–1183.
- (29) Alluis, B.; Dangles, O. Quercetin (2-(3,4-dihydroxyphenyl)-3,5,7-trihydroxy-4*H*-1-benzopyran-4-one) glycosides and sulfates: Chemical synthesis, complexation, and antioxidant properties. *Helv. Chim. Acta* **2001**, *84*, 1133–1156.
- (30) Lucarini, M.; Mugnaini, V.; Pedulli, G. F. Bond dissociation enthalpies of polyphenols: The importance of cooperative effects. *J. Org. Chem.* **2002**, *67*, 928–931.
- (31) Haribal, M.; Feeny, P.; Lester, C. C. Caffeoylcyclohexane-1-carboxylic acid derivative from *Asimina triloba*. *Phytochemistry* **1998**, *49*, 103–108.
- (32) Wang, Y.; Hamburger, M.; Gueho, J.; Hostettmann, K. Cyclohexanecarboxylic acid derivatives from *Psiadia trinervia*. *Helv. Chim. Acta* **1992**, *75*, 269–275.

Received for review December 3, 2002. Revised manuscript received March 7, 2003. Accepted March 8, 2003.

JF026187Q