



Restoring working forests in human dominated landscapes of tropical South Asia: An introduction



Mark S. Ashton^{a,*}, Uromi M. Goodale^b, Kamal S. Bawa^c, Peter S. Ashton^d, J. David Neidel^e

^a School of Forestry and Environmental Studies, Yale University, New Haven, CT 06511, USA

^b Key Laboratory of Tropical Forest Ecology, Xishuangbanna Tropical Botanical Garden, Chinese Academy of Sciences, Menglun, Mengla, Yunnan 666303, PR China

^c Department of Biology, University of Massachusetts, Boston, MA 02125, USA

^d Department of Organismic and Evolutionary Biology, Harvard University, Cambridge, MA 02138, USA

^e Environmental Leadership and Training Initiative, School of Forestry and Environmental Studies, Yale University, New Haven, CT 06511, USA

ARTICLE INFO

Article history:

Available online 14 June 2014

Keywords:

Bird flocks
Forest carbon
Land use
Non-timber forest products
Restoration
Sacred groves

ABSTRACT

The resource issues around restoring human dominated landscapes in tropical South Asia are complex and can be divided into topics concerning forest fragmentation and restoration. Issues that focus on effects of forest fragmentation include studies that show declines in forest structure and standing carbon stocks within forest fragments as compared to contiguous forest; changes in bird composition and flock density in relation to land use – especially between forests, the forest buffer zones and agricultural lands (tea, coffee); and the potential origins of many sacred groves as forest fragments and their future roles as cultural, social and ecological centers of reforestation within deforested landscapes. Studies that focus on restoration have demonstrated the benefits of plantations as mechanisms to establish second growth forests and native species plantings for both economic and conservation purposes; the incorporation of indigenous plants that produce non-timber forest products in forest restoration programs; and the control of fire, soil preparation and protection from herbivory as treatments that can facilitate natural forest regeneration in montane grasslands.

© 2014 Elsevier B.V. All rights reserved.

1. Introduction

Tropical forests provide a variety of goods and services to humanity, including the maintenance of biodiversity, carbon sequestration, watershed protection, timber and non-timber forest products and open space recreation (Myers 1988; Chopra 1993; Godoy and Bawa, 1993; Balmford et al., 2002; MEA 2003; Ricketts et al., 2004; Wong and Yu, 2005; Foley et al., 2007; Dasgupta, 2010; Rands et al., 2010; Gamfeldt et al., 2013). Tropical South Asia comprises India and Sri Lanka. Tropical South Asia has high population density, and ancient cultures and land use histories. It was one of the first regions in the tropics that underwent large land clearance of forests (Goldewijk, 2001) for agriculture during the colonial era under the British from 1850 to 1950 mainly due to land conversion for plantation crops. This continued in the post-colonial era from 1950 to 1970 primarily for development (Vincent, 1883; Wimaladharm, 1977; Flint, 1994; Wickramagamage, 1998; Kurosaki, 2009; Näsström and Mattsson, 2011). Forest conversion therefore started over

150 years ago and, unlike many regions, progressed at lower rates of deforestation for a longer period (Kurosaki, 2009; Näsström and Mattsson, 2011). Many studies within the region have highlighted the damaging effects on biodiversity and other forest values because of land use policies that favored extensive forest conversion to commercial tree and shrub crops such as tea, rubber and coffee, or a few favored timber species (Flint, 1994; Dixon et al., 1994; Wickramagamage, 1998).

In the last thirty years, scientists and policy makers have recognized the important role that tropical forests play in maintaining biodiversity and therefore providing a multitude of ecosystem services (Balmford et al., 2002; Gibson et al., 2011), sequestering carbon (Brown and Lugo, 1982; Phillips et al., 1998; Lewis et al., 2009), and providing clean surface water supplies for downstream irrigated agriculture and drinking water (Pringle, 2001; Foley et al., 2005). The Intergovernmental Panel on Climate Change has proposed reforestation of tropical abandoned agricultural land and pastures as a strategy to increase carbon sequestration and help mitigate anthropogenic carbon emissions (Ding et al., 2001). Through reforestation ecosystems are able to recover function and biodiversity and serve as potential sinks for atmospheric carbon by storage in both plant

* Corresponding author. Tel.: +203 432 9835.

E-mail address: mark.ashton@yale.edu (M.S. Ashton).

biomass and soil organic matter (Lamb et al., 2005). In addition, reforestation has long been recognized for its potential to reduce erosion locally and to impact hydrological cycles (Eshleman, 2004; Ellison et al., 2012).

Plantation forestry has played a major role in the reforestation history of South Asia. It has a complex history of trial and error, dominated by monoculture plantations of exotic species such as *Eucalyptus*, *Acacia* and *Pinus*. The globalization of *Eucalyptus* increased with planting and success rates in the tropics from 1960 to 1990 (FAO, 1985; Saxena et al., 1993; Bennett 2010). However, criticism of planting *Eucalyptus* was vocal in the 1980s during the intense period of planting in India and Thailand (Bennett, 2010). At the same time, a mix of state, government, industrial cooperation, small holder land ownership and community-based reforestation programs were implemented within degraded forest areas and shrub and grasslands that were previously covered by forests, resulting in further experimentation with *Eucalyptus* as well as other exotics such as *Acacia* and *Pinus* species (Rai, 1999; Pomeroy et al., 2003; Ramachandra et al., 2004). Although exotic species may greatly out produce native tree species, especially under short rotations (Nath et al., 1990), exotic plantations in some cases had detrimental effects on hydrology for adjacent agriculture, and provided only single values – timber for the State (Kumar et al., 2003). Small-holders and subsistence farmers do not like the exotic plantations chiefly because none could be used for their benefit (FAO, 1985). Further, they have been assumed to have low conservation value (Lamb, 1998; Chazdon, 2008). In recent years, however, efforts have increased to better understand the processes that govern restoration of tropical degraded land (see. e.g. Yirdaw, 2001; Parrotta and Knowles, 2002; Vincent and Davies, 2003; Sharam et al., 2006; Slocum et al., 2006; Goodale et al., 2012; Salek and Vylupek, 2012) and exotic timber plantations are recognized as one potential approach (Lamb, 1998; Ashton et al., 1998; Lamb et al., 2005; Wishnie et al., 2007).

Worldwide, integrated approaches, although unlikely to achieve complete restoration, are now favored. These approaches range from using native species, landscape design, targeting site matching in species selection, using species mixtures, silvicultural management for greater productivity, to fostering understory diversity (see. e.g. Ashton et al., 1997a; Menalled et al., 1998; Miyawaki, 1999; Mark, 2007; Benayas et al., 2008; Chazdon, 2008; Cole et al. 2010; Park et al., 2010; Richards and Schmidt 2010). In using any of these approaches for reforestation and land restoration in South Asia, and in particular the Western Ghats and Sri Lanka, there has been a shift towards integrating native tree species (Raman et al., 2009; Nagaraja et al., 2011; Chetana and Ganesh, 2012). In fact, native species are now often promoted as being more ecologically and socially sustainable than exotic species for a series of reasons. First, recent studies have highlighted the importance of native trees for agricultural communities, emphasizing the goods and services that these trees provide to local farmers (Walters, 1997; Aguilar and Condit, 2001; DeClerck et al., 2006; Garen et al., 2009). Second, much work has been done on the recovery of second growth forest as a way of combating forest fragmentation and in the development of corridors for ecosystem restoration for local communities (Chazdon, 2003, 2008).

Given these trends, the goal of this special issue in Forest Ecology and Management is to review the scientific data and experiences of more than twenty-five years of research relating to the ecology and management of native tree species and second growth forests in order to inform reforestation and land restoration activities in the tropical South Asian region. The papers within this special issue cover largely ecological and silvicultural

topics of reforestation and restoration. The issue is dedicated to Professors Savithri and Nimal Gunatilleke of the University of Peradeniya, Sri Lanka. Building on the rich tradition of village home garden woodlands in their nation, Sri Lanka, they with their students have applied continuing research, into the landscape ecology and historical biogeography of its rich wet zone flora, in path breaking experiments on forest restoration in the context of the contemporary village economy (Gunatilleke et al., 1995; Ashton et al., 1997a; Ashton et al., 1997b; Gunatilleke et al., 2004; Tennakoon et al., 2005; Gunaratne et al., 2010; Gunaratne et al., 2011).

2. The papers in the special issue: organization and content

This special issue comprises six papers. They cover topics around tropical forest restoration that range from ecological aspects of native species reforestation, effects of land use on birds flocks and diversity, estimates of carbon in relation to forest fragmentation and underlying drivers of forest diversity in sacred forests.

In the first paper, Gunaratne and colleagues provided a synthesis of their work investigating the ecological barriers to tropical montane forest succession on anthropogenic grasslands (Gunaratne et al., 2014). Their findings suggest that forest regeneration on grasslands is impeded by limited seed dispersal, herbivory of seedlings that do establish and fire. Planting studies show that tree seedlings do better when grasses are removed and herbivores are excluded. *Macaranga indica* and *Symplocos cochinchinense* are two early successional pioneers that are potential nurse trees; planted on grasslands, they facilitate other tree regeneration beneath their canopy. Other potential techniques that they suggest include the creation of fire breaks, and scarification and exposure of mineral soil of grassland adjacent to forest patches. In the second paper, Mark Ashton and colleagues describe the use of pine plantations as a technique, both to secure the natural recruitment of native tree regeneration beneath the pine canopy and to facilitate the plantings of native trees (Ashton et al., 2014a). They provide a case for restoration for both conservation and utilitarian values, and make a strong economic case for the economic restoration of native species in the everwet regions of South Asia in comparison to the intensive cultivation of tea.

Rural communities have traditionally valued forests for a diversity of products and services, with timber serving a minor role. The demand for 'non-timber forest products' (NTFPs) must change from subsistence to monetary values, given the development of the region, if they are to be included in future values harvested from the forest. In the third paper, Ashton and colleagues review the use and cultivation of exemplar non-timber forest plants that yield a mix of products comprising resins, latexes, sugars, medicines, fibers, fruits and nuts (Ashton et al., 2014b). The paper provides ways of incorporating NTFP's into restoration plantings. In particular, the major species identified to have future commercial values in plantation systems include *Aegle marmelos*, *Calamus* spp., *Caryota urens*, *Coscinium fenestratum*, *Diospyros melanoxylon*, *Elettaria cardamomum*, *Gyrinops walla*, *Phyllanthus emblica*, *Santalum album*, and *Vateria* spp.

Forest fragments are thought to store a disproportionate amount of some of the remaining terrestrial carbon in degraded and deforested landscapes. Anand et al. (2014) evaluate the change in tree allometry and forest structure in forest fragments of the Western Ghats. They compared above-ground carbon stocks in evergreen forest fragments to stocks in relatively undisturbed, contiguous forests. Their results suggest forest

fragments store 40% less carbon per hectare above ground than contiguous forests. They relate the lower carbon estimates to change in forest stature, with decline if canopy height and increased mortality from edge effects, including the eventual death of surviving large trees. While improved protection of fragments might play an important role in mitigating carbon loss, a more promising approach is to actively manage restored fragments focusing on increasing stand density, height to diameter ratios, and late successional species composition.

The response of birds and mixed-species bird flocks to forest fragmentation has been investigated by Goodale et al. (2014). Their work has been carried out over the past ten years in South Asia. They rarely found flocks in agricultural areas but the numbers of flocks in buffer areas was similar to that in intact forests, although buffer flocks were smaller and had a different composition. Considering buffer transects alone, pine plantations retained fewer forest interior species in flocks than did broad-leaved forests, while small areas of agriculture and abandoned agriculture attracted open-landscape species. Though clearly not equivalent to protected forests, degraded forests and agroforests in buffer areas still hold some conservation value. Mixed-species flocks are particularly important as vehicles for retaining forest species in human-modified habitats.

South Asia still retains more than one thousand forest fragments held sacred by village communities and religious associations. The last paper in this special issue, by Bagwat et al. (2014) concerns cultural drivers of reforestation in tropical forest groves of the Western Ghats of India. Many of these sacred forest groves are considered ancient woodlands, but there is very little information on their origins. Based on palaeoecological reconstruction of two such sacred forest patches, Bhagwat and associates examine the historical literature in a search for linkages between the 'social' and the 'ecological' systems of sacred forest groves. They conclude that some at least of these forest fragments transitioned from grassland to tree-covered landscapes fairly recently in paleo-geographic time. This finding challenges the common perception that sacred forest groves are remnants of once-continuous aboriginal forest; indicating that they may often instead be regenerated forest patches. This raises questions concerning appropriate incentives for encouraging reforestation.

3. Future research

Though considerable research has been done within the region on forest restoration within degraded and deforested lands more needs to be done. In particular silvicultural studies on the domestication of promising non-timber forest products needs to continue. Certain forest types, especially montane forests and dry and semi-evergreen tropical forests, have been poorly studied in South Asia. Little work has been done on understanding their forest dynamics, and in then using this knowledge to test reforestation techniques for native species plantations and second growth forests.

Further studies need to be done on understanding the social and cultural dimensions of reforestation. Reforestation can only be successful if social desirable and acceptable to various stakeholders that have both local and national interests. What kinds of markets, incentives in land-use regulation, taxation and land tenure are needed to facilitate reforestation with native species for state and local values.

Lastly, more studies need to be done on conservation and ecosystem service values that reforestation and existing forest fragments in deforested regions can have. Understanding the role that reforestation can play in influencing climate, and affecting

hydrology and soil productivity has been done elsewhere, but studies in this region are lacking and need to be done to understand the specifics of the region. Similarly work has been done on understanding wildlife behaviors and population demographics but more work needs to be done in understanding wildlife use of forest fragments, corridors and new forests in human dominated landscapes.

4. Acknowledgments

We would like to thank the authors for putting together the papers that make up this special issue. We also wish to acknowledge support from the Environmental Leadership and Training Initiative (ELTI: <https://environment.yale.edu/elti/>) of Yale University that hosting and the University of Peradeniya and Peradeniya Botanical Gardens for facilitating the conference that stimulated the idea of publishing the presentations. ELTI is a capacity building organization devoted to training environmental professionals from tropical forest countries around forest restoration. The program is supported by a generous gift from the Arcadia Fund.

References

- Aguilar, S., Condit, R., 2001. Use of native tree species by an Hispanic community in Panama. *Econ. Bot.* 55, 223–235.
- Anand, M.O., Kumar, V., Sankaran, M., 2014. Altered stand density and tree allometry reduce carbon storage in evergreen forest fragments in India's Western Ghats. *Forest Ecol. Manage.* 329, 30–36.
- Ashton, P.M.S., Gamage, S., Gunatilleke, I.A.U.N., Gunatilleke, C.V.S., 1997a. Restoration of a Sri Lankan rainforest: using Caribbean pine *Pinus caribaea* as a nurse for establishing late-successional tree species. *J. Appl. Ecol.* 34, 915–925.
- Ashton, P.M.S., Samarasinghe, S.J., Gunatilleke, I.A.U.N., Gunatilleke, C.V.S., 1997b. Role of legumes in release of successional arrested grasslands in the Central Hills of Sri Lanka. *Restor. Ecol.* 5, 36–43.
- Ashton, P.M.S., Gamage, S., Gunatilleke, I.A.U.N., Gunatilleke, C.V.S., 1998. Using Caribbean pine to establish a mixed plantation: testing effects of pine canopy removal on plantings of rain forest tree species. *Forest Ecol. Manage.* 106, 211–222.
- Ashton, M.S., Gunatilleke, C.V.S., Gunatilleke, I.A.U.N., Singhakumara, B.M.P., Gamage, S., Shibayama, T., Tomimura, C., 2014a. Restoration of rain forest beneath pine plantations: a relay floristic model with special application to South Asia. *Forest Ecol. Manage.* 329, 351–359.
- Ashton, M.S., Gunatilleke, C.V.S., Gunatilleke, I.A.U.N., Tennakoon, K., Ashton, P.S., 2014b. Use and cultivation of plants that yield products other than timber from South Asian tropical forests, and their potential in forest restoration. *Forest Ecol. Manage.* 329, 360–374.
- Bagwat, S.A., Economou, A., Nogue, S., Willis, K.J., 2014. Cultural drivers of reforestation in tropical forest groves of the Western Ghats of India. *Forest Ecol. Manage.* 329, 393–400.
- Balmford, A., Bruner, A., Cooper, P., Costanza, R., Farber, S., Green, R.E., Jenkins, M., Jefferiss, P., Jessamy, V., Madden, J., Munro, K., Myers, N., Naem, S., Paavola, J., Rayment, M., Rosendo, S., Roughgarden, J., Trumper, K., Turner, R.K., 2002. Economic reasons for conserving wild nature. *Science* 297, 950–953. <http://dx.doi.org/10.1126/science.1073947>.
- Benayas, J.M.R., Bullock, J.M., Newton, A.C., 2008. Creating woodland islets to reconcile ecological restoration, conservation, and agricultural land use. *Front. Ecol. Environ.* 6, 329–336.
- Bennett, B.M., 2010. The El Dorado of forestry: the eucalyptus in India, South Africa, and Thailand, 1850–2000. *Int. Rev. Soc. Hist.* 55, 27–50. <http://dx.doi.org/10.1017/S0020859010000489>.
- Brown, S., Lugo, A.E., 1982. The storage and production of organic matter in tropical forests and their role in the global carbon cycle. *Biotropica* 14, 161–187.
- Chazdon, R.L., 2003. Tropical forest recovery: legacies of human impact and natural disturbances. *Perspect. Plant Ecol.* 6, 51–71.
- Chazdon, R.L., 2008. Beyond deforestation: restoring forests and ecosystem services on degraded lands. *Science* 320, 1458–1460. <http://dx.doi.org/10.1126/science.1155365>.
- Chetana, H.C., Ganesh, T., 2012. Importance of shade trees (*Grevillea robusta*) in the dispersal of forest tree species in managed tea plantations of southern Western Ghats, India. *J. Trop. Ecol.* 28, 187–197.
- Chopra, K., 1993. The value of non-timber forest products: an estimation for tropical deciduous forests in India. *Econ. Bot.* 47, 251–257.
- Cole, R.J., Holl, K.D., Zahawi, R.A., 2010. Seed rain under tree islands planted to restore degraded lands in a tropical agricultural landscape. *Ecol. Appl.* 20, 1255–1269.
- Dasgupta, P., 2010. Nature's role in sustaining economic development. *Philos. T. Roy. Soc. B* 365, 5–11.

- DeClerck, F., Ingram, J.C., Rumbaitis del Rio, C.M., 2006. The role of ecological theory and practice in poverty alleviation and environmental conservation. *Front Ecol. Environ.* 4, 533–540.
- Ding, Y.D.J.G., Griggs, D.J., Noguer, M., van der Linden, P.J., Dai, X., Maskell, K., Johnson, C.A., 2001. *Climate change 2001: the scientific basis*, vol. 881. Cambridge university press, Cambridge.
- Dixon, R.K., Solomon, A.M., Brown, S., Houghton, R.A., Trexler, M.C., Wisniewski, J., 1994. Carbon pools and flux of global forest ecosystems. *Science* 263, 185–190.
- Ellison, D., Futter, M.N., Bishop, K., 2012. On the forest cover–water yield debate: from demand- to supply-side thinking. *Global Change Biol.* 18, 806–820. <http://dx.doi.org/10.1111/j.1365-2486.2011.02589.x>.
- Eshleman, K.N., 2004. Hydrological consequences of land use change: a review of the state-of-science. *Geophysical Monograph Series* 153. American Geophysical Union. DOI: 10.1029/153GM03.
- FAO (Food and Agriculture Organization), 1985. The ecological effects of *Eucalyptus*. FAO Forestry Paper.
- Flint, E.P., 1994. Changes in land use in South and Southeast Asia from 1880 to 1980: a database prepared as part of a coordinated research program on carbon fluxes in the tropics. *Chemosphere* 29, 1015–1062.
- Foley, J.A., DeFries, R., Asner, G.P., Barford, C., Bonan, G., Carpenter, S.R., Chapin, F.S., Coel, M.T., Daily, G.C., Gibbs, H.K., Helkowski, J.H., Holloway, T., Howard, E.A., Kucharik, C.J., Monfreda, C., Patz, J.A., Prentice, I.C., Ramankutty, N., Snyder, P.K., 2005. Global consequences of land use. *Science* 309, 570–574.
- Foley, J.A., Asner, G.P., Costa, M.H., Coe, M.T., DeFries, R., Gibbs, H.K., Howard, E.A., Olson, S., Patz, J., Ramankutty, N., Snyder, P., 2007. Amazonia revealed: forest degradation and loss of ecosystem goods and services in the Amazon Basin. *Front. Ecol. Environ.* 5, 25–32.
- Gamfeldt, L., Snäll, T., Bagchi, R., Jonsson, M., Gustafsson, L., Kjellander, P., Ruiz-Jaen, M.C., Fröberg, M., Stendahl, J., Philipson, C.D., Mikusiński, G., Andersson, E., Westerlund, B., Andrén, H., Moberg, F., Moen, J., Bengtsson, J., 2013. Higher levels of multiple ecosystem services are found in forests with more tree species. *Nat. Commun.* 4, 1340.
- Garen, E.J., Saltonstall, K., Slusser, J.L., Mathias, S., Ashton, M.S., Hall, J.S., 2009. An evaluation of farmers' experiences planting native trees in rural Panama: implications for reforestation with native species in agricultural landscapes. *Agroforest. Syst.* 76, 219–236.
- Gibson, L., Lee, T.M., Koh, L.P., Brook, B.W., Gardner, T.A., Barlow, P., Peres, C.A., Bradshaw, C.J.A., Laurance, W.F., Lovejoy, T.E., Sodhi, N.S., 2011. Primary forests are irreplaceable for sustaining tropical biodiversity. *Nature* 478, 378–381.
- Godoy, R.A., Bawa, K.S., 1993. The economic value and sustainable harvest of plants and animals from the tropical forest: assumptions, hypotheses, and methods. *Econ. Bot.* 47, 215–219.
- Goldewijk, K.K., 2001. Estimating global land use change over the past 300 years: the HYDE database. *Global Biogeochem. Cy* 15, 417–433.
- Goodale, U.M., Berlyn, G.P., Ashton, M.S., Gregoire, T.G., Sinhakumara, B.M.P., Tennakoon, K.U., 2012. Disturbance and tropical pioneer species: patterns of association across life history stages. *Forest Ecol. Manage.* 227, 54–66.
- Goodale, E., Kotagama, S.W., Shankar Raman, T.R., Sidhu, S., Goodale, U.M., Parker, S., Chen, J., 2014. The response of birds and mixed bird flocks to human-modified landscapes in Sri Lanka and southern India. *Forest Ecol. Manage.* 329, 384–392.
- Gunaratne, A.M.T.A., Gunatilleke, C.V.S., Gunatilleke, I.A.U.N., Weerasinghe, H.M.S.P.M., Burslem, D.F.R.P., 2010. Barriers to tree seedling emergence on human-induced grasslands in Sri Lanka. *J. Appl. Ecol.* 47, 157–165.
- Gunaratne, A.M.T.A., Gunatilleke, C.V.S., Gunatilleke, I.A.U.N., Madawala Weerasinghe, H.M.S.P., Burslem, D.F.R.P., 2011. Release from root competition promotes tree seedling survival and growth following transplantation into human-induced grasslands in Sri Lanka. *Forest Ecol. Manage.* 262, 229–236.
- Gunaratne, A.M.T.A., Gunatilleke, C.V.S., Gunatilleke, I.A.U.N., Madawala, H.M.S.P., Burslem, D.F.R.P., 2014. Overcoming the ecological barriers of tropical montane forest succession on anthropogenic grasslands: synthesis and future prospects. *Forest Ecol. Manage.* 329, 340–350.
- Gunatilleke, C.V.S., Gunatilleke, I.A.U.N., Ashton, P.M.S., 1995. Rain forest research and conservation: the Sinharaja experience in Sri Lanka. *Sri Lanka Forest.* 22, 49–60.
- Gunatilleke, C.V.S., Gunatilleke, I.A.U.N., Ethugala, A.U.K., Weerasekara, N.S., Esufali, S., Ashton, P.S., Ashton, M.S., Wijesundara, S.A., 2004. Community ecology in an everwet forest in Sri Lanka. In: Losos, Elizabeth C., Leigh, Egbert G. (Eds.), *Tropical Forest Diversity and Dynamism: Findings from a Large-scale Network*. University of Chicago Press, Chicago, USA; London, U.K., pp. 119–144.
- Kumar, A., Sinha, A.K., Singh, D., 2003. Studies of eucalyptus plantations under the farm forestry and agroforestry systems of UP in Northern India. *Forest. Trees Livelihoods* 13, 313–330. 10(1080/14728028), 2003, 9752468.
- Kurosaki, T., 2009. Land-use changes and agricultural growth in India, Pakistan, and Bangladesh, 1901–2004. In: Dutta, B., Ray, T., Somanathan, E. (Eds.), *New and Enduring Themes in Development Economics*. World Scientific Publishing, Chennai/Singapore, pp. 303–330.
- Lamb, D., 1998. Large-scale ecological restoration of degraded tropical forest lands: the potential role of timber plantations. *Restor. Ecol.* 6, 271–279.
- Lamb, D., Erskine, P.D., Parrotta, J.A., 2005. Restoration of degraded tropical forest landscapes. *Science* 310, 1628–1632.
- Lewis, S.L., Lopez-Gonzalez, G., Sonké, B., Affum-Baffoe, K., Baker, T.B., Ojo, L.O., Phillips, O.I., Reitsma, J.M., White, L., Comiskey, J.A., Djukouo, M., Ewango, C.E.N., Feldpausch, T.R., Hamilton, A.C., Gloor, M., Hart, T., Hladik, A., Lloyd, J., Lovett, J.C., Makana, J., Malhi, Y., Mbago, F.M., Ndangalasi, H.J., Peacock, J., Peh, K.S.H., Sheil, D., Sunderland, T., Swaine, M.D., Taplin, J., Taylor, D., Thomas, S.C., Votere, R., Wöhl, H., 2009. Increasing carbon storage in intact African tropical forests. *Nature* 457, 1003–1006.
- Mark, S., 2007. Technical and financial analysis of enrichment planting in logging gaps as a potential component of forest management in the eastern Amazon. *Forest Ecol. Manage.* 255, 866–879.
- MEA (Millennium Ecosystem Assessment), 2003. *Ecosystems and human well-being*. Washington, DC, USA, Island Press.
- Menalled, F.D., Kelt, M.J., Ewel, J.J., 1998. Canopy development in tropical tree plantations: a comparison of species mixtures and monocultures. *Forest Ecol. Manage.* 104, 249–263.
- Miyawaki, A., 1999. Creative ecology: restoration of native forests by native trees. *Plant Biotechnol. J.* 16, 15–25.
- Myers, N., 1988. Tropical forests: much more than stocks of wood. *J. Trop. Ecol.* 4, 209–221. <http://dx.doi.org/10.1017/S0266467400002728>.
- Nagaraja, B.C., Maskey, N., Somasekara, R.K., 2011. Regeneration of native woody species under plantations in Kudremukh National Park, Western Ghats of South India. *Int. J. Biodivers. Sci. Ecosys. Serv. Manage.* 7, 77–83.
- Näsström, R., Mattsson, E., 2011. Country report Sri Lanka land-use change and forestry at the national and sub-national level. Focall, Sweden.
- Nath, S., Das, P.K., Gangopadhyay, S.K., Singh, B., Banerjee, S.K., Singh, B., 1990. Suitability of different forest species for social forestry programme under different soil conditions. Part II. Lateritic Soil Indian 116, 464–473.
- Park, A., van Breugel, M., Ashton, M.S., Wishnie, M., Mariscal, E., Deago, J., Ibarra, D., Cedeno, N., Hall, J.S., 2010. Local and regional environmental variation influences the growth of tropical trees in selection trials in the Republic of Panama. *Forest Ecol. Manage.* 260, 12–21.
- Parrotta, J.A., Knowles, O.H., 2002. Restoration of tropical moist forests on bauxite-mined lands in the Brazilian Amazon. *Restor. Ecol.* 7, 103–116.
- Phillips, O.L., Malhi, Y., Higuchi, N., Laurance, W.F., Núñez, P.V., Vázquez, R.M., Laurance, S.G., 1998. Changes in the carbon balance of tropical forests: evidence from long-term plots. *Science* 282, 439–442.
- Pomeroy, M., Primack, R., Rai, S.N., 2003. Changes in four rainforest plots of the Western Ghats, India, 1939–1993. *Conserv. Soc.* 1, 113–135.
- Pringle, C.M., 2001. Hydrologic connectivity and the management of biological reserves: a global perspective. *Ecol. Appl.* 11, 981–998.
- Rai, S.N., 1999. *Nursery and planting techniques of forest trees in tropical South-Asia*. Punarvasu Publications Dharwad. Karantaka – Eastern Press, Bangalore, India.
- Ramachandra, T.V., Kamakshi, G., Shrutthi, B.B., 2004. Bio resource status in Karnataka. *Renew. Sustain. Energy Rev.* 8, 1–47.
- Raman, T.R., Mudappa, D., Kapoor, V., 2009. Restoring rainforest fragments: Survival of mixed-native species seedlings under contrasting site conditions in the Western Ghats. *India Restor. Ecol.* 17, 137–147.
- Rands, M.R., Adams, W.M., Bunnell, L., Butchart, S.H., Clements, A., Coomes, D., Entwistle, A., Hodge, I., Kapos, V., Scharlemann, J.P.W., Sutherland, W.J., Vira, B., 2010. Biodiversity conservation: challenges beyond 2010. *Science* 329, 1298–1303.
- Richards, A.E., Schmidt, S., 2010. Complementary resource use by tree species in a rain forest tree plantation. *Ecol. Appl.* 20, 1237–1254.
- Ricketts, T.H., Daily, G.C., Ehrlich, P.R., Michener, C., 2004. Economic value of tropical forests in coffee production. *Proc. Natl Acad. Sci. USA* 101 (579–12), 582.
- Salek, L., Vylupek, O., 2012. Contribution to the restoration of mixed forests in Central Vietnam. *J. Sust. Forest.* 31, 549–562.
- Saxena, K.G., Rao, K.S., Purohit, A.N., 1993. Sustainable forestry prospects in India. *J. Sustain. Forest.* 1, 69–95. http://dx.doi.org/10.1300/J091v01n02_04.
- Sharam, G., Sinclair, A.R.E., Turkington, R., 2006. Establishment of broad-leaved thickets in Serengeti, Tanzania: the influence of fire, browsers, grass competition, and elephants. *Biotropica* 38, 599–605.
- Slocum, M.G., Aide, T.M., Zimmerman, J.K., Navarro, L., 2006. A strategy for restoration of montane forest in anthropogenic fern thickets in the Dominican Republic. *Restor. Ecol.* 14, 526–536.
- Tennakoon, M.M.D., Gunatilleke, I.A.U.N., Hafeel, K.M., Seneviratne, G., Gunatilleke, C.V.S., Ashton, P.M.S., 2005. Ectomycorrhizal colonization and seedling growth of *Shorea* (Dipterocarpaceae) species in simulated shade environments of a Sri Lankan rain forest. *Forest Ecol. Manage.* 208, 399–405.
- Vincent, F.d'A., 1883. *Forest Administration of Ceylon. Report on the conservation and administration of the Crown forests in Ceylon. With a map. Papers laid before the Legislative Council of Ceylon. 1882–1883*. Sessional Paper 43.
- Vincent, A., Davies, S.J., 2003. Effects of nutrient addition, mulching and planting-hole size on early performance of *Dryobalanops aromatica* and *Shorea parvifolia* planted in secondary forest in Sarawak. *Malaysia Forest Ecol. Manage.* 180, 261–271.
- Walters, B.B., 1997. Human ecological questions for tropical restoration: experiences from planting native upland trees and mangroves in the Philippines. *Forest Ecol. Manage.* 99, 275–290.
- Wickramagamage, P., 1998. Large-scale deforestation for plantation agriculture in the hill country of Sri Lanka and its impacts. *Hydrol. Process.* 12, 2015–2028.
- Wimaladharma, K.P., 1977. Land use adjustments in the Mid-country of Sri Lanka: an examination of the Nilambe Atabage catchment development project. MA Dissertation, University of Colombo, Sri Lanka.
- Wishnie, M.H., Dent, D.H., Mariscal, E., Deago, J., Cedeno, N., Ibarra, D., Condit, R., Ashton, P.M.S., 2007. Initial performance and reforestation potential of 24

- tropical tree species planted across a precipitation gradient in the Republic of Panama. *Forest Ecol. Manage.* 243, 39–49.
- Wong, N.H., Yu, C., 2005. Study of green areas and urban heat island in a tropical city. *Habitat Int.* 29, 547–558.
- Yirdaw, E., 2001. Diversity of naturally-regenerated native woody species in forest plantations in the Ethiopian highlands. *New Forest.* 22, 159–177.