

Genetic Analysis of Osmotic and Cold Stress Signal Transduction in Arabidopsis: Interactions and Convergence of Abscissic Acid-Dependent and Abscissic Acid-Independent Pathways

Manabu Ishitani, Liming Xiong, Becky Stevenson, and Jian-Kang Zhu¹

Department of Plant Sciences, University of Arizona, Tucson, Arizona 85721

To dissect genetically the complex network of osmotic and cold stress signaling, we constructed lines of Arabidopsis plants displaying bioluminescence in response to low temperature, drought, salinity, and the phytohormone abscissic acid (ABA). This was achieved by introducing into Arabidopsis plants a chimeric gene construct consisting of the firefly luciferase coding sequence (*LUC*) under the control of the stress-responsive *RD29A* promoter. *LUC* activity in the transgenic plants, as assessed by using in vivo luminescence imaging, faithfully reports the expression of the endogenous *RD29A* gene. A large number of *cos* (for constitutive expression of osmotically responsive genes), *los* (for low expression of osmotically responsive genes), and *hos* (for high expression of osmotically responsive genes) mutants were identified by using a high-throughput luminescence imaging system. The *los* and *hos* mutants were grouped into 14 classes according to defects in their responses to one or a combination of stress and ABA signals. Based on the classes of mutants recovered, we propose a model for stress signaling in higher plants. Contrary to the current belief that ABA-dependent and ABA-independent stress signaling pathways act in a parallel manner, our data reveal that these pathways cross-talk and converge to activate stress gene expression.

INTRODUCTION

Drought, salinity, and low temperature are common adverse environmental factors encountered by land plants (Boyer, 1982; Thomashow, 1994; Bohnert et al., 1995). Water deficit caused by drought and high salinity has been a major selective force in plant evolution and an important factor limiting crop productivity. On the other hand, low temperature is perhaps the most important environmental constraint for plant distribution on land. To cope with these environmental stresses, plants execute a number of physiological and metabolic responses (Bartels and Nelson, 1994; Thomashow, 1994; Bohnert et al., 1995). Knowledge of the mechanisms by which plants perceive and transduce the stress signals is the key to understanding these responses and to genetic improvement of stress tolerance through biotechnology.

In response to osmotic stress elicited by water deficit or conditions of high salt, the expression of numerous genes is altered in plants (Skriver and Mundy, 1990; Bray, 1993; Bartels and Nelson, 1994; Zhu et al., 1997). Some of the osmotic stress-responsive (OR) genes can also be induced by low-temperature stress (Nordin et al., 1991; Thomashow, 1994; Giraudat, 1995). Both osmotic and cold stresses increase the level of the phytohormone abscissic acid (ABA; Zeevaart and Creelman, 1988; Skriver and Mundy, 1990;

Chandler and Robertson, 1994). The expression of many OR genes can be induced by the application of ABA (Skriver and Mundy, 1990; Bray, 1993; Zhu et al., 1997). Accordingly, ABA is known to mediate OR gene expression in response to osmotic and cold stresses (Skriver and Mundy, 1990; Chandler and Robertson, 1994; Zhu et al., 1997). However, analysis of OR gene expression in ABA-deficient and ABA-insensitive mutants has indicated that the expression of some OR genes can act independently of ABA (Gilmour and Thomashow, 1991; Nordin et al., 1991; Gosti et al., 1995). In addition, a *cis*-acting DNA regulatory element, termed the dehydration-responsive element (DRE)/C-repeat, which responds to cold or osmotic stress but not to ABA, has been found in some OR promoters (Yamaguchi-Shinozaki and Shinozaki, 1994; Stockinger et al., 1997). In contrast, the ABA-responsive element/complex (ABRE) is known to mediate gene expression in response to ABA (Guiltinan et al., 1990; Yamaguchi-Shinozaki and Shinozaki, 1994; Shen and Ho, 1995; Vasil et al., 1995). Therefore, ABA-dependent and ABA-independent signal transduction pathways have been proposed to function in a parallel manner to mediate gene expression in response to cold and osmotic stresses (Yamaguchi-Shinozaki and Shinozaki, 1994; Gosti et al., 1995; Shen and Ho, 1995; Stockinger et al., 1997). However, the results described in this study reveal that ABA-dependent and ABA-independent stress signaling

¹To whom correspondence should be addressed. E-mail jkzhu@ag.arizona.edu; fax 520-621-7186.

pathways do not act in a parallel manner; rather, they interact and converge to activate stress genes.

Despite the rapid progress in dissecting osmotic signaling pathways in the unicellular model eukaryote *Saccharomyces cerevisiae* (Brewster et al., 1993; Maeda et al., 1994; Posas and Saito, 1997) and the wealth of information on the identification and expression of plant OR genes (Zhu et al., 1997), understanding of the osmosensing mechanism in plants still remains a major challenge. Phospholipase C (Hirayama et al., 1995), many putative transcription factors (Urao et al., 1993; Stockinger et al., 1997), and protein kinases (Urao et al., 1994; Nishihama et al., 1995) have been described—some of which are encoded by OR genes—but their role in osmotic signaling is unclear. Mutants defective in osmotic responses would be valuable for dissecting osmotic signaling pathways. Traditional approaches to isolating mutants with altered osmotic responses are problematic because of difficulties in identifying a reliable phenotype for mutant screening. Hence, except for several ABA signaling mutants (e.g., ABA deficient [*aba*], ABA insensitive [*abi*], and enhanced response to ABA [*era*]), which were identified by their aberrant seed germination response to ABA (Koornneef et al., 1982, 1984; Finkelstein, 1994; Cutler et al., 1996; Leon-Kloosterziel et al., 1996), it has not been possible to isolate other plant mutants defective in osmotic signal transduction.

To begin a comprehensive genetic analysis of the osmotic and cold signal transduction pathways in plants, we identified Arabidopsis mutants that show altered regulation of OR gene expression. Arabidopsis plants were transformed with a chimeric gene construct (*RD29A-LUC*) containing a firefly luciferase reporter (Millar et al., 1992, 1995) driven by the DRE/C-repeat- and ABRE-containing *RD29A* promoter (Yamaguchi-Shinozaki and Shinozaki, 1994). The resulting plants emit bioluminescence in response to cold, osmotic stress, or exogenous application of ABA. Seeds from transgenic plants homozygous for the transgene were mutagenized by ethyl methanesulfonate (EMS), and the M_2 seedlings were screened for mutants with altered bioluminescence. Hundreds of such mutants were obtained, and their responses to osmotic stress, cold, and ABA were characterized. Surprisingly, the phenotypes of many of the mutants cannot be explained by current models of osmotic signal transduction. Based on the analysis of these mutants, we propose an alternative scheme of osmotic and cold signal transduction in which ABA-dependent and ABA-independent pathways interact and converge to activate stress genes.

RESULTS

Regulation of Bioluminescence in *RD29A-LUC* Transgenic Arabidopsis Plants by Osmotic Stress, Low Temperature, and ABA

The choice of an appropriate promoter and reporter gene is critical for utilizing a promoter-reporter approach to screen

for gene regulation/signaling mutants. For the selection of osmotic signaling mutants, we chose the *RD29A* promoter (Yamaguchi-Shinozaki and Shinozaki, 1994) and the firefly *LUC* reporter gene (Millar et al., 1992). Although the function of the *RD29A* gene (also known as *cor78* or *lti78*; Horvath et al., 1993; Nordin et al., 1993) product is not known, the *RD29A* promoter is one of the well-characterized promoters that can be activated by osmotic and cold stresses (Yamaguchi-Shinozaki and Shinozaki, 1994). In addition to an ABRE/ABA response complex mediating ABA regulation, the *RD29A* promoter also contains the DRE element, which can be activated by osmotic and cold stresses but not by ABA (Yamaguchi-Shinozaki and Shinozaki, 1994). Thus, this promoter makes it possible to identify mutants in both ABA-dependent and ABA-independent pathways. The *LUC* reporter gene was chosen because its expression in plants can be measured noninvasively by using low-light video imaging, making it practical to screen a large population of plants. Because of the presumed complex nature of osmotic and cold signaling in plants, a high-throughput video-imaging method of screening is necessary to isolate a large number of mutants to recover mutations in many signaling components.

The *RD29A-LUC* construct used to transform Arabidopsis is illustrated in Figure 1A. The construct was introduced into Arabidopsis (ecotype C24) via Agrobacterium-mediated root transformation (Valvekens et al., 1985). From nine independent transformants, one line was chosen for subsequent experiments because it displayed the highest *LUC* activity under osmotic and cold stresses (data not shown). Kanamycin-resistant and kanamycin-sensitive plants segregated 3:1 in the T_1 population, indicating that there is a single functional *RD29A-LUC* transgene in this line. Plants homozygous for the transgene were identified from the progeny of this line. Seeds from these plants were collected and used for analysis of *LUC* expression and mutagenesis.

To test the suitability of the *RD29A-LUC* transgenic plants for isolation of mutants, we first examined the effect of low-temperature treatment on *LUC* expression. One-week-old seedlings grown in agar plates were exposed to cold (0°C) for 48 hr, and their luminescence was measured with a low-light CCD imaging system. As shown in Figure 1B, luminescence was detected in the *RD29A-LUC* plants only after imposing cold stress. In contrast, transgenic plants containing the *PC-LUC* transgene (*LUC* being under the control of the light-responsive plastocyanin promoter; Dijkwel et al., 1996) emitted luminescence without cold stress; after the cold treatment, luminescence in the *PC-LUC* plants decreased (Figure 1B). As expected, control plants without the *RD29A-LUC* or *PC-LUC* transgene did not emit luminescence before or after the cold stress (Figure 1B). Further imaging experiments showed that cold, ABA, or high-salt stress strongly induced bioluminescence in roots, stems, leaves, and cotyledons (data not shown).

The response of *RD29A-LUC* to cold, drought, or ABA is shown in Figure 2A. Cold stress, osmotic stresses (NaCl or




Figure 1. Construction of Transgenic Arabidopsis with Stress-Inducible Bioluminescence.

(A) Diagrammatic representation of the *RD29A-LUC* construct used for transformation. LB and RB, left and right T-DNA borders, respectively; NPT-II, neomycin phosphotransferase II (for kanamycin resistance).

(B) The *RD29A-LUC* plants emit luminescence in response to cold stress. Control, untransformed plants; PC-LUC, plants transformed with *PC-LUC* (i.e., LUC under the control of the light-responsive plastocyanin promoter; Dijkwel et al., 1996); *RD29A-LUC*, plants transformed with *RD29A-LUC*. Left, bright-field illumination of all plants; center, luminescence image of the plants before cold treatment; right, luminescence after cold stress (0°C for 2 days). The color scale on the right shows the luminescence intensity from dark blue (lowest) to white (highest).

polyethylene glycol [PEG]), or ABA strongly induced bioluminescence in the *RD29A-LUC* plants. Under NaCl treatments, the luminescence level became higher as the NaCl concentration increased. The luminescence reached its peak level at 300 mM NaCl (Figure 2B). Higher concentrations of NaCl resulted in decreased levels of luminescence (Figure 2B). Figures 2C and 2D show the time course of *RD29A-LUC* expression in response to 300 mM NaCl or 100 μ M ABA treatment, respectively. The response to both treatments was rapid; significant luminescence was detected 1 hr after treatment, and the expression reached near peak levels by 3 hr after the treatments (Figures 2C and 2D). The patterns of these bioluminescence responses to ABA and various stresses are similar to that of the endogenous *RD29A* gene, as determined by RNA gel blot analysis (Horvath et al., 1993; Yamaguchi-Shinozaki and Shinozaki, 1993; M. Ishitani and J.-K. Zhu, unpublished data), and to responses reported in studies using a promoter- β -glucuronidase gene reporter (Horvath et al., 1993; Yamaguchi-Shinozaki and Shinozaki, 1994). We conclude that *RD29A-LUC* faithfully reflects stress and ABA regulation and appears to be reliable for genetic analysis of stress and ABA responses.

Isolation of Mutants with Altered Bioluminescence with Respect to Osmotic Stress, Low Temperature, and ABA Regulation

To conduct a large-scale screening of stress and ABA signaling mutants, it was necessary to have a large population of mutagenized plants. One gram of the *RD29A-LUC* transgenic seeds was mutagenized by EMS, divided into 28 pools, and germinated, and the resulting plants were allowed to self-pollinate. M_2 seeds from each pool (~ 9 g of seeds representing ~ 1200 M_1 plants) were collected independently. We were interested in three types of mutants designated as *cos* (for constitutive expression of osmotically responsive genes), *los* (for low expression of osmotically responsive genes), and *hos* (for high expression of osmotically responsive genes), respectively.

Figure 3 outlines the screening strategy that was used to select for the *cos*, *los*, and *hos* mutants. Agar plates, each containing 500 to 1000 1-week-old seedlings (Figure 4A), were first imaged for luminescence before stress treatment (Figure 4C). Individuals emitting significant luminescence were marked as putative *cos* mutants. The plates were then




Figure 2. Characterization of the Regulation of Luminescence from *RD29A-LUC* Plants by Cold, Osmotic Stress, or ABA.

(A) Induction of luminescence by NaCl (300 mM for 5 hr), PEG (30% for 5 hr), ABA (100 μ M for 3 hr), or cold stress (0°C for 2 days).

(B) Concentration curve of NaCl response (5-hr treatments).

(C) Time course of NaCl (300 mM) response.

(D) Time course of ABA (100 μ M) response.

Each point represents the average luminescence from 48 seedlings. Error bars represent standard deviation.

placed at 0°C in the dark for 48 hr followed by a second luminescence imaging. Seedlings with abnormally low or high luminescence levels were noted as putative *los* and *hos* mutants, respectively (Figure 4D). The plates were then placed at room temperature (20 to 22°C) for 2 days to allow the luminescence to drop to pre-cold treatment levels. The plants were then sprayed with 100 μ M ABA, and a third luminescence image was taken 3 hr later. Individuals with abnormally low or high luminescence levels were again marked as putative *los* and *hos* mutants, respectively. All putative mutants were then transferred from the plates to grow in soil. Some of the remaining seedlings were transferred onto filter papers soaked with 300 mM NaCl to identify additional putative *los* and *hos* mutants.

During the screening of *cos* mutants, it was difficult to

match accurately the luminescence image of a putative *cos* mutant directly with the individual seedling in a plate containing several hundred plants. To solve this problem, we took advantage of the natural fluorescence emitted by the seedlings. Green plants exposed to light emit fluorescence during the first 1 or 2 min after being transferred to the dark. Before luminescence imaging for *cos* mutants, a fluorescence image of all seedlings can be collected with a 30-sec exposure (Figure 4B). To identify precisely a putative *cos* mutant out of hundreds of seedlings in an agar plate, we simply looked for a seedling in the fluorescence image with the same coordinates as the one in the luminescence image. An example of the use of fluorescence and luminescence imaging before and after cold treatment is shown in Figure 4. Normally, to avoid the interference of fluorescence during

luminescence imaging, we spray seedlings with luciferin first and then store the plants in the dark for 5 min before collecting a luminescence image.

Approximately 300,000 M_2 seedlings from the 28 pools were screened, and 3000 were selected as putative mutants for growth in soil. Of these, 1768 lines survived and set seeds, and their progeny were rescreened to eliminate false positives (Figure 3). Approximately 20 seedlings from each putative mutant were examined for luminescence expression before and after cold and ABA treatments. Five to 10 seedlings from each putative mutant were also transferred from the agar plates onto a filter paper soaked with 300 mM NaCl and incubated for 5 hr. Luminescence images were then taken from the NaCl-treated seedlings. Of these, 833 were confirmed as true *cos*, *los*, or *hos* mutants during the rescreening process; 103 of them exhibited the strongest luminescence phenotypes and were further characterized.

Characterization of Stress and ABA Responses in the Mutants

To categorize the mutants, we quantitated their responses to cold, NaCl, and ABA. The 103 mutants with the strongest phenotypes fall into 13 categories: *cos* (Figure 5A); hyperresponsive to osmotic stress, cold, and ABA (*hos_{all}*; Figure 5B); hyperresponsive to cold only (*hos_{cold}*; Figure 5C); hyperresponsive to osmotic stress only (*hos_{NaCl}*; Figure 5D); hyperresponsive to ABA only (*hos_{ABA}*; Figure 5E); hyperresponsive to osmotic stress and cold (*hos_{cold/NaCl}*; Figure 5F); hyperresponsive to cold and ABA (*hos_{cold/ABA}*; Figure 5G); hyperresponsive to osmotic stress and ABA (*hos_{NaCl/ABA}*; Figure 5H); low response to osmotic stress, cold, and ABA (*los_{all}*; Figure 6A); low response to cold only (*los_{cold}*; Figure 6B); low response to osmotic stress only (*los_{NaCl}*; Figure 6C); low response to osmotic stress and cold (*los_{cold/NaCl}*; Figure




Figure 3. The Protocol Used for the Screening for Mutants.




Figure 4. An Example of Screening *cos*, *los*, and *hos* Mutants for Cold Stress.

(A) Plate containing 6-day-old M_2 seedlings.

(B) Fluorescence image of the seedlings.

(C) Luminescence image taken before stress treatments.

(D) Luminescence image taken after cold stress treatment.

Putative *cos*, *los*, and *hos* mutants are marked. The fluorescence image was taken to assist in the accurate identification of the putative *cos* plant. For relative luminescence intensities, refer to the color scale in Figure 1.

6D); and low response to osmotic stress and ABA ($los_{NaCl/ABA}$; Figure 6E). Table 1 lists the number of lines in all mutant classes obtained. Figure 7 shows the luminescence images of some of the *cos*, *los*, and *hos* mutants under various treatments.

The *cos* mutant shown in Figures 5A and 7 emits lumines-

cence constitutively. However, the responses to cold, ABA, and NaCl are not changed in this mutant. This is not true in all *cos* mutants. One *cos* mutant (line 396) showed reduced responses to cold, NaCl, and ABA, whereas others had enhanced responses to these treatments (data not shown). We also recovered *cos* mutants with enhanced responses to

just one or two of the treatments (M. Ishitani and J.-K. Zhu, unpublished data). The majority of the *cos* mutants that we obtained exhibited slow growth and appeared to be “constitutively stressed.”

The two classes that include the largest numbers of mutant lines are *los_{all}* and *hos_{all}* (Table 1). In several of the *los_{all}* mutants, the responses to cold, NaCl, or ABA were nearly eliminated. Most other *los_{all}* mutants still responded to the




Figure 5. Luminescence of Representative *cos* and *hos* Mutants in Response to Cold, High-Salt, and ABA Treatments.

M₄ plants were grown for 10 days and then subjected to control (untreated), cold (0°C for 2 days), ABA (100 μ M for 3 hr), or NaCl (300 mM for 5 hr) treatments. At the end of the treatments, luminescence was determined, and the mean level was calculated from data for 10 to 20 single seedlings. White bar, wild type; black bar, mutant. Error bars represent standard deviation.

- (A) A *cos* mutant with constitutive expression of luminescence.
- (B) A *hos* mutant with enhanced responses to cold, ABA, and high salt.
- (C) A *hos* mutant with a high response to only low-temperature stress.
- (D) A *hos* mutant with a high response to only osmotic stress.
- (E) A *hos* mutant with an enhanced response to only ABA.
- (F) A *hos* mutant with enhanced responses to cold and osmotic stresses.
- (G) A *hos* mutant with enhanced responses to cold and ABA.
- (H) A *hos* mutant with enhanced responses to ABA and osmotic stress.




Figure 6. Luminescence of Representative *los* Mutants in Response to Cold, High-Salt, and ABA Treatments.

Treatment and assay conditions were the same as described for Figure 5. White bar, wild type; black bar, mutant. Error bars represent standard deviation.

(A) A *los* mutant with reduced responses to cold, ABA, and osmotic stress.

(B) A *los* mutant with a reduced response to only cold stress.

(C) A *los* mutant with a reduced response to only osmotic stress.

(D) A *los* mutant with reduced responses to cold and osmotic stresses.

(E) A *los* mutant with reduced responses to ABA and osmotic stress.

stress and ABA treatments, but their responses were reduced. Nearly all *hos_{all}* mutants also exhibited substantial luminescence in the absence of a stress treatment (Figures 5B and 6; M. Ishitani and J.-K. Zhu, unpublished data). However, the constitutive expression in *hos_{all}* mutants generally was not as high as those in the *cos* class of mutants.

Expression of the endogenous *RD29A* gene was examined with RNA gel blot analysis in the 103 mutants with the

strongest luminescence phenotypes. It is easy to envision that a mutation in the *LUC* coding region or even in the *RD29A* promoter could result in a *los* phenotype. These *cis* mutants would not be of interest. Figure 8 shows the expression of the endogenous *RD29A* gene in *los10*, *los1*, and *los5* mutants, which represent the *los_{all}*, *los_{cold}*, and *los_{cold/NaCl}* classes, respectively. Complementation tests showed that these mutations are not allelic to each other (data not

shown). Whereas the *RD29A* transcript was induced by cold, ABA, NaCl, or PEG in the wild-type parent, this induction was greatly reduced in the *los10* mutant (Figure 8). The *los1* mutation specifically blocked *RD29A* induction by cold, but induction by ABA, NaCl, or PEG was not affected (Figure 8). In *los5* mutant plants, *RD29A* induction by NaCl or PEG was eliminated, and induction by cold was reduced, whereas ABA regulation was normal (Figure 8). As a control, expression of the actin gene was not changed in these mutants (Figure 8). These results confirmed the phenotypes of the mutants as revealed by their luminescence responses. In fact, in at least one mutant from each of the 13 categories presented in Figures 5 and 6, the steady state *RD29A* mRNA level was found to correlate with *LUC* reporter activity (data not shown), indicating that the luminescence phenotypes in these mutants are caused by mutations in signaling components and not in the *RD29A-LUC* construct.

DISCUSSION

In this study, we present results on the isolation of hundreds of Arabidopsis mutants defective in the regulation of *RD29A* gene expression by cold, osmotic stress, or ABA. In addition to many expected mutants, several unexpected classes of mutants were recovered. These mutants provide a foundation for comprehensive analysis of stress and ABA signal transduction. Categorization of the mutants provides a global view of cold, osmotic stress, and ABA signaling.

A High-Throughput Method for the Selection of Stress and ABA Signal Transduction Mutants

Because of the complexity of the responses of plants to environmental stresses, it has not been possible to identify osmotic and cold stress signal transduction mutants by their morphological or physiological phenotypes. Osmotic signaling mutants of yeast were identified by NaCl inhibition of growth (Brewster et al., 1993; Maeda et al., 1994). Our group has isolated many *sos* (salt overly sensitive) mutants of Arabidopsis (Wu et al., 1996; J.-K. Zhu, unpublished data). However, these mutants are all specifically hypersensitive to Na⁺ concentrations and not to general osmotic stress. Arabidopsis is a glycophyte that is very sensitive to NaCl (Wu et al., 1996), with concentrations >150 mM resulting in near complete inhibition of growth. In contrast, *S. cerevisiae* can tolerate almost 1 M NaCl (Haro et al., 1993). Therefore, the very salt-sensitive nature of Arabidopsis makes it impossible to isolate osmotic response mutants by screening for NaCl-hypersensitive growth.

We have developed a very efficient method for the isolation of osmotic as well as cold and ABA signaling mutants by using transgenic plants with stress-inducible bioluminescence. This method takes advantage of low-light video im-

Table 1. Numbers of Mutant Lines in Various Mutant Classes

Mutant Classes	1 ^a	2 ^b	3 ^c
<i>cos</i>	9	16	8
<i>los</i>			
All	34	139	198
Cold	3	7	23
NaCl	3	15	47
ABA	0	6	18
Cold/NaCl	3	2	17
NaCl/ABA	2	4	12
Cold/ABA	0	6	9
<i>hos</i>			
All	21	12	10
Cold	9	9	32
NaCl	1	20	32
ABA	3	11	16
Cold/NaCl	1	9	12
NaCl/ABA	11	10	11
Cold/ABA	3	5	14

^a Mutants with strong luminescence phenotypes (total of 103).

^b Mutants with intermediate phenotypes (total of 271).

^c Mutants with weak phenotypes (total of 459).

aging and thus makes it easy to screen large populations of plants. Gene expression in hundreds of plants can be quantitatively detected by luminescence imaging in a matter of several minutes. A Petri dish containing several hundred to 1000 seedlings can be subjected to multiple stress and hormonal treatments and imaged repeatedly. Our screening procedure (Figure 4) is comparable to replica plating in prokaryotic genetics.

Using this procedure, we were able to screen efficiently >300,000 M₂ plants. The limitation in screening more plants is in time spent growing Arabidopsis in agar plates. More than 100 mutants exhibiting strong phenotypes were readily recovered. These mutants include not only constitutive expressers but also low and high expressers in response to cold, high salt, or ABA. Some of the mutants may turn out to be caused by mutations in the transgene and are not of interest. However, many have been found to be *trans* mutants and should be useful for studies on stress and ABA signal transduction. The frequency of these mutations may appear very high. However, considering that there are >10 categories of these mutants, the frequency for any particular mutant class is not too high. A high frequency of these mutations may also reflect the complexity of osmotic and cold responses in higher plants.

Interactions and Convergence of ABA-Dependent and ABA-Independent Stress Signaling Pathways

Recovery of those mutants defective specifically in their responses to cold, osmotic stress, or ABA is expected. Because




Figure 7. Images of Luminescence from Selected Examples of *hos* and *los* Mutants in Response to Cold, ABA, and Osmotic Stress Treatments.

(A) Arrangement of the wild type (WT) and mutant lines. Clockwise: wild type, the *RD29A-LUC* transgenic parent; *cos*, a mutant with constitutive expression of *RD29A* (Figure 5A); *los* to all, a *los* mutant with reduced responses to cold, ABA, and osmotic stress (Figure 6A); *hos* to cold, a *hos* mutant with an enhanced response to cold (Figure 5C); *los* to cold & NaCl, a *los* mutant with reduced responses to cold and osmotic stress; *hos* to NaCl & ABA, a *hos* mutant with an enhanced response to ABA and osmotic stress (Figure 5H); *los* to cold, a *los* mutant with a reduced response to cold stress (Figure 6B); *hos* to all, a *hos* mutant with an enhanced response to cold, ABA, and osmotic stress (Figure 5H).

(B) Morphology of the seedlings.

(C) Image of luminescence from the plants before stress and ABA treatments. Note that the *hos* mutants had constitutive luminescence.

(D) Image of luminescence from the plants that had been cold treated for 2 days.

(E) Image of luminescence from the plants after ABA treatment. The plants were placed at room temperature (20 to 22°C) for 2 days to allow the luminescence to drop to pre-cold treatment levels before being sprayed with 100 μ M ABA and then imaged 3 hr later.

(F) Image of luminescence from the plants under osmotic stress. Three days after the ABA treatment, luminescence from the plants dropped to pretreatment levels. The plants were then treated with high salt for 5 hr by flooding the plate with 300 mM NaCl. Some of the *cos* mutant seedlings died after the cold and ABA treatments and thus did not emit luminescence upon NaCl treatment.

For relative luminescence intensities, refer to the color scale in Figure 1.

nearly all of the mutations were recessive (data not shown), the *los_{cold}* and *hos_{cold}* mutants most likely define positive and negative regulators, respectively, in cold signal transduction. Similarly, the *los_{NaCl}* and *hos_{NaCl}* mutants most likely define positive and negative regulators, respectively, in osmotic signaling.

However, the phenotypes of several other categories of mutants are not expected and cannot be explained by current knowledge of signal transduction in response to abiotic stresses. Existing schemes of osmotic and cold signaling in-

voke parallel ABA-dependent and ABA-independent pathways (Baker et al., 1994; Yamaguchi-Shinozaki and Shinozaki, 1994; Gosti et al., 1995). However, mutations that abolish (Figure 6A) or reduce responses to cold, osmotic stress, and ABA simultaneously suggest that ABA-dependent and ABA-independent pathways converge to activate *RD29A* and other OR gene expression. Similarly, mutations resulting in an enhanced response to osmotic stress, cold, and ABA simultaneously also reveal convergence of the two pathways. Although a mutation causing enhanced response to ABA could

theoretically also result in enhanced response to osmotic stress and cold, data shown in Figure 5E indicate otherwise. Mechanistically, it is not obvious how ABA-dependent and ABA-independent pathways converge, because they target independent *cis*-regulatory elements in the *RD29A* promoter. One possibility is that the convergence points represent transcriptional enhancers and repressors that interact directly or indirectly with both the ABRE and DRE/C-repeat DNA elements. In yeast, for example, a large number of different stimuli controlling mating, haploid invasive growth, or pseudohyphal development were found to converge at the transcriptional factor Ste12 (Madhani and Fink, 1997). This common factor is able to direct selective responses to a stimulus through interactions with more specific transcriptional factors (Madhani and Fink, 1997).

Our results also reveal specific interactions between cold and ABA pathways as well as between osmotic and ABA pathways. Mutations causing enhanced response to osmotic stress and ABA (Figure 5H) define common steps in osmotic and ABA signaling pathways. Interactions between osmotic and ABA pathways are also revealed by mutations that cause low response to osmotic stress and ABA but not to cold (Figure 6E). Mutations causing a hyperresponse to cold and ABA (Figure 5G) reveal a step(s) at which cold and ABA signaling pathways interact. Although it is conceivable that different mutations in one gene could result in phenotypes found in more than one category of the above mutants, allelism tests thus far have shown that mutants in different categories are complements (data not shown).

A Genetic Model of Cold and Osmotic Signal Transduction in Plants

Mutations that fail to synthesize ABA or cause ABA insensitivity or hypersensitivity have been previously isolated by using seed germination assays (Koornneef et al., 1982, 1984; Meyer et al., 1994; Cutler et al., 1996; Leon-Kloosterziel et al., 1996; Leung et al., 1997). These mutations proved instrumental in our understanding of ABA function and signaling. Our work adds many novel types of mutations to the

study of ABA and stress signaling. This new collection of mutants provides the foundation for comprehensive analysis of osmotic, low-temperature, and ABA signal transduction in plants. This comprehensive genetic analysis should complement and facilitate integration of current molecular and biochemical studies of stress signal transduction (Nishihama et al., 1995; Jonak et al., 1996; Sheen, 1996). A scheme that places the various mutations in putative positions in stress signaling pathways is presented (Figure 9). In line with existing experimental evidence (Baker et al., 1994; Yamaguchi-Shinozaki and Shinozaki, 1994; Gosti et al., 1995), the scheme retains the notion of ABA-dependent and ABA-independent cold and osmotic signal transduction. The fact that mutations affecting osmotic stress and cold but not ABA signaling were recovered in the screening (Figure 6D) strongly supports this relationship.

A distinct feature of the proposed scheme is that ABA-dependent and ABA-independent pathways for osmotic stress and low temperature interact and converge (Figure 9). The interactions and convergence may provide added levels of coordination between the stress signals and ABA in the regulation of OR gene expression. It is currently not known whether the points of interactions (i.e., $HOS_{cold/ABA}$, $LOS_{NaCl/ABA}$, and $HOS_{NaCl/ABA}$) are downstream or upstream of the *ABI1* and *ABI2* gene products (Leung et al., 1994, 1997; Meyer et al., 1994). However, because the *abi1* mutation affects seed germination, stomata opening, and OR gene expression, *ABI1* most likely functions early in the ABA pathway (Koornneef et al., 1984; Leung et al., 1994) (Figure 9). Although the cold- or osmotic-specific components are placed upstream of the respective interaction points, their relationships are also unclear (Figure 9).

We hypothesize that *cos* mutations may occur either at early steps of stress signal perception or at the end points of the signal transmission (e.g., transcriptional repressors). A mutation in a very early component of osmotic signaling is expected to result in pleiotropic phenotypes. It is possible that some of the *cos* mutants may accumulate ABA constitutively. We have observed that many of the *cos* mutants are seriously retarded in growth and that their leaves appear vitreous in culture.




Figure 8. RNA Gel Blot Analysis of *RD29A* Gene Expression in Three *los* Mutants.

Twenty micrograms of total RNA from the wild type (WT) or individual mutants was loaded per lane. C, untreated; cold, ABA, NaCl, and PEG treatments are as described in the legend to Figure 2.




Figure 9. A Schematic Representation of Cold, Osmotic Stress, and ABA Signal Transduction.

Gene products defined by the various signaling mutants are placed in putative positions in the pathways. The scheme indicates that ABA-dependent and ABA-independent pathways interact and eventually converge to activate the transcription of *RD29A* and other OR genes. Dashed lines depict pathways that, by themselves, are not sufficient to activate OR gene transcription.

Osmotic stress is the single most limiting environmental factor for crop yield worldwide (Boyer, 1982; Bartels and Nelson, 1994). Unlike that of the unicellular yeast (Brewster et al., 1993; Haro et al., 1993), plant responses to osmotic stress have largely been unamenable to genetic analysis. Our results establish a genetic framework for the elucidation of osmotic and cold stress sensing and intracellular signaling mechanisms and should facilitate the improvement of osmotic stress tolerance of crop plants.

METHODS

Plant Materials and Growth Conditions

In all of the experiments described here, the wild type refers to the unmutagenized *RD29A*-luciferase (*LUC*) line in the *Arabidopsis thaliana* ecotype C24 background. Growth conditions were as described by Wu et al. (1996). During mutant screening and subsequent

experiments with the mutants, we routinely planted seeds in nutrient agar media that were supplemented with 30 μ g/mL kanamycin.

Construction of *RD29A-LUC* Transgenic Plants

The *RD29A* promoter (Yamaguchi-Shinozaki and Shinozaki, 1994) was obtained by polymerase chain reaction by using *Arabidopsis* genomic DNA and two primers: 5'-TCGGGATCCGGTGAATTAAGAGGAGAGAGGAGG-3' and 5'-GACAAGCTTTGAGTAAACAGAGGAGGGTCTCAC-3'. The promoter fragment was inserted into a plant transformation vector containing the firefly *LUC* coding sequence (Millar et al., 1992). *Arabidopsis* plants were transformed with the chimeric *RD29A-LUC* construct via *Agrobacterium tumefaciens* infection of the roots (Valvekens et al., 1988). Plants homozygous for the *RD29A-LUC* chimeric gene were selected from T_1 seeds and used for subsequent experiments.

Mutagenesis and Mutant Isolation

RD29A-LUC seeds were mutagenized with ethyl methanesulfonate (EMS) (Redei and Koncz, 1992). M_2 seeds were sterilized and planted

individually in 150 × 15 mm plates (500 to 1000 seeds per plate) containing Murashige and Skoog salts (Murashige and Skoog, 1962), 3% sucrose, and 0.8% agar, pH 5.7. Five- to 7-day-old seedlings grown under light were sprayed with luciferin and placed immediately under a CCD camera (see next section on LUC imaging) for fluorescence imaging. Fluorescence images were collected with 30-sec exposures. Four minutes after the fluorescence imaging, a LUC image was acquired with a 5-min exposure to identify *cos* mutants. The plates of seedlings were then incubated at 0°C for 2 days. After the cold treatment, the plates were sprayed immediately with luciferin and then incubated at room temperature for 5 min in the dark. A luminescence image was then collected to identify *los* and *hos* mutants. The plates were returned to grow under light for ~2 days before being sprayed with 100 μ M abscisic acid (ABA).

Three hours after the ABA treatment, the seedlings were sprayed with luciferin, incubated in the dark for 5 min, and placed under the CCD camera (model CCD-512SB; Princeton Instruments, Inc., Trenton, NJ) for luminescence collection. Putative *cos* mutants, *los_{cold}* and *hos_{cold}* mutants, and *los_{ABA}* and *hos_{ABA}* mutants were transferred to grow in soil. For a small portion (15 plates) of the M₂ population screened, the remaining plants in the plates were treated with NaCl by transferring individual seedlings onto filter papers soaked with 300 mM NaCl in a Murashige and Skoog solution. To collect luminescence, the NaCl solution was drained before spraying the seedlings with luciferin. Putative *los_{NaCl}* and *hos_{NaCl}* mutants were also transferred to grow in soil. To eliminate false positives, putative mutants were rescreened as described in Results.

Stress and ABA Treatments

For cold treatment, seedlings in agar plates were incubated at 0 ± 2°C in the dark or with dim light. Osmotic stress was imposed by NaCl or polyethylene glycol (PEG) (average molecular mass 6000) treatment. Seedlings were pulled out of agar media and laid down on filter papers soaked with appropriate concentrations of NaCl or PEG solutions for specified time periods. For ABA treatment, 100 μ M ABA in water was sprayed on the seedlings until solution runoff. Osmotic stress and ABA treatments were conducted under light.

LUC Imaging

Imaging with the firefly LUC reporter requires application of the exogenous substrate luciferin. Luciferin (Promega, Madison, WI) was dissolved in sterile water and stored frozen in small aliquots as 100 mM stock solution. One millimolar concentration of working solution of luciferin in 0.01% Triton X-100 was applied uniformly onto seedlings by spraying five times. For LUC imaging, the seedlings were kept for 5 min in the dark after the luciferin application. The imaging system consists of a high-performance CCD camera mounted in a dark chamber, a camera controller, and a computer. Image acquisition and processing were performed with the WinView software provided by the camera manufacturer. Exposure time was 5 min, unless stated otherwise.

RNA Gel Blot Analysis

Total RNA was extracted from seedlings and analyzed as described previously (Liu and Zhu, 1997). The *RD29A* gene-specific probe was from the 3' noncoding region (Liu and Zhu, 1997).

ACKNOWLEDGMENTS

We thank Shaw-Jye Wu and Lei Ding for technical assistance and Drs. Paul P. Dijkwel and Sjef C.M. Smeekeens of Utrecht University for the *PC-LUC* transgenic seeds. We also thank Dr. Robert Leonard for critical reading of the manuscript. This study was supported by grants from the Southwest Consortium on Plant Genetics and Water Resources and the United States Department of Agriculture National Research Initiative Competitive Grants Program (Plant Responses to the Environment) to J.-K.Z.

Received July 10, 1997; accepted September 2, 1997.

REFERENCES

- Baker, S.S., Wilhelm, K.S., and Thomashow, M.F. (1994). The 5'-region of *Arabidopsis thaliana cor 15a* has cis-acting elements that confer cold-, drought- and ABA-regulated gene expression. *Plant Mol. Biol.* **24**, 701–713.
- Bartels, D., and Nelson, D.E. (1994). Approaches to improve stress tolerance using molecular genetics. *Plant Cell Environ.* **17**, 659–667.
- Bohnert, H.J., Nelson, D.E., and Jensen, R.G. (1995). Adaptation to environmental stresses. *Plant Cell* **7**, 1099–1111.
- Boyer, J.S. (1982). Plant productivity and environment. *Science* **218**, 443–448.
- Bray, E.A. (1993). Molecular responses to water deficit. *Plant Physiol.* **103**, 1035–1040.
- Brewster, J.L., de Valoir, T., Dwyer, N.D., Winter, E., and Gustin, M.C. (1993). An osmosensing signal transduction pathway in yeast. *Science* **259**, 1760–1763.
- Chandler, P.M., and Robertson, M. (1994). Gene expression regulated by abscisic acid and its relation to stress tolerance. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **45**, 113–141.
- Cutler, S., Ghassemian, M., Bonetta, D., Cooney, S., and McCourt, P. (1996). A protein farnesyl transferase involved in abscisic acid signal transduction in *Arabidopsis*. *Science* **273**, 1239–1241.
- Dijkwel, P.P., Kock, P.A.M., Bezemer, R., Weisbeek, P.J., and Smeekeens, S.C.M. (1996). Sucrose represses the developmentally controlled transient activation of the plastocyanin gene in *Arabidopsis thaliana* seedlings. *Plant Physiol.* **110**, 455–463.
- Finkelstein, R.R. (1994). Mutations at two new *Arabidopsis* ABA response loci are similar to the *abi3* mutations. *Plant J.* **5**, 765–771.
- Gilmour, S.J., and Thomashow, M.F. (1991). Cold acclimation and cold-regulated gene expression in ABA mutants of *Arabidopsis thaliana*. *Plant Mol. Biol.* **17**, 1233–1240.
- Giraudat, J. (1995). Abscisic acid signaling. *Curr. Opin. Cell Biol.* **7**, 232–238.
- Gosti, F., Bertauche, N., Vartanian, N., and Giraudat, J. (1995). Abscisic acid-dependent and -independent regulation of gene expression by progressive drought in *Arabidopsis thaliana*. *Mol. Gen. Genet.* **246**, 10–18.

- Guiltinan, M.J.W., Marcotte, R., and Quatrano, R.S. (1990). A plant leucine zipper protein that recognizes an abscisic acid response element. *Science* **250**, 267–271.
- Haro, R., Banuelos, M.A., Quintero, F.J., Rubio, F., and Rodríguez-Navarro, A. (1993). Genetic basis of sodium exclusion and sodium tolerance in yeast. A model for plants. *Physiol. Plant.* **89**, 868–874.
- Hirayama, T., Ohto, C., Mizoguchi, T., and Shinozaki, K. (1995). A gene encoding a phosphatidylinositol-specific phospholipase C is induced by dehydration and salt stress in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. USA* **92**, 3903–3907.
- Horvath, D.P., McIarney, B.K., and Thomashow, M.F. (1993). Regulation of *Arabidopsis thaliana* L. (Heynh.) *cor78* in response to low temperature. *Plant Physiol.* **103**, 1047–1053.
- Jonak, C., Kiegerl, S., Ligterink, W., Barker, P., Huskisson, N.S., and Hirt, H. (1996). Stress signaling in plants: A mitogen-activated protein kinase pathway is activated by cold and drought. *Proc. Natl. Acad. Sci. USA* **93**, 11274–11279.
- Koornneef, M., Jorna, M.L., Brinkhorst-Van der Swan, D.L.C., and Karssen, C.M. (1982). The isolation of abscisic acid (ABA) deficient mutants by selection of induced revertants in non-germinating gibberellin sensitive mutants of *Arabidopsis thaliana* (L.) Heynh. *Theor. Appl. Genet.* **61**, 385–393.
- Koornneef, M., Reuling, G., and Karssen, C.M. (1984). The isolation and characterization of abscisic acid-insensitive mutants of *Arabidopsis thaliana*. *Physiol. Plant.* **61**, 377–383.
- Leon-Kloosterziel, K.M., Alvarez Gil, M., Ruijs, G.J., Jacobsen, S.E., Olszewski, N.E., Schwartz, S.H., Zeevaert, J.A.D., and Koornneef, M. (1996). Isolation and characterization of abscisic acid-deficient *Arabidopsis* mutants at two new loci. *Plant J.* **10**, 655–661.
- Leung, J., Bouvier-Durand, M., Morris, P.-C., Guerrier, D., Chefdor, F., and Giraudat, J. (1994). *Arabidopsis* ABA-response gene *ABI1*: Features of a calcium-modulated protein phosphatase. *Science* **264**, 1448–1452.
- Leung, J., Merlot, S., and Giraudat, J. (1997). The *Arabidopsis* *ABSCISIC ACID-SENSITIVE2* (*ABI2*) and *ABI1* genes encode homologous protein phosphatases 2C involved in abscisic acid signal transduction. *Plant Cell* **9**, 759–771.
- Liu, J., and Zhu, J.-K. (1997). Proline accumulation and salt stress-induced gene expression in a salt-hypersensitive *Arabidopsis* mutant. *Plant Physiol.* **114**, 591–596.
- Madhani, H.D., and Fink, G.R. (1997). Combinatorial control required for the specificity of yeast MARK signaling. *Science* **275**, 1314–1317.
- Maeda, T., Wurgler-Murphy, S.M., and Saito, H. (1994). A two-component system that regulates an osmosensing MAP kinase cascade in yeast. *Nature* **396**, 242–245.
- Meyer, K., Leube, M.P., and Grill, E. (1994). A protein phosphatase 2C involved in ABA signal transduction in *Arabidopsis thaliana*. *Science* **264**, 1452–1455.
- Millar, A.J., Short, S.R., Chua, N.-H., and Kay, S.A. (1992). A novel circadian phenotype based on firefly luciferase expression in transgenic plants. *Plant Cell* **4**, 1075–1089.
- Millar, A.J., Carre, I.A., Strayer, C.A., Chua, N.-H., and Kay, S.A. (1995). Circadian clock mutants in *Arabidopsis* identified by luciferase imaging. *Science* **267**, 1161–1163.
- Murashige, T., and Skoog, F. (1962). A revised medium for rapid growth and bioassays with tobacco tissue culture. *Physiol. Plant.* **15**, 473–497.
- Nishihama, R., Banno, H., Shibata, W., Hirano, K., Nakashima, M., Usami, S., and Machida, Y. (1995). Plant homologues of pathways in yeast and animals. *Plant Cell Physiol.* **36**, 749–757.
- Nordin, K., Heino, P., and Palva, E.T. (1991). Separate signal pathways regulate the expression of a low-temperature-induced gene in *Arabidopsis thaliana* (L.) Heynh. *Plant Mol. Biol.* **17**, 1233–1240.
- Nordin, K., Vahala, T., and Palva, E.T. (1993). Differential expression of two related, low-temperature-induced genes in *Arabidopsis thaliana* (L.) Heynh. *Plant Mol. Biol.* **21**, 641–653.
- Posas, F., and Saito, H. (1997). Osmotic activation of the HOG MAPK pathway via Ste11p MAPKKK: Scaffold role of Pbs2p MAPKK. *Science* **276**, 1702–1705.
- Redei, G.P., and Koncz, C. (1992). Classical mutagenesis. In *Methods in Arabidopsis Research*, C. Koncz, N.-H. Chua, and J. Schell, eds (Singapore: World Scientific Publishing Co.), pp. 16–82.
- Sheen, J. (1996). Ca^{2+} -dependent protein kinases and stress signal transduction in plants. *Science* **274**, 1900–1902.
- Shen, Q., and Ho, T.-H.D. (1995). Functional dissection of an abscisic acid (ABA)-inducible gene reveals two independent ABA-responsive complexes each containing a G-box and a novel *cis*-acting element. *Plant Cell* **7**, 295–307.
- Skriver, K., and Mundy, J. (1990). Gene expression in response to abscisic acid and osmotic stress. *Plant Cell* **2**, 503–512.
- Stockinger, E.J., Gilmour, S.J., and Thomashow, M.F. (1997). *Arabidopsis thaliana* *CBF1* encodes an AP2 domain-containing transcriptional activator that binds to the C-repeat/DRE, a *cis*-acting DNA regulatory element that stimulates transcription in response to low temperature and water deficit. *Proc. Natl. Acad. Sci. USA* **94**, 1035–1040.
- Thomashow, M.F. (1994). *Arabidopsis thaliana* as a model for studying mechanisms of plant cold tolerance. In *Arabidopsis*, C. Somerville and E.M. Meyerowitz, eds (Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press), pp. 807–834.
- Urao, T., Yamaguchi-Shinozaki, K., Urao, S., and Shinozaki, K. (1993). An *Arabidopsis* *myb* homolog is induced by dehydration stress and its gene product binds to the conserved MYB recognition sequence. *Plant Cell* **5**, 1529–1539.
- Urao, T., Katagiri, T., Mizoguchi, T., Yamaguchi-Shinozaki, K., Hayashiba, N., and Shinozaki, K. (1994). Two genes that encode Ca^{2+} -dependent protein kinases are induced by drought and high-salt stresses in *Arabidopsis thaliana*. *Mol. Gen. Genet.* **244**, 331–340.
- Valvekens, D., Van Montagu, M., and Van Lijsebettens, M. (1988). *Agrobacterium tumefaciens*-mediated transformation of *Arabidopsis thaliana* root explants by using kanamycin selection. *Proc. Natl. Acad. Sci. USA* **85**, 5536–5540.

- Vasil, V., Marcotte, W.R., Jr., Rosenkrans, L., Cocciolone, S.M., Vasil, I.K., Quatrano, R.S., and McCarty, D.R.** (1995). Overlap of Viviparous1 (VP1) and abscisic acid response elements in the *Em* promoter: G-box elements are sufficient but not necessary for VP1 transactivation. *Plant Cell* **7**, 1511–1518.
- Wu, S.-J., Ding, L., and Zhu, J.-K.** (1996). *SOS1*, a genetic locus essential for salt tolerance and potassium acquisition. *Plant Cell* **8**, 617–627.
- Yamaguchi-Shinozaki, K., and Shinozaki, K.** (1993). Characterization of the expression of a desiccation-responsive *rd29* gene of *Arabidopsis thaliana* and analysis of its promoter in transgenic plants. *Mol. Gen. Genet.* **236**, 331–340.
- Yamaguchi-Shinozaki, K., and Shinozaki, K.** (1994). A novel *cis*-acting element in an *Arabidopsis* gene is involved in responsiveness to drought, low-temperature, or high-salt stress. *Plant Cell* **6**, 251–264.
- Zeevaart, J.A.D., and Creelman, R.A.** (1988). Metabolism and physiology of abscisic acid. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **39**, 439–473.
- Zhu, J.-K., Hasegawa, P.M., and Bressan, R.A.** (1997). Molecular aspects of osmotic stress in plants. *Crit. Rev. Plant Sci.* **16**, 253–277.